

EFFECT OF DRYING METHOD ON THE ANTIOXIDANT PROPERTIES OF SOME LOCAL SPICES

*¹Zubair, A . B , ²Isah, L . R, ³ Femi, F. A and ⁴ Jiya, M . J

^{1,3,4} Department of Food Science and Technology, Federal University of Technology,

P.M.B. 65 Minna, Niger State Nigeria

²Department of Food and Home Science, Kogi State University Anyigba
Corresponding author: b.zubair@futminna.edu.ng, +2348035889779

Introduction

Spices are a large group of natural ingredients such as dried seeds, fruits, roots, rhizomes, barks, leaves, flowers and other vegetative substances used in a very small quantity as food additives either as color, aroma, flavor or preservative [1]. They are mainly used to improve the palatability/taste and the visual appearance of diets. Most Spices contain phenolic compound that is responsible for their medicinal, antioxidant and preservative properties [2]. The three spices studied include ginger leaf, curry leaf and scent leaf dried using traditional sundried method and oven dried method. The objective of this study is to determine the antioxidant properties of the selected spice.

Materials and Method

The spices (ginger, curry and scent leaf) were purchased from a local market in Minna, Niger State. The spices were sundried 5 h daily for 2 consecutive days and turned over at 1 h interval to achieve uniform drying. After sun drying, the sample was milled to powder form. The second sample was oven dried at a temperature of 50°C for 5 h and turned over periodically to aid uniform drying and milled to powder form. The results obtained from analysis of the antioxidant properties were subjected to one-way ANOVA and the means were separated by Duncan Multiple Range Test using SPSS version 20.

Results and Discussion

The antioxidant properties of the spices is presented in Table 1. Total phenol was in the range of 185 to 290 mg/100g. There was a significant difference across the samples. Total flavonoid ranged from 19 to 27 mg/100. There was significant difference in the values with sample oven dried having the highest value. The decrease in the values of the sundried sample may be due to rapid loss of phenolic compound when the spices were exposed to uncontrolled sun drying temperature [3]. Total flavonoid and phenol have the ability to act as antioxidant, ameliorate inflammation, modulate enzyme activity and regulate gene expression [3].

Table 1 Antioxidant properties of the samples

Samples	Parameters (mg/100g)	
	Total phenol	Total flavonoid
Scent leaf SD	185.0 ^{d1} ± 0.06	25.96 ^b ± 1.15
Scent leaf OD	290.52 ^a ± 4.60	27.56 ^{ab} ± 0.40
Ginger SD	222.58 ^c ± 0.13	19.06 ^d ± 1.00
Ginger OD	249.31 ^b ± 3.46	21.12 ^c ± 0.09
Curry leaf SD	247.65 ^b ± 0.16	27.96 ^a ± 0.23
Curry leaf OD	221.97 ^c ± 1.57	19.85 ^d ± 0.35

Values are means ± standard deviation of duplicate determination. Means in the same column followed by different superscripts are significantly different (p<0.05).

Key:

SD = Sun dried samples

OD = Oven dried samples

Conclusion

The result shows that sun drying and oven drying methods are effective in the processing of spices as well as affect the functionality of the spices in food system. Oven dried samples were found to exhibit a better properties in the parameter investigated. Therefore, the use of oven drying method is recommended in the drying of spices due to its control temperature of drying to minimize the destruction of the antioxidant properties of the spices and also to reduce incidence of contamination of the spices when exposed to open environment for sun drying.

References

1. Birt, D. A. (2006). Phytochemicals and cancer prevention: from epidemiology to mechanism of action. *Journal of the American Dietetic Association* 106: 20 – 24.
2. Burt, S. (2004). Essential oils: Their antibacterial properties and potential applications in foods – A review. *International Journal of Food Microbiology* 94(3): 223–253.
3. Mueller-Harvey, I. (2001). Analysis of hydrolysable tannins. *Anim, Feed Sci, Technol*, 9, 3-20 Podsedek, A. (2007). Natural antioxidants and antioxidant capacity of Brassica vegetables: A review. *Food Science and Technology*, 40,

1-11.