

EVALUATION OF NUTRITIONAL COMPOSITION, PHYSICAL AND CHEMICAL CHARACTERISTICS OF SELECTED RAW AND PROCESSED TOMATO BRAND

Esther Udensi*, Rasaan Bolakale Salau and Simon Olonkwoh Salihu

Department of Chemistry, Federal University of Technology, P. M. B. 65, Minna, Niger State, Nigeria
*Corresponding author's email: Success4j@yahoo.com, +2348166144613

ABSTRACT

Tomato (Lycopersicon esculentum) belonging to the solanaceae family of plants. It is one of the most preferred beneficial vegetable consumed in the world. The aim of this study was to evaluate the nutritional, physical and chemical characteristics of selected local and processed tomatoes sold in Minna, Niger State. Three species of fresh local tomatoes and ten brands of processed tomatoes were sampled in triplicates across markets in Minna city. The proximate, vitamins, phenolic composition physical and chemical parameters were carried out using standard techniques. Proportionate application of HNO₃, H₂O₂ and de-ionised water aided the digestion of the samples. Colorimetric techniques were used to analyze for Phosphorus while FES and FAAS analyse other elements. Statistical Analysis of Variance (ANOVA) was carried out with IBM SPSS statistic version 21. The proximate values (%) are in ranges of: Fat (0.29 – 11.31); Protein (6.65 – 20.15); Fibre (0.92 – 15.70) ash (0.08 – 3.18); Carbohydrate (53.07 – 80.53). The physiochemical parameters values are in ranges: pH (2.25–4.83); Conductivity (3.73 – 12.97), Titratable acidity (0.11 – 1.07); Total soluble solids (15.23–40.70). The minerals values (mg/100g) are in ranges of: K (650.10 – 2187.50); Na (21.55–2912.49); Ca (117.26 – 4297.75); Fe (2.09 – 15.15); Mn (0.30 – 1.24); Zn (8.15–20.01); P (24.45 – 42.17). The vitamins and phenolic values are in ranges: Vitamin A (0.47 – 8.20); vitamin C (1.31–18.49); Phenolic (0.041– 0.360). Prominent values of fat and protein were found in raw samples. High values of physicochemical properties are observed among processed tomatoes relative to raw ones. The raw tomatoes have more prominent content of vitamin C, Mn and K. These unique contents, especially the moderate physical and chemical properties, make raw tomatoes to be more relevant in management of malnutrition.

Key words: Raw tomato and processed tomato brands proximate, physicochemical, FES, FAAS

INTRODUCTION

Tomato (*Lycopersicon esculentum*) belonging to the *solanaceae* family of plants (Eke-Ejiofor, 2015). It is one of the most preferred beneficial vegetable consumed in the world. Tomatoes are cultivated widely in home gardens and large farms for fresh consumption and commercial processing (Aditi *et al.*, 2011). It is typically over 90% water and, once they are harvested, begins to undergo higher rates of respiration, resulting in moisture loss, quality deterioration and potential microbial spoilage (Abdullahi *et al.*, 2016).

Tomato has many nutrients with secondary metabolites that are important for human health such as mineral matter, vitamins, antioxidants, phenolic compounds (Demirbas, 2010) besides other components such as dietary fiber and

consumption reduces the risk of certain types of cancer, cardiovascular, osteoporosis and chronic degenerative diseases (Chang *et al.*, 2006; Hernández *et al.*, 2008; Bhowmik *et al.*, 2012). Minerals are involved in many important functions in the body, such as enzymatic reactions, bone mineralization, as well as the protection of cells and lipids in biological membranes. Low intake of minerals leads to deficiencies which could cause impairment of body functions (Melø *et al.*, 2008). They detoxify free radicals which are produced during normal metabolism that affect DNA. Major and minor element contents of tomatoes depend on cultivar, cultivation method, region of cultivation, sampling period and growing conditions (Demirbas, 2010; Hernández *et al.*, 2008).

10
9
8
7
6
5
4
3
2
1

The first part of the document discusses the importance of maintaining accurate records of all transactions. It emphasizes that proper record-keeping is essential for the success of any business and for the protection of the interests of all parties involved.

ACCOUNTING AND BUSINESS

Accounting
Accounting is the process of recording, summarizing, and explaining the transactions and events which in part at least are financial in nature, and which in part at least are of a financial character, and interpreting the results thereof.

The second part of the document discusses the various methods used in accounting to record and summarize transactions. It covers the double-entry system, the use of journals and ledgers, and the preparation of financial statements. It also discusses the importance of internal controls and the role of the auditor.

CONCLUSION

In conclusion, accounting is a vital part of any business. It provides the information needed to make informed decisions and to ensure the success of the organization. Proper accounting practices are essential for the long-term survival and growth of any business.

The document concludes by emphasizing the importance of accuracy and integrity in accounting. It states that accountants have a duty to provide reliable and unbiased information to their clients and the public.

Accounting is a dynamic field that evolves with the needs of the business world. It is essential for the success of any business and for the protection of the interests of all parties involved.

The second part of the document discusses the various methods used in accounting to record and summarize transactions. It covers the double-entry system, the use of journals and ledgers, and the preparation of financial statements. It also discusses the importance of internal controls and the role of the auditor.

Accounting

Accounting is the process of recording, summarizing, and explaining the transactions and events which in part at least are financial in nature, and which in part at least are of a financial character, and interpreting the results thereof.

Accounting

The second part of the document discusses the various methods used in accounting to record and summarize transactions. It covers the double-entry system, the use of journals and ledgers, and the preparation of financial statements. It also discusses the importance of internal controls and the role of the auditor.

ACCOUNTING AND BUSINESS

The document discusses the importance of maintaining accurate records of all transactions. It emphasizes that proper record-keeping is essential for the success of any business and for the protection of the interests of all parties involved.

The document concludes by emphasizing the importance of accuracy and integrity in accounting. It states that accountants have a duty to provide reliable and unbiased information to their clients and the public. Accounting is a dynamic field that evolves with the needs of the business world. It is essential for the success of any business and for the protection of the interests of all parties involved.

brand 7 the highest Titratable acidity (0.11–1.07) did not exceed the maximum acidity (7 %) recommended by CAC (2012). Total soluble solids (15.23–40.70) with brand 6 having the least and brand 7 the highest. The total soluble solid (15.23–40.70) of the samples were higher

than the standard level of 20 to 22 % required by CAC (2011). The higher total soluble solid might be due to lower moisture content as reported by Eke-Ejiofor (2015).

Table 1; Physiochemical parameters

S/N	Sample	pH	Conductivity	Titratable Acids (%)	Total Soluble Solids
1	KI	4.79±0.01 ^e	3.73±0.01 ^a	0.41±0.01 ^a	28.42±0.01 ^k
2	MG	4.85±0.01 ^h	5.72±0.01 ^d	0.77±0.01 ^b	18.17±0.02 ^h
3	ZU	4.79±0.00 ^f	6.48±0.00 ^f	0.41±0.01 ^a	15.81±0.01 ^d
4	B1	4.78±0.01 ^e	8.87±0.01 ⁱ	0.18±0.00 ^a	17.53±0.01 ^g
5	B2	4.68±0.00 ^c	6.72±0.00 ^g	0.31±0.01 ^a	18.65±0.01 ⁱ
6	B3	4.86±0.01 ^h	8.34±0.00 ^h	1.02±0.01 ^b	15.21±0.01 ^b
7	B4	2.25±0.00 ^a	9.87±0.00 ^k	1.02±0.01 ^b	17.27±0.01 ^e
8	B5	4.72±0.00 ^d	12.97±0.00 ^m	1.04±0.00 ^b	15.23±0.03 ^a
9	B6	4.56±0.00 ^b	6.36±0.00 ^e	1.04±0.00 ^b	15.61±0.01 ^c
10	B7	4.86±0.01 ^h	5.25±0.00 ^c	1.07±0.00 ^b	40.61±0.01 ^l
11	B8	4.73±0.01 ^d	9.03±0.01 ^j	1.02±0.01 ^b	18.15±0.05 ^h
12	B9	4.88±0.01 ⁱ	9.92±0.00 ^l	0.11±0.01 ^a	19.01±0.01 ^j
13	B10	4.83±0.00 ^g	3.93±0.00 ^b	0.44±0.03 ^a	17.41±0.01 ^f

Values are reported as mean ± standard error of means. Values with the same letter on the column are not significant while values on the same column with different alphabetic superscript are significant at $p \leq 0.05$ DMRT test.

Key: KI = Kano-India variety, MG = Minna-Gwari variety, ZU= Zaria UTC variety, B1–B10 are processed tomato brands 1-10

Table 2 showed proximate composition (%). The proximate values are in ranges: Moisture (0.74–11.84) with brand 7 having the least and brand 6 the highest. The moisture contents values (0.74 – 11.84) with brand 5 having highest moisture. These were low compared to the values (69.00 – 84.85 %) reported by Eke-Ejiofor (2015). These decreases in moisture of the processed tomatoes increase the shelf life, hence beneficial to the consumers (Joel *et al.*, 2020). Crude fat (0.29–11.31) with brand 3 having the least fat and Minna-Gwari species the highest (11.31 %). The fat content of the fresh tomato (Minna-Gwari species) is higher than all the processed brands and as well as its raw counterparts. The reason might be due to different geographical location and different in processing methods. Crude protein (6.66–20.15) with brand 1 having the least and Minna-Gwari species the highest. The crude protein was higher in some of the raw than all the processed brands. The difference in crude protein contents could be attributed to species differences as well as

differences in the processing conditions of the and high-water content of the raw. The crude fibre pastes. The higher protein content of the raw content of high-water content of the raw tomato, this could be as a result of higher variance with the reported higher result of high-water content of the raw tomato, similar observation. Crude fibre (53.07-80.53) with brand 8 the highest; Ash (0.08-3.18) with brand 10 having the least and brand 3 the highest. Carbohydrate percentage content in the raw content in both raw and processed brands differed significantly, however, one of the processed tomatoes have higher ash than others. This might be as a result of addition of salt to the processed tomatoes.

Table 2: Proximate Composition (%)

Samples	Moisture	Crude Fat	Crude Protein	Crude Fibre	Ash	Carbohydrate
---------	----------	-----------	---------------	-------------	-----	--------------

K1	4.77±0.01 ^b	9.08±0.00 ^f	19.26±0.01 ^k	11.57±0.01 ⁱ	2.28±0.00 ^a	53.07±0.00 ^a
MG	5.68±0.01 ^d	11.31±0.00 ^m	20.15±0.01 ⁱ	3.91±0.01 ^e	2.26±0.05 ⁱ	56.68±0.00 ^c
ZU	5.78±0.01 ^e	9.81±0.00 ^f	13.35±0.05 ⁱ	1.03±0.01 ^b	1.43±0.00 ^f	68.58±0.00 ⁱ
B1	7.93±0.01 ^j	9.11±0.00 ^g	6.66±0.01 ^a	2.92±0.01 ^d	2.01±0.00 ^f	71.39±0.00 ^k
B2	8.18±0.01 ^k	9.67±0.00 ^h	11.39±0.01 ^s	11.68±0.01 ^j	1.54±0.00 ^g	68.96±0.00 ^l
B3	5.47±0.01 ^c	0.29±0.00 ^a	10.07±0.01 ^d	2.60±0.00 ^e	1.07±0.00 ^d	80.53±0.00 ^m
B4	7.03±0.01 ^h	8.57±0.00 ^d	9.19±0.01 ^c	6.75±0.01 ^h	1.46±0.00 ^f	66.91±0.00 ^h
B5	8.37±0.01 ⁱ	9.73±0.00 ^k	10.07±0.01 ^d	11.94±0.01 ^k	3.18±0.00 ^k	56.73±0.00 ^d
B6	11.84±0.01 ^m	6.82±0.00 ^b	8.32±0.01 ^b	11.94±0.01 ^k	0.86±0.00 ^c	60.25±0.00 ^e
B7	0.74±0.01 ^a	8.58±0.00 ^c	10.51±0.01 ^e	3.94±0.01 ^f	0.52±0.00 ^b	75.74±0.01 ^j
B8	7.52±0.01 ⁱ	9.73±0.00 ^l	10.94±0.01 ^f	15.61±0.01 ^l	1.69±0.00 ^h	54.45±0.00 ^b
B9	5.99±0.01 ^f	9.19±0.00 ^h	12.26±0.01 ^h	4.23±0.01 ^s	1.39±0.01 ^e	66.98±0.00 ^g
B10	6.52±0.01 ^s	7.76±0.00 ^c	17.93±0.01 ^j	0.92±0.01 ^a	0.08±0.00 ^a	66.79±0.01 ^f

Values are reported as mean ± standard error of means. Values with the same letter on the column are not significant while values on the same column with different alphabetic superscript are significant at p≤0.05 DMRT test

Key: Kano = Kano-India variety, Minna = Minna-Gwari variety, Zaria = Zaria UTC variety

Phenolic and vitamins (antioxidants) are

shown in Table 3.

the highest value of vitamin A, C and Phenolic were found in Brand 9, Kano-India species and and 7 respectively. Vitamins A, C and Phenolic important quality parameters used in assessing tomato paste. They act as antioxidant, preventing the oxidation of some fatty acid component and play some important vital role in the body metabolism.

Table 3: Antioxidants Analysis (mg/100 g)

Samples	Vitamin A	Vitamin C	Phenolic content
KI	0.47±0.00 ^a	18.49±0.01 ^j	0.25±0.01 ^f
MG	1.28±0.02 ^d	10.60±0.10 ^h	0.11±0.00 ^d
ZU	1.93±0.00 ^b	15.89±0.01 ⁱ	0.16±0.00 ^c
B1	1.89±0.00 ^f	2.61±0.01 ^b	0.20±0.00 ^d
B2	3.57±0.00 ⁱ	1.31±0.01 ^a	0.46±0.00 ^j
B3	1.06±0.00 ^c	7.81±0.01 ^f	0.04±0.00 ^a
B4	4.28±0.00 ^j	7.91±0.01 ^f	0.24±0.00 ^f
B5	5.62±0.00 ^k	8.51±0.01 ^g	0.12±0.00 ^b
B6	0.76±0.00 ^b	5.31±0.01 ^e	0.22±0.02 ^e
B7	1.81±0.00 ^e	4.62±0.02 ^d	0.36±0.00 ⁱ
B8	6.55±0.00 ^l	5.35±0.05 ^e	0.32±0.00 ^g
B9	8.20±0.00 ^m	3.81±0.00 ^c	0.34±0.00 ^h
B10	2.32±0.00 ^h	2.65±0.05 ^b	0.32±0.00 ^g

Values are reported as mean ± standard error of means. Values with the same letter on the column are not significant while values on the same column with different alphabetic superscript are significant at $p \leq 0.05$ DMRT test.

Key: KI = Kano-India variety, MG = Minna-Gwari variety, ZU= Zaria UTC variety, zB1–B10 are processed tomato brands 1-10

Table 3 showed the Mineral composition (%). The minerals values are in range: Potassium (650.10–2187.50) brand 9 having the least and Kano-India species the highest. Potassium (K) which has numerous functions in the biochemical and physiochemical functions of the body according to report of Abdullahi *et al.*, (2016). They found the raw kano variety to have the highest value of K. The concentrations of sodium were higher in processed brands. This was attributable to the addition of table salt during the course of processing to improve preservation. This makes it inimical to hypertensive patients if recommend. With regard to Iron concentration, the processed brands tomato was found to be highest. Such a difference might arise due to possible deposition of iron from the iron plates used in drying of the tomato samples and differences in geographical location (Abdullahi *et al.*, 2016). Calcium, Zinc

and Phosphorus were significantly different in respect to the fresh species and processed tomato brands; however, their peak values were found with the brand 2, the processed tomato. Both are required in our dietary intake, as calcium and zinc support bone mineral density (Goodson, 2018) and phosphorus help to maintain a regular heartbeat and facilitate nerve conduction among others (Madell, 2020). There was also an assertion that foods that high in calcium are also high in phosphorous (Madell, 2020). Manganese (0.30–1.24) brand1 having the least and Minna-Gwari the highest; Manganese (0.30–1.24) is considered an essential nutrient and can be found especially, in vegetables. The highest value is found in fresh tomato (Minna-Gwari species). Manganese may play a positive role in bone health by working in concert with other vitamins and minerals to improve bone mineral density (Goodson, 2018)

Table 4: Mineral compositions

Samples	K (mg/100g x10 ³)	Na (mg/100g x10 ³)	Ca (mg/100g x10 ³)	Fe (mg/100g)	Mn (mg/100g)	Zn (mg/100g)	Phosphorus (P) (mg/100g)
KI	2.19±0.01 ^k	0.05±0.03 ^c	2.76±0.22 ^l	2.39±0.00 ^b	0.80±0.00 ^j	15.43±0.00 ^j	37.47±0.00 ⁱ
MG	1.31±0.01 ^f	0.04±0.01 ^b	1.29±0.050 ^h	3.93±0.00 ^e	1.24±0.01 ^l	16.30±0.00 ^k	36.65±0.00 ^h
ZU	1.76±0.01 ⁱ	0.02±0.05 ^a	1.03±0.05 ^s	3.06±0.00 ^c	0.72±0.00 ⁱ	9.56±0.01 ^e	39.47±0.00 ^j
B1	0.76±0.01 ^c	1.14±0.11 ^k	1.66±0.10 ^j	2.09±0.01 ^a	0.21±0.00 ^a	8.53±0.00 ^c	40.87±0.00 ^j
B2	1.56±0.00 ^s	0.25±0.45 ^e	4.21±0.26 ^m	3.65±0.00 ^d	0.51±0.00 ^e	20.06±0.01 ^m	42.17±0.00 ^m
B3	1.56±0.01 ^s	0.61±0.01 ⁱ	1.56±0.01 ⁱ	7.87±0.00 ^j	0.56±0.00 ^f	14.01±0.01 ^h	24.76±0.00 ^b
B4	0.76±0.05 ^c	0.71±0.05 ^j	0.17±0.01 ^b	4.73±0.24 ^s	0.43±0.00 ^c	8.55±0.00 ^d	32.97±0.00 ^e
B5	1.93±0.01 ^j	2.91±0.01 ^m	0.44±0.01 ^c	7.18±0.00 ^h	1.02±0.00 ^k	13.65±0.01 ^s	39.76±0.00 ^k
B6	0.93±0.05 ^d	0.59±0.05 ^h	0.12±0.01 ^a	8.89±0.00 ^k	0.47±0.00 ^d	8.31±0.01 ^b	33.93±0.00 ^s
B7	1.21±0.01 ^e	0.22±0.01 ^d	0.59±0.02 ^e	15.15±0.00 ^m	0.69±0.00 ^h	14.33±0.01 ⁱ	32.02±0.00 ^d
B8	1.59±0.05 ^h	2.84±0.02 ^l	0.51±0.05 ^f	8.21±0.00 ^j	0.61±0.00 ^s	10.41±0.00 ^f	28.28±0.00 ^c
B9	0.65±0.10 ^a	0.27±0.05 ^s	0.46±0.04 ^d	4.44±0.00 ^f	0.42±0.00 ^b	8.15±0.00 ^a	24.45±0.00 ^a
B10	0.68±0.01 ^b	0.26±0.01 ^f	1.78±0.02 ^k	14.34±0.00 ^l	0.69±0.00 ^h	17.51±0.01 ^l	33.81±0.00 ^f

Values are reported as mean ± standard error of means. Values with the same letter on the column are not significant while values on the same column with different alphabetic superscript are significant at p≤0.05 DMRT test.

Key: KI = Kano-India variety, MG = Minna-Gwari variety, ZU= Zaria UTC variety, B1–B10 are processed tomato brands 1-10

CONCLUSION

From the study, both raw tomatoes and processed brands were source of nutrients. Carefully taking according to the body requirement will benefit humanities. The processed tomatoes were found to have significant concentration in some of essential nutrients than the raw species in some cases. However, prominent values of fat and protein were found in raw samples. High values of physicochemical properties are observed among

processed tomatoes relative to local ones. The raw tomatoes have more prominent content of vitamin C, Mn and K. These unique contents, especially the moderate physical and chemical properties, make raw tomatoes to be more relevant in management of malnutrition.

REFERENCES

Abdullahi, I.I., Abdullahi, N., Abdu, M.A and Ibrahim, S.A. (2016). Proximate, mineral and vitamin analysis of fresh and canned tomato. Journal of

- Aditi Gupta, Kawatra, A and Sehgal, S. (2011). Physicalchemical properties and nutritional evaluation of newly developed tomato genotypes. *African Journal of Food Science and Technology*, 2(7), 167-172.
- Bhowmik, D., Kumar, K. S., Paswan, S. and Srivastava, S. (2012). Tomato-a natural medicine and its health benefits. *Journal of Pharmacognosy and Phytochemistry*, 1(1), pp. 33-43.
- CAC (Codex Alimentarius Commission) (2011). Technical specification for tomato paste. CODEX STAN 57, 1-6.
- Chang, C. H., Lin, H. Y., Chang, C. Y. And Liu, Y. C. (2006). Comparisons on the antioxidant properties of fresh, freeze-dried and hot-air-dried tomatoes. *Journal of Food Engineering*, 77(3), 478-485.
- Demirbas, A. (2010). Oil, micronutrient and heavy metalcontents of tomatoes. *Food Chemistry*, 118, 504-507.
- Eke-Ejiofor J. (2015). Comparative evaluation of lycopene content and some chemical properties of commonly consumed brands of tomato paste in port -harcourt, South-South, Nigeria. *Journal of Food and Nutrition Sciences*, 3(2), 35-3.
- Goodson, A. (2018). 10 Evidence-Based Benefits of Manganese. <https://www.healthline.com/nutrition/manganese-benefits>. Accessed on 15th April, 2020.
- Hernández, M.S., Rodríguez, E.R. & Díaz, C.R. (2008). Chemical composition of tomato (*Lycopers icon esculentum*) from Tenerife, the Canary Islands. *Food Chemistry*, 10(6), 1046-1055.
- Joel, N., Anselm, U.O. and Oluwafunmike, O. (2020). Comparative evaluation of the qualities of some selected tomato-paste brands sold in kano market. *Journal of Food. Stability*, 3 (1), 1-11.
- Katırcı, N., Isık, N., Güpür, C., Guler, O.H., Gursoy, O. and Yilmaz, Y. (2018). Differences in antioxidant activity, total phenolic and flavonoid contents of commercial and homemade tomato pastes, *Journal of the Saudi Society of Agricultural Sciences*, 30(40), 1-7
- Madell, R. (2020). Phosphorus in Your Diet. <https://www.healthline.com/phosphorus-in-diet/#food-source>. Accessed on 15th April, 2020.
- Melø, R., Gellein, K., Evje, L. And Syversen, T. (2008). Minerals and trace elements in commercial infant food. *Food and chemical toxicology*, 46 (10), 3339-3342.