



STUDIES ON THE IMPACT OF GAS-FLARING ON THE QUALITY OF RAIN WATER, GROUNDWATER AND SURFACE WATER IN PARTS OF EASTERN NIGER DELTA, NIGERIA

A. N. Amadi*, P. I. Olasehinde*, Y. B. Alkali* and J. O. Jacob**

*Department of Geology, Federal University of Technology, Minna

**Department of Chemistry, Federal University of Technology, Minna

Email address: geonma76@gmail.com; amadi.nwanosike@futminna.edu.ng

ABSTRACT

The environmental and health impact associated with over 50 years of oil exploration and exploitation in Eastern Niger Delta region of Nigeria is of great concern. Flaring is the controlled burning of the natural gas associated with oil production. This study has clearly shown that gas-flaring constitutes a major source of water pollution in the oil producing region of Eastern Niger Delta. The results of the laboratory analysis of the water samples revealed that the water sources in the area have been negatively impacted acid-rain and NO_2 , SO_2 and CO_2 from the burning gas. The sulphate, bicarbonate and nitrate content of rain water samples within the radius of 20 km from the gas-flaring station are high but the concentration decreases away from the gas-flaring stations. It was ascertained that the acidity and hence the nitrate, sulphate and bicarbonate content of rain water is a function of the frequency and duration of the rainfall as well as the direction of the prevailing wind prior to the rainfall and it gives rise to the formation of acid-rain. Corrosion of buildings, the dominance of respiratory problems and skin diseases are some of the signatures on the host communities. However, there may be a serious long term effect on the water resources in region in terms of quality if gas-flaring in the area does not stop.

Keywords: Impact of Gas-Flaring, Water Quality, Eastern Niger Delta, Nigeria

INTRODUCTION

Globally, gas flaring is one of the hottest debated environmental issues currently and in Nigerian oil producing Niger Delta in particular. Flaring is the controlled burning of the gas associated with oil production. Flaring is a means of eliminating gas when the volume is insufficient in terms of recovery and collecting it would be uneconomical. Gas flaring is a serious contributor to acid-rain, the impacts of which are already being felt in the Niger Delta region in terms of vegetation damage (Plate 1), corrosion and caving-in of roofing sheets (Plate 2) and death of aquatic lives (Plate 3). According to [1], about 115 billion cubic meters of gas are flared every year world-wide. Similarly, [2],

ascertained that gas flaring in Niger Delta had contributed more greenhouse gases to the atmosphere more than all other sources in Sub-Saharan Africa Combined. Oil production in Niger Delta began over 50 years ago and so did the practice of flaring associated gas. In developed countries, the associated gas is re-injected into the ground while in Nigeria it is burnt off into the atmosphere without considering the environmental and health impact. Despite regulations introduced more than 20 years ago to outlaw the practice, most associated gas are flared by the oil companies, causing local pollution and contributing to climate change [3].



Plate 1: Degradation of the Vegetation in the area due to acid rain
 Plate 2: A forest of site trees with various degree of rusting/leaving in



Plate 3: Death of aquatic animals in the rivers in the area due to acid-rain caused by gas flaring

According to an interview with a village head, from one of the local communities living around the gas flaring station, the unpleasant smell, roaring noise and intense heat emanating from the gas flaring are dehumanizing as many of them are now suffering from skin diseases, cancer, ear problem, respiratory problems such as asthma and bronchitis. He confirmed that these diseases were new to them and are possible caused by continuous flaring of gas in the area. The impact of gas flaring on the environment and health of host communities in Niger Delta, Nigeria is of great concern. In view of the economic activities domiciled in the region, it becomes imperative to undertake a comprehensive study of the effects of gas flaring on the water resources of the area. This study was carried out in order to assess the impact of the gas flaring on the quality of rain water, surface water and groundwater in the vicinity of two representative gas flaring stations in Eastern Niger Delta within a radius of 40km centered at each station.

MATERIALS AND METHODS

Physiography, Geology and Hydrogeology of the area
 The area is a low land and characterized by two distinct seasons: a dry season (November to March) and rainy season (April to October). The study area is underlain by the coastal plain sand of Miocene age (Fig.1). The study area is underlain by unconfined aquifer of regional extent. Recharge into the unconfined aquifer is by direct infiltration of rainwater. The formation is made up of friable, fine to coarse grained sands with minor clay intercalations [5]. It consists of water-bearing, porous and highly permeable continental sands. Records from drilled boreholes indicate shallow groundwater table while pumping test analyses show high hydraulic characteristics in terms of yield, transmissivity and storativity. Petrographic study on several thin sections [5] shows that quartz makes up more than 96% of all the sand grains. The general thickness of the coastal plain sands is variable and ranged from 180 m to 2000 m [6]. The annual groundwater recharge ranged between 20% and 30% of the annual rainfall of about 2500 mm [7].



Sampling and Laboratory Analysis

Altogether 38 samples comprising of 20 groundwater samples, 10 rain water samples and 8 surface water samples were collected and analyzed. All the samples were preserved by refrigeration and analyzed within 24 hours of collection. The analyses were carried out in accordance with [8], standard for analyzing water and waste water. The physical parameters pH and conductivity were determined on the field using a Gallenkamp pH meter and Hach conductivity meter WPA 400 digital model respectively. The coordinates of each samples location was taken using the global positioning system and the data was used to generate the digital terrain model of the area overlap with the contour map and groundwater flow direction (Fig. 2).




Fig. 1: Geology map of Eastern Niger Delta showing the study area




Fig. 2: Digital terrain model of the study area overlapped with contour and flow lines

RESULTS AND DISCUSSION

The mean concentrations of some of the parameters analyzed are shown in figures 3 and 4 while the scatter plot and correlation graph of the different water sources analyzed are illustrated in figures 5 and 6 respectively. The effect of gas flaring on the quality of the water sources was studied using hydrochemical data from water samples in the vicinity of the gas flaring stations. The concentration of the major anions (bicarbonate, sulphate and nitrate), temperature, conductivity, heavy metal content (HM) and total hydrocarbon content (TH) increase appreciable in the vicinity of the gas flaring stations and consistently decreased away from the gas flaring points (Figures 3 and 4). Their enrichment in the water sources may be attributed to the gas flaring activity in the area [9, 10]. The rain water samples show concentration and variation trends similar to those of surface water and groundwater (Fig.3). The concentrations of pH, moisture content (MC) and bacteria count (BC) were lowest at vicinity of the gas flaring stations but increases away from the gas flaring points. The increased acidity of the rain water in the gas flaring stations may be responsible for the low pH (5.2) as control samples taken far away the gas flaring stations have higher pH (6.7). Similarly, the low concentration of

MC and BC can be attributed to the intense heat emanating from the gas flaring station. Such heat (elevated temperature) may be capable of drying up the soil moisture and kill the bacteria in the soil/water as bacteria cannot withstand high temperature (Fig. 4).

The similarity in distribution pattern of the geochemical constituents from the different water sources (Fig. 5) in the vicinity of the gas flaring stations is a clear indication that the enrichment of these ions are associated with the gas flaring activity. A comparison of the distribution pattern (Fig. 6) and the groundwater flow direction (Fig. 2) revealed a striking similarity. The gas flaring station overlies a groundwater recharge zone and the direction of the decreasing ionic concentration is parallel to the direction of groundwater flow away to the discharge zone. This relationship is an indication that the movements of contaminants (TH, HM, SO_4^{2-} , HCO_3^- and NO_3^-) ions are likely assisted by advection, which is the contaminant transport by the bulk volume of the migrating groundwater. Surface water sampled at the discharge area of the groundwater flow from the gas flaring stations also showed evidence of geochemical enrichment. The results are consistent with the




Fig. 3: Graph of water sources and their concentrations




Fig. 4: Graph of concentrations versus distance (km) to gas flaring station

Fig. 5: Scatter plot of water sources versus concentrations in mg/l







Fig.6: Correlation analyses of water sources versus concentrations in mg/l



CONCLUSION AND RECOMMENDATIONS

This study has clearly demonstrated that gas flaring constitutes a major source of water pollution in the oil producing area of Eastern Niger Delta, Nigeria. The signatures are felt in heavily on the water sources in the area. The dispersion of the enriched contaminant is heaviest on the direction of the groundwater flow but reduces on the opposite direction. The deterioration of the water resources in region will continue except an end is put to gas flaring in the region. Evaluation of the impact of gas flaring on plants, soil and air should be undertaken in the area in order to have comprehensive baseline information on the impact of gas flaring on the environment.

REFERENCE

- [1]. Stan, S., (2002). Gas flaring: A mounting environmental concern in western Canada. *Saskatchewan Business Magazine*, June/July-2002, pp 21.
- [2]. World Bank, (2004). Gas flaring in Nigeria. *Friends of the Earth Media Briefing*, October 2004, pp 1. www.foe.co.uk.
- [3]. Tell Magazine, (2008). The 50 years of oil in Nigerian. *Tell Magazine*, 18th February, 2008, 9-109.
- [4]. Daily Trust Newspaper, (2008). Nigeria: Gas flaring in the Niger Delta and its health hazards. 8th March, 2003, pp 1-2.
- [5]. Onyeagocha, A. C., (1980). Petrography and depositional environment of the Benin Formation, Nigeria. *Journal of Mining and Geology*, 17, 147-151.
- [6]. Avbovbo, A. A., (1978). Tertiary lithostratigraphy of Niger Delta: *Bulletin of American Association of Petroleum Geology*, 62, 297-306.
- [7]. Uma K. O. & Egboka, B. C. E., (1987). Water resource of Owerri and its environs, Imo State, Nigeria. *Journal of Mining and Geology*, 22(1-2), 57-64.
- [8]. APHA, (1995). Standards methods for the examination of water and wastewater. 19th Edition *American Water Works Association*, Washington DC.
- [9]. Uma, K. O., (1989). Assessment of the impact of gas-flaring on the quality of rain water, surface water and groundwater in parts of the oil-producing region of Nigeria. *Water Resources*, 1(2), 200-204.
- [10]. Amadi, A. N., Nwankwoala, H. O., Eze, C. I., Alkali, Y. B. and Waziri, S. H., (2012). A review of waste management techniques in parts of Niger Delta, Nigeria. *Centre for Human Settlement and Urban Development Journal*, 3(1), 98-108.