

Effect of Rates and Duration of *Gliricidia* Biochar and Poultry Manure on Chemical Properties of Soil in Minna, North Central, Nigeria

¹Tsado, P.A., ²Igwe, C.A., ¹Lawal, B.A., ¹Osunde, A. O., ¹Eze, P.C. and ²C.C. Okolo

¹Department of Soil Science and Land Management, Federal University of Technology, Minna, Nigeria

²Department of Soil Science and Land Management, University of Nigeria, Nsukka, Nigeria

*Corresponding authors' e-mail address: tsadophilips@yahoo.com / p.tsado@futminna.edu.ng

Abstract

Pot experiment was carried out in the screen house of School of Agriculture and Agricultural Technology, Federal University of Technology, Minna. The objective of the study was to investigate the effectiveness of different rates and duration of *Gliricidia Biochar* (GB) and Poultry Manure (PM) on some chemical properties of soil sampled every two months in Minna. The treatments consisted of nine rates of GB and PM (control, 10, 20, 30, 40 GB and 10, 20, 30, 40 PM t ha⁻¹) and three duration of sampled at 2, 4 and 6 months and analyzed for their chemical properties. The experiment was a factorial fitted to Complete Randomized Design (CRD) with three replications. Laboratory analysis of soil chemical properties was conducted. All data collected were subjected to ANOVA at 5% level of significant and the means were separated using Duncan Multiple Range Test (DMRT). Irrespective of the rates of application of GB and PM, there was a significant increase in pH, organic carbon, effective cation exchange capacity, total nitrogen and available phosphorus over the control. Also, the results showed a significant increase of the above parameters with the durations of these amendments. No significant interaction existed between the rates and duration of GB and PM application on the above parameters recorded during the experiment.

Keywords: : *Gliricidia* Biochar, Poultry Manure, Chemical properties, Rates of application

EFFET DES TAUX ET DE LA DURÉE DU GLIRICIDIA BIOCHAR ET DU FUMIER DE VOLAILLE SUR LES PROPRIÉTÉS CHIMIQUES DU SOL À MINNA, NORD CENTRAL, NIGERIA

Abstrait

Résumé

L'expérience en pot a été réalisée dans le crible de l'École d'agriculture et de technologie agricole, Université fédérale de technologie, Minna. L'objectif de l'étude était d'étudier l'efficacité de différents taux et durées de *Gliricidia* Biochar (GB) et de fumier de volaille (PM) sur certaines propriétés chimiques du sol échantillonné tous les deux mois à Minna. Les traitements étaient regroupés en neuf taux de GB et PM (contrôle, 10, 20, 30, 40 GB et 10, 20, 30, 40 PM t ha⁻¹) et trois durées d'échantillonnage à 2, 4 et 6 mois et analysées pour leurs propriétés chimiques. Le dispositif expérimental était un factoriel ajusté à la conception aléatoire complète (CRD) avec trois répétitions.

Une analyse en laboratoire des propriétés chimiques du sol a été effectuée. Toutes les données collectées ont été soumises à une analyse ANOVA au seuil de significativité 5% pour la comparaison des moyennes suivies du test de comparaison multiple Duncan Multiple Range (DMRT) pour la comparaison des différents traitements. Indépendamment des taux d'application de GB et de PM, il y avait une augmentation significative du pH, du carbone organique, de la capacité d'échange cationique effective, de l'azote total et du phosphore disponible par rapport au témoin. De plus, les résultats ont montré une augmentation significative des paramètres ci-dessus avec la durée de ces modifications. Aucune interaction significative n'existait entre les taux et la durée de l'application de GB et de PM sur les paramètres ci-dessus enregistrés pendant l'expérience.

recommandé qu'un atelier soit organisé pour les agriculteurs par les agences agricoles compétentes en vue de promouvoir un écosystème de qualité et d'accroître la participation à une agriculture respectueuse de l'environnement.

Mots clés: Gliricidia Biochar, Fumier de volaille, Propriétés chimiques, Taux d'application

Introduction

Arable soils of the tropics are degraded quickly physically, chemically and biologically as a result of continuous cropping with persistent use of inorganic fertilizers, increase soil acidity and physical degradation while crop yield may fall. Some of the fertilizers aggravate nutrient imbalance due to lack of soil testing programme (Ojeniyi, 2002). These soils are low in organic matter, basic cations, available phosphorus and total nitrogen. This low level of organic matter has made the soils susceptible to major chemical, physical and biological limitations which reduce crop yields (Jones and Wild, 1975).

The use of organic materials as a source of nutrients is a well established practice among the small farming sector in many countries including Nigeria.

Biochar is a recent organic material used as a soil amendment to improve soil nutrients for crop production. They are produced during a process known as pyrolysis from the thermo-conversion of biomass under little or no oxygen supply for use in soil as an amendment. When biochar is applied to the soil, it increases the soil chemical properties. This is observed in the amount of N retention in the soil, increased organic carbon content, pH cations exchange capacity, decreased exchange acidity, sulphur and zinc (Cheng, *et al.*, 2008).

They are produced from a wide variety of feed stocks. *Gliricidia* feedstock for the production of biochar was from *Gliricidia sepium*, a N fixing tree species that grow wild on fallow lands in the southern and middle belt agro – ecological zones of Nigeria and which does not appear to have any other higher net resources value than to be converted to biochar (Fagbenro, *et al.*, (2016).

Poultry manure is also a major organic material that is used by a wide range of the farmers as a soil amendment that provides high nutrient content (N P K) and other essential nutrients (Farhad, *et al.*, 2009). When compared with chemical fertilizers, it also add organic matter to the soil to improve soil structure, aeration, soil

moisture holding capacity and water infiltration rates (Nguyen, 2010). Increase in soil nitrogen and phosphorus was reported by (Thomas and Aluko, 2016) as the rates of application of poultry, goat and dairy cow manures increases.

Therefore, the objective of the study was to investigate the effectiveness of different rates and duration of *Gliricidia* Biochar (GB) and Poultry Manure (PM) on some chemical properties of soil in Minna.

Results and Methods

A pot trial was conducted at the screenhouse of the Department of Soil Science and Land Management, Federal University of Technology, Minna (Latitude 9053'N and Longitude 6045'E). Minna falls under the Southern Guinea Savannah agro – ecological zone of Nigeria. The climate of the area is sub – humid tropical with Mean Annual Rainfall of about 1284 mm with the Mean Maximum Temperature of around 35.5 °C (Ojanuga, 2006). The soil used in the pot trial was collected from the Teaching and Research Farm of Federal University of Technology, Minna at the depth of 0 – 20cm using soil auger. The soils were air dried, crushed and sieved through 2mm mesh and the pre-treatment analysis carried out (Table 1).

The treatment consisted of: (i) control (no amendment), four rates of *Gliricidia* biochar (GB), (10, 20, 30 and 40 t ha⁻¹) and four rates of Poultry manure (PM) (10, 20, 30 and 40 t ha⁻¹). (ii). Three sampling duration (2, 4 and 6 months after the establishment of the experiment). The experiment was a 9 x 3 factorial fitted to Completely Randomized Design (CRD) with three replications.

Two kilogram of soil samples thoroughly mixed with the amendments was weighed into eighty one poly pots (drainage holes were earlier made on each of the poly pot but covered with lining materials before filling with the soil). And on weekly basis, the soil columns were wetted with 700 ml of water. Five hundred

grams soil sample were collected at interval of 2 months after the establishment of the experiments.

Soil samples were air-dried and gently crushed to pass through 2 mm and 0.5 mm sieve. pH values of the samples were determined in 1.0N CaCl₂ solution using a soil - solution ratio of 1:2.5 (McLean, 1982). Organic carbon was determined by the Walkley-Black wet oxidation method (Allison, 1965). Exchangeable basic cations were extracted with neutral 1N NH₄OAc with potassium (K) and sodium (Na) determined by flame photometry and calcium (Ca) and magnesium (Mg) by atomic absorption spectrophotometry. Exchange acidity was determined by shaking the samples with 1.0 M KCl and titrating them with 0.1 M NaOH. Available P was determined by the Bray P1 method (Bray and Kurtz, 1945). Effective cations exchange capacity was obtained by summation of the exchangeable basic cations. Total nitrogen (TN) was determined by the micro Kjeldahl method.

The data generated was subjected to analysis of variance (ANOVA) at 5 % level of significance, while the means were separated using Duncan Multiple Range Test (DMRT).

Results and Discussions

The properties of the soil before the application of the treatment (Table 1) shows that the soil was sandy loam in texture with 668 gkg⁻¹, 160gkg⁻¹ and 172gkg⁻¹ of sand, silt and clay respectively. The soil is moderately acidic in reaction and low in organic carbon (8.24 gkg⁻¹), total nitrogen (0.45gkg⁻¹) and available phosphorus (8.89mgkg⁻¹). These values and the others in Table 1 imply that the soil used for the trial was low in fertility. Jones and Wild (1975) reported that low to medium organic carbon rate for savanna soil was attributed to paucity of vegetation cover, rapid mineralization of organic matter, inadequate return of crop residue, bush burning and short fallow periods.

In Table 2, the result shows that all the treatments applied increased the soil pH more than the control. However, the increment of pH when 10tha⁻¹ was not significant, but the significant increase was observed as the rates of application increases for both GB and PM. It was also observed that there was a significant difference in pH as the duration increases. When organic materials are applied to the soil, basic cations are released which will certainly improved the base saturation of the soil and thereby reducing the soil acidity. Soil reaction is an important factor in controlling the nutrient availability and processes that take place in the soil (Onwuka and Nwangu, 2016).

Results for the organic carbon (OC) and total nitrogen (TN) (Table 2) shows that there is a significant increase in both OC and TN content of the soil as the rates of application of GB and PM increases. It was further observed that the TN increases with the OC simultaneously. This could be due to the fact that organic matter is the major source of N in the soil. Higher OC and TN of the treated soil over non treated confirm the fact that GB and TN is an excellent source of soil amendment providing both organic matter and N. It could also be attributed to the presence of N P and K contained in the GB and PM. The positive effect of biochar may be due to a number of factors which may include gradual

abiotic and biotic oxidative release of nutrients and humic substances contained in the biochar (Chan *et al.*, 2008;Fagbenro *et al.*, 2016).

The level of available phosphorus (Table 2) increased significantly with the addition of GB and PM over the control and observed that there is a greater increase at different levels of their application. The increase in N and P after the application of GB and PM could be as a result of increase in microbial activities which could aid the decomposition of organic forms of N and P in the soil (Uwah *et al.*, 2014). The level available P is also observed to be dependent of the duration.

Effective Cation Exchange Capacity (ECEC) in Table 2 reveals that there is a significant increase as both GB and PM were added. The increment was observed to be significantly consistence as the rates of these amendment increases. This was corroborated by Mbah and Mbagwu (2006) and Thomas and Aluko (2016) that reported an increase in ECEC after the application of PM which was further attributed to the availability and adequate supply of organic matter. The ECEC of the soil increases significantly as the duration increases.

Generally, it was observed that all the chemical parameters recorded increased as the duration increases. The increment of these chemical constituents with duration could be attributed to the fact that the nutrients contained in organic materials are slowly released over time as also reported by Onwuka and Nwangu (2016).

Table 1: Physical and chemical properties of the soil before treatment

Parameters	Values
Particle size (g kg ⁻¹)	
Sand	668
Silt	160
Clay	172
Textural Class	Sandy loam
pH(H ₂ O)	5.86
pH(CaCl ₂)	5.52
Organic Carbon (g kg ⁻¹)	8.24
Total Nitrogen	0.45
Avail. Phosphorus (mg kg ⁻¹)	8.89
Exchangeable bases (Cmol kg ⁻¹)	
Ca ²⁺	4.23
Mg ²⁺	1.52
Na ⁺	0.43
K ⁺	0.78
ECEC	4.77

Source: Field work, 2018

Total Nitrogen	0.45
Avail. Phosphorus (mg kg ⁻¹)	8.89
Exchangeable bases (Cmol kg ⁻¹)	
Ca ²⁺	4.23
Mg ²⁺	1.52
Na ⁺	0.43
K ⁺	0.78
ECEC	4.77

Source: Field work, 2018

Table 2: Effect of rates and durations of GB and PM on selected soil chemical properties

Rate (R) (t ha ⁻¹)	pH (CaCl ₂)	Org. Carbon (g kg ⁻¹)	Total Nitrogen (g kg ⁻¹)	Available phosphorus (mg kg ⁻¹)	ECEC (Cmol kg ⁻¹)
0	5.52 ^{bc}	8.20 ^d	0.39 ^c	8.99 ^c	4.62 ^c
10 GB	5.87 ^b	10.55 ^c	0.77 ^b	10.69 ^b	6.88 ^b
10 PM	5.94 ^b	10.03 ^c	0.68 ^{bc}	10.98 ^b	6.76 ^b
20 GB	6.33 ^{ab}	10.98 ^c	0.74 ^b	13.75 ^b	6.97 ^b
20 PM	6.42 ^{ab}	12.45 ^{bc}	0.82 ^{ab}	13.44 ^b	7.11 ^b
30 GB	6.85 ^a	14.98 ^b	0.94 ^a	15.58 ^{ab}	13.65 ^{ab}
30 PM	6.72 ^a	16.66 ^b	0.95 ^a	17.32 ^a	12.98 ^{ab}
40 GB	6.82 ^a	20.42 ^a	0.97 ^a	22.16 ^a	16.85 ^a
40 PM	6.85 ^a	18.87 ^{ab}	0.94 ^a	21.68 ^a	17.77 ^a
SE	0.61	1.19	0.17	4.24	4.34
Duration(D)					
2 Months	5.88 ^b	11.24 ^b	0.74 ^b	9.11 ^c	8.43 ^b
4 Months	6.38 ^a	18.33 ^a	0.79 ^b	14.67 ^b	16.96 ^a
6 Months	6.79 ^a	18.86 ^a	0.96 ^a	22.33 ^a	16.75 ^a
SE	0.27	1.62	0.11	2.01	0.35
Interaction					
R*D	NS	NS	NS	NS	NS

Source: Field Work, 2018

Means with the same letter in a column are not significantly different at 5%. ECEC = Effective Cations Exchange Capacity. NS = Not significant at 5%

Two kilogram of soil samples thoroughly mixed with the amendments was weighed into eighty one poly pots (drainage holes were earlier made on each of the poly pot but covered with lining materials before filling with the soil). And on weekly basis, the soil columns were wetted with 700 ml of water. Five hundred

Conclusion

The results of this study have shown that the addition of organic materials such as *Gliricidia* Biochar and Poultry Manure greatly

improved the chemical properties of the soil. From the study, application of beyond 20t ha⁻¹ is thus recommended. It can also be concluded and recommended that farmers should apply these organic materials not less than two months before cultivation so as to achieve the full benefits of the application in terms of the release of the nutrients.

References

- Adeyemi M.O., Fakore M.A., Edema A.O. 1999. Effect of poultry Manure and cutting height on the performance of Amaranthus hybridus. *Nigerian Journal of Agronomy*, 2 (1): 12–20.
- Allison, I. E. (1965). Organic carbon : In C. A. Black (ed). *Methods of Soil Analysis*. Agron. 9 Amer. Soc. Agron. Madison Wis pp 374–390.
- Bray, R.H. and L.T. Kurtz, (1945). Determination of total, organic and available forms of phosphorus in soils. *Soil Science.*, 59; 39–45.
- Chan, K., Van Zwieten, I., Messzaros, I., Downie, A and oseph, S. (2007). Agronomic values of green waste biochar as a soil amendment. *Australian Journal of Soil Research* 45: 629–634.
- Chang, C. H., Lehmann, J. and Engelhard, M. H. (2008). Natural oxidation of black carbon in soils: changes in molecular form and surface charge along a climo-sequence. *Geochim. Cosmochim. Acta* 72(6):1598–1610.
- Fagbenro, J. A., Ajilowura, J. O., Sangoyomi, T. E., Oshunsanya, S. O. and Aliku, O. (2016). Effects of post – treated and incubated *Gliricidia* biochar samples on the growth of maize seedlings. *Nigeria Journal of Soil Science* 26:17–26.
- Farhad, W., Saleem, M. F., Cheema, M. A. and Hammad, H. M. (2009). Effect of poultry manure levels on the productivity of spring maize (*Zea mays* L). *Journal of Animal and Plant Sciences* 19(3):122–125
- Jones, M.J. and A.Wild, (1975). *Soil of West Africa Savannah*. Commonwealth Agricultural Bureaux, Harpenden.U.K.pp.24.
- Mbah, C.N and Mbagwu, S. C. (2006). Effect of animal wastes on physic-chemical properties, yield and root growth of Amaranthus (*Amaranthuscruentus* L.). *African Journal of Science and Technology* 1(4):14–21.
- McLean, E.O. 1982. Soil pH and lime requirement. In: A. L. Page *et al.* (eds), *Methods of soil analyses* (No. 9, part 2), 199–224. *American Society. of Agronomy;* *Soil Science. Society. American;* Inc. Madison, Wisconsin, U.S.A.
- Nguyen, H. Q. (2010). Long – term effects of land application of poultry manure on crop production and soil and water quality under a corn-soybean rotation system in Iowa. Graduate Theses and Dissertation paper 11718.
- Ojeniyi, S. O. (2000). Effect of goat manure on soil nutrient contents on okra yield in rainforest area of Nigeria. *Applied Tropical Agriculture* 5:20–23
- Okon, E. A., Usoro, I. E. and Effiong, G. S. (2016). Effect of goat and poultry manure application on some selected soil properties and yield of garden egg on acid sand of Akwa Ibom State of Nigeria. *Nigeria Journal of Soil Science* 26:319–324.
- Onwuka, M. I. and Nwangwu, B. C. (2016). Characterization of biochar produced from diverse feedstocks used as amendments on an acidic ultisols at Umudike, Abia State *Nigeria Journal of Soil Science* 26:254–264.
- Uwah, D. F., Undie, U.L. and John, N. M. (2014). Comparative evaluation of animal manures on soil properties, growth and yield of sweet maize (*Zea mays* L. *Saccharata* Strut). *Journal of Agriculture and Environmental Sciences* 3(2):315–331.
- Thomas, E. Y. and Aluko, A. P. (2016). Effect of poultry manure on early growth (*Treculia africana* (Decne) seedlings and its impact on the soil chemical properties. *Nigeria Journal of Soil Science* 26:181–189.