



PROCEEDINGS
Of the
35th Annual National Conference
Of
FISHERIES SOCIETY OF NIGERIA
(FISON)

Held at
Raw Materials Research and Development,
Abuja

“MINNA 2020”

Published by
Fisheries Society of Nigeria
Old College Road, Wilmot Point Road, Ahmadu Bello Way,
Bar Beach Busstop, V. I. Lagos State. P. O. Box 2001, Apapa, Lagos
Email: fison2011@yahoo.com, fison1976@gmail.com, www.fison.ng.org
Tel: [08023325185](tel:08023325185), [08023545803](tel:08023545803), [08185477818](tel:08185477818)

Editor's Comment	4
Communique of the 35 th Annual National Conference of FISON	5
NATIONAL EXECUTIVE COUNCIL AND LOC, MINNA 2020	7
AQUACULTURE	15
Evaluation of growth performance of the predatory <i>Megalops atlanticus</i> reared in different culture systems <i>Okoro, C. B., Anwa-Udondiah, E. P., Ejiogu, I. N. and A. O. Olurole</i>	16
Performance of African catfish (<i>Heterobranchus bidorsalis</i> and <i>Clarias gariepinus</i>) hybrid eggs and hatchlings in clay pot, plastic and glass aquarium <i>K. S. Tukura (Mrs)¹, S.M. Tsadu and A.T. Yisa</i>	20
Standardization of <i>Clarias gariepinus</i> fingerlings using age, weight and length estimations <i>Nwangwu, D.C., Yisa, M., Bake, G.G., Orire, A.M., Ogbuebunu, K., Mogaji, O., Goni, I.M., Abubakar, G.R., Yahaya, M., Aremu, A., & Mohammed, U.F.</i>	24
Proximate analyses of sex reversed <i>Oreochromis niloticus</i> (Linnaeus, 1758) fry treated with different doses of 17 α methyl testosterone <i>K. E. Ogbuebunu, V. O. Onyeche, M.I. Goni, and S. A. Olupinyo</i>	28
Comparison of haematological profiles of <i>Oreochromis niloticus</i> cultured in fibre tank and earthen pond. <i>Lawal, B.M., Oke, A.I., Olurole, A.O., Aboluwodi, A.S., Awonuga, A. A.</i>	33
Some morphometric and meristic characteristic of cross between albino <i>Clarias</i> and normal <i>Clarias gariepinus</i> <i>Shettima, A, Yisa T. A Tsado M.S, Orire A.M, Sogbesan O. A and Apollos T. G</i>	38
Effects of using different water sources in the hatchability, growth and survival rate of <i>Clarias gariepinus</i> fry <i>Kigbu A.A., Imgbian T.D. and Mohammed, R.</i>	42
Spawning periodicity and fecundity in <i>Coptodon guineensis</i> reared in sheltered systems <i>Oyenyi, O. V. Soboyejo, I. O. and E. P. Anwa-Udondiah</i>	46
Reproductive indices of female <i>Clarias gariepinus</i> fed different inclusion levels of fertility enhancing seed powder <i>Onyia, L. U., Bishara, Y. A., Onyia, E.C. and Duwal, S.D.</i>	49
Growth performance of monosex cultured <i>Clarias gariepinus</i> Juveniles in concrete flow-through and stagnant water tanks. <i>Robert E.A., Yisa A.T. and Tsadu S.M.</i>	53
The Role of Aquaculture in Enhancing Nutrition and Food Security in the World <i>Godwin, S.O. and Akinrotimi O.A</i>	57
FISH DISEASES	60
Gills and intestinal parasites of elephant snout fish <i>Mormyrus rume</i> (Valenciennes, 1847) at Uke River, Karu Local Government Area, Nassarawa State, Nigeria. <i>Banyigy, A. H., Akpanva, Y. R., Makpo, J. K. and Ameh, S. M.</i>	61
Sensitivity of Bacteria Isolated from Smoked <i>Clarias gariepinus</i> Treated with Scent leaf (<i>Ocimum gratissimum</i>) Juice <i>Alagun O. Olateju G. B. and Akinyemi, A. A</i>	66

Microbial Composition of Effluent and Fish Samples from Selected Water Bodies and Farms in Lagos State, Nigeria <i>Oguntade, O.R. and Olajuyigbe, O.O.</i>	70
FISH NUTRITION	74
Effects of Replacing Soybean with Beniseed, <i>Sesame indicum</i> Seed Meal on Some Haematological parameters Of <i>Clarias gariepinus</i> Juveniles. <i>Inya, F. U., I.B. Okey, S.N. Ochang & J. J. Ekpenyong</i>	75
Dietary Effects of Pineapple (<i>Anana comosus</i>) Crush Waste on the Proximate Composition of <i>Clarias gariepinus</i> Juveniles <i>Binyotubo, O.I., Binyotubo, T.E., Kwen, K., Anwuri, P.A., Akegbejo, S.Y. and Omoniyi, I.T.</i>	81
Effect of storage on nutritive value of Eco float and vital commercial Fish feed <i>Gusau, I. Mamman, T., Jibrin, H., Hambali, I., Abubabakar, M. A and Muhammad U.</i>	87
Effect of garlic (<i>Allium sativum</i>) as feed additive on the performance and survival of African catfish (<i>Clarias gariepinus</i>) <i>Kolo R.J., Orire, A.M., Jidelola F. B. and Haruna, M.A.</i>	92
Growth performance, hematology and intestinal microflora of <i>Clarias gariepinus</i> (Burchell, 1822) fingerlings fed turmeric (<i>Curcuma longa</i>) supplemented diets <i>Joda O. A., Akinde A.O. and Ibidapo-Obe E.O.</i>	97
Nutritional analysis of <i>Clarias gariepinus</i> cultured with <i>Spirulina</i> supplemented diets <i>Okwuosa, Obinna B. and Ekeledo, Chukwunyere B.</i>	104
Effect of graded level of <i>Ziziphus mauritania</i> leaf meal on growth performance and reproductive indices of <i>Oreochromis niloticus</i> <i>Suleiman, S. B., Babagana, A., Yusuf, Z. A., Mohammed, Z.B., Aliyu, M.H.</i>	110
Nutritional evaluation of animal by-products meal as partial substitute for fishmeal in the diet of African catfish, <i>Clarias gariepinus</i> (BURCHELL, 1822). <i>Kenge, B. N., Ofojekwu, P.C. and Mohammed, S. A.</i>	116
Growth Performance and Feed Utilization Of African Catfish (<i>Clarias gariepinus</i>) Juveniles Fed Varying Inclusion Levels Of Butterfly Pea (<i>Clitoria ternatea</i>) Seed Meal <i>Adedokun, M.A; Adelodun, O.B., Rafiu, R.A,</i>	121
Assessment of some fatty acids composition of cultured African mud catfish (<i>Clarias gariepinus</i>) in Lagos State <i>Salaudeen, M. M., Ezekiel, M. O., Esan, O. M., Uzoalu, I. A., Egbe, F. C..</i>	125
What Fish Can Do To Immune System:A Case Study Of Corona Virus Relief <i>Garba, A.A</i>	129
Nutritional evaluation of animal by-products meal as partial substitute for fishmeal in the diet of African catfish, <i>Clarias gariepinus</i> (Burchell, 1822). <i>Kenge, B. N., Ofojekwu, P.C. and Mohammed, S. A.</i>	133
Growth performance and haematology parameters of <i>Clarias gariepinus</i> juveniles fed <i>Moringa Oleifera</i> leaf meal diet <i>Elizabeth. O. Ibidapo-Obe, Khadijah O. Yusuff, and Abiodun O. Akinde</i>	139
Nutritional Composition of Commercial Fishes from Ikpoba Reservoir and Their Potential Contribution to Recommended Nutrient Intake. <i>Oboh, I. P., Egun, N. K. And Olowo, U. C.</i>	145

Survival rate and performance characteristics of African catfish (<i>Clarias gariepinus</i>) fed varying inclusion levels of cow rumen epithelial tissue Rafiu, R.A, Adelodun, O.B., Ajao, F.S., Nworgu, U.C and Adeolu, S.P.	149
Black soldier fly: A perfect replacement of protein source of feed in Nigeria livestock industry (Review) Abubakar, G.R, Goni, I.M, Musa, A., Idris, H., Nwagwu, D.C, Yisa, M, Ibiyo, L.M.O, Ogbuebunu, K.E.	153
The use of non-conventional feed resources for fish nutrition in Nigeria: A review Agbo, A.N. and Onimisi, H. U.	156
Potentials of sweet potato (<i>Ipomoea batatas</i>) leaf meal as dietary ingredient for <i>Oreochromis niloticus</i> fingerlings Joshua, F.O, Ibiyo, L.M.O, Mogaji, O.Y, Woru, H, Azeez, A, Mohammed, A	161
Evaluation of commercial and locally made feed in the culture of African catfish (<i>Clarias gariepinus</i>) in Nigeria Mogaji, O.Y, Ajani, E.K, Olafur, S.	166
Assessment of Some Fatty Acids Composition of Cultured African Mud Catfish (<i>Clarias gariepinus</i>) In Lagos State Salaudeen, M. M., Ezekiel, M. O., Esan, O. M., Egbe, F.C.	171
Comparative growth performance and haematological profile of <i>Clarias gariepinus</i> (Burchell, 1822) fed dietary inclusions of <i>Zingiber officinale</i> , <i>Allium sativum</i> and <i>Azadirachta indica</i> extracts Agatha A. Nwabueze and Ijeoma Q. Atokpe	175
Substitution of soyabean with <i>Moringa oleifera</i> on the growth performance of <i>Clarias gariepinus</i> . Onuoha, P. C. And Okata, J.	179
Effects of dietary lipid on the growth performance and serum biochemistry in carp (<i>Barbonymus schwanenfeldii</i> BLEEKER 1853) fingerling in an aquaponics system. M. A. Sulaiman, L. O. Ibiyo and M. S. Kamarudin	182
Growth performance of maggot meal as a Substitute for Imported Fishmeal in the Culture of <i>Clarias gariepinus</i> Juveniles. Ayo-Olalus, C.I., Oresgun A., Adeleke T.A., Akweeb W.A., Adeyemi, B.Y., Edah, B., Chidume, I; Jibrin, H., Eze, N.C., Olagunju, G.E. and Ayokhai, J.O.	185
Effects of replacement of fish meal with pig blood meal in catfish feeds on their growth, survival and heamotology Agwu, S. C., Oko, A. O., Akinrotimi, O.A., Nwezza, S.N., Onwe, E.U., Jibrin, H., Nwuzor, E. O ² , Ogbuzuru-Ebe, S. L. and Chidume, I.	189
FISHERIES ECONOMICS	194
Cultural practices adopted by farmers in Lafia Local Government Area, Nasarawa State for sustainable aquaculture Adebote, E. E. and Adelakin, C. S	195
Production Constraints Effects On Small Scale Aquaculture In Oyo State Morawo B.O., Ajao F.S.	199
Adaptation Of Fish Farmers To Covid-19 Pandemic In Epe Local Government Area Of Lagos State, Nigeria. E. E. Adebote and Adelakin C.S.	203

COVID-19: The Role of Aquaculture in Employment Generation in Nigeria <i>Jim-Saiki, Ojo Lawal; Ogunbadejo, Kehinde Hussain and Giwa, Esemuze Joseph</i>	207
Post Covid-19 economic recovery: importance of aquaculture fish supply to economic recovery plan. <i>Landu, E. J., Sanni, O.A., Olowosegun, T., Ovwie, P.O., Ibrahim, S.O., Bewaji, O. S and Umar, K.J.</i>	212
Economic analysis of fresh fish marketing in Kede-Tifin district of Mokwa local government area, Niger state, Nigeria. <i>Ndanitsa, M. A., Sallawu, H., Mohammed, D. and Ndako, N.</i>	216
Fish Farmers Socio-economic characteristics predicting the challenges faced during COVID-19: A case of Akinyele Local Government, Nigeria <i>Omega, S., Ametepey, E T.K., and Adebote, E.E</i>	222
Effect Of Income, Socio-Economic Factors On Consumers' Preferences And Quantity Of Fish Consumed By Staff In Different Campuses Of Lagos State. Polytechnic (Laspotech) Nigeria, <i>Sunnuvu T.F, Okunsebor, S.A., Zubairu Z.A, Faleti B.E</i>	226
Blue economy as antedote for crude oil market recession, social insecurity and sustainable development in post COVID' 19 pandemic in Nigeria: Extentionist perspective. <i>Olaniran Olakunle Lateef (Ph.D), Buba Saleh Aliyu and Adetunji Adeyemi Olalekan</i>	232
Factors influencing market price of fish in some coastal communities of Rivers State, Nigeria <i>Okenwa, U., Edun O. M. and Akinrotimi O.A.</i>	237
Factors Affecting Adoption of Innovative Fish Farming Technologies in the Niger Delta <i>Opara, J.Y and Ani, N.A.</i>	241
FISHERIES	244
Some aspects of the biology of the silver catfish <i>Chrysichthys nigrodigitatus</i> (Lacepède, 1803) from great Kwa river, Nigeria <i>Edem Thomas Edem, Victor Oscar Eyo, Bridget Effiong Udobong</i>	245
Relationship between fish species abundance and physico-chemical parameters of Tagwai Lake, Minna, Nigeria. <i>Chukwuemeka V. I.; Tsadu S. M.; Kolo, R. J.; Ojutiku, R. O.</i>	249
Length-Weight Relationship and Condition Factor of Fresh Water Turtle (<i>Pelusios castaneus</i>) Dry Season I Study. <i>Sule, S.O., Mohammed, R. And Sotolu, A.O.</i>	256
Stomach content analysis of mangrove oyster <i>crassostrea gasar</i> from Badagry Creek, Lagos Nigeria <i>Kolawole-Daniels Aghogho, Popoola, K.O.K., A.A. Sowunmi</i>	260
Length-weight relationship and condition factor (K) of <i>Clarias gariepinus</i> (Burchell, 1822) from Kwaru Lake, Tuarare, Dutsin-ma Local Government Area, Katsina State, Nigeria. <i>Nababa, A.S; Yusuf, Z.A; Dambatta, M. and Sadauki, M.A.</i>	265
Aspect of the biology of <i>Hepsetus odoe</i> in Ebonyi River, Southeast, Nigeria <i>Ude, E.F., Ogbonnaya, H. F., Amadi-Ibiam, C. O. And Nwokpor, E.S.</i>	269
Gonadosomatic index, fecundity and sex ratio of <i>Alestes baremose</i> (JOANISS, 1835) in Zobe reservoir, Dutsinma, Katsina State <i>Muhammad, A. Y.</i>	273

Dietary composition and feeding strategy of the big eye grunt <i>Brachydeuterus auritus</i> (VALENCIENNES, 1832) off The Lagos coast <i>Fola-Matthews, O. O. And Fakayode, O. S.</i>	278
Species Composition, Dominance And Similarity Index of <i>Macrobrachium vollehovenii</i> and <i>Macrobrachium felicinum</i> In Akor River, Ibere Ikwuano, Nigeria <i>Ukagwu, J. I., Anyanwu, D. C., Orgi, M. C., Ohaturonye, S, And Dan-Uchegbu, C.</i>	281
Diet of the Common Cuttlefish, <i>Sepia officinalis</i> (Cephalopoda: Sepiidae) (Linnaeus, 1758), Off Lagos Coast, Nigeria <i>Oluboba, T. F. and Lawal-Are, A. O.</i>	284
Species composition, dominance and similarity index of <i>Macrobrachium vollehovenii</i> and <i>Macrobrachium felicinum</i> in Akor River, Ibere Ikwuano, Nigeria <i>Ukagwu, J. I., Anyanwu, D. C., Orgi, M. C., Ohaturonye, S, And Dan-Uchegbu, C.</i>	290
Industrial Fishing In Nigeria: A Recent Assessment Of Fish Landings <i>Jim-Saiki L. O., Eke M., Ogunbadejo H. K., Alhaji T. A. and Giwa J. E.</i>	293
Fish Species Composition And Abundance of Kondo River At New Libata, Kebbi State, Nigeria <i>Ibrahim, Baba Usman and Eitokpah, Priscilla</i>	296
Variations of fishing gears used by fishermen in some communities of Lagos East coast. <i>Bolaji, D.A, Akinnigbagbe, R.O, Obienu, A.J., Oluwakayode-Oluyi, .O.O., Akande, M.O.</i>	300
Variation of Fishing Gears used By Fishers Folks in Some Coastal Communities of Lagos State <i>Oluwakayode-Oluyi Oluwakemi</i>	304
Fish Weight and Species Diversity of Traditional and Modified Malian Traps in Tagwai Dam of Niger State, Nigeria <i>J. Yusuf, S.L. Lamai, M. M. Abdullahi., A. Ibrahim, S.U. Ibrahim., U.P. Yakubu</i>	309
Stomach content analysis of mangrove oyster <i>Crassostrea gasar</i> from Badagry creek, Lagos Nigeria <i>Kolawole-Daniels Aghogho, Popoola, K.O.K, A.A. Sowunmi</i>	314
An assessment of marine fish resource exploitation in Ogun state, Nigeria. <i>Ogunbadejo H. K., Jim-Saiki L. O. and Alhaji T. A.</i>	319
Fishing and natural mortalities as factors depleting the silver catfish, <i>Chrysichthys nigrodigitatus</i> resource of the Cross River, Nigeria. <i>Daniel Ama-Abasi and Edak Aniedi Uyoh.</i>	323
Stomach Content and Feeding Habit of West African Fiddler Crab <i>Uca tangeri</i> From a Tropical Mangrove Lagoon, Lagos, Nigeria. <i>Adejumobi, K.O. Oluwakayode, O.O. Olubola, T.O. and Kasim, O.I.</i>	327
Food and feeding habit and condition factor of tiger shrimp, <i>Penaeus monodon</i> (Fabricius), in the Lagos Lagoon, Nigeria. <i>Aderonke Omolara Lawal-Are, Oluboba, T.F, Oluwajoba, E.O, Oluwakayode-Oluyi O.O, Adejumobi, K.O, Kassim O.I. and Jerome, F.C</i>	333
Assessment of Sea Turtle Bycatch in Artisanal Gillnet Fisheries off The Niger Delta <i>Justina .A. Obienu and Adegbile O.M.</i>	337
Nutritional composition of commercial fishes from Ikpoba reservoir and their potential contribution to recommended nutrient intake. <i>Oboh, I.P., Egun, N. K. And Olowo, U.C.</i>	341
Catch effects of monofilament and multifilament gillnet of different mesh size in Shabu stream	345

Mohammed, R., Kigbu, A.A. and Usman, M.	350
The Effects of Imported Fishes on Nigeria Fishing Industry Alhaji T.A, Jim-Saiki L.O., Ogunbadejo H.K and Mohammed R.	353
POST HARVEST	354
<i>Dermestes maculatus</i> degeer, 1774 (Coleoptera: Dermestidae): A threat to fish preservation (review) Nasir U. A., Suleiman, M. and Abdulkarim, B.	358
Preference and Perception of Fish Processors on the Influence of Wood Types on the Quality of Smoke-dried Fish in Katsina State Dauda, A.B., Oladapo O.O., and Garuba A.M.	363
Development and evaluation of cookies supplemented with fish protein concentrate. Omojowo F.S; Ihuahi J.A.; Ayuba A.B; Isah M.A. and Unogwu, A.	367
Preference and Perception of Fish Processors on the Influence of Wood Types on the Quality of Smoke-dried Fish in Katsina State Dauda, A.B., Oladapo O.O., and Garuba A.M.	372
Effect of some spices' combination on the proximate composition of <i>Clarias gariepinus</i> processed under hot-flame from charcoal (Chorkor oven) Umar, K.A., Yusuf, J.K. and Abdullateef, M.M.	377
Organoleptic characteristics and eating quality acceptability of traditionally smoke-dried freshwater fishes in Toru-orua, Bayelsa State Tina Enize and John F. Alfred-Ockiya	380
TOXICOLOGY AND WATER QUALITY	381
Assessment of Nutrient Load in Wastewater and Sediment-Pore Water from Catfish Ponds in Monai Cluster Fish Farm, Southern Basin of Kainji Lake Mu'azu, M.M., Atiribom, R.Y. and Mbagwu, I.G	385
Stoichiometric Nutrient Limitation and Microalgal Flora of A Former Sewage Disposal Site, at Iddo, Lagos Lagoon, Nigeria. Elegbeleye, O. W., Yusuf, W.A., Offiah, V.C., Okonkwo, C.E.	389
Behavioural Responses and Mortality of <i>Clarias gariepinus</i> Juveniles Exposed to Acute Concentrations of Paraquat Okey, I. B., E.O. Ayotunde, B. U. Patrick & E. E. Ewelu	394
Macro Benthic Invertebrates Assemblage in different Anthropogenic Activity Zones in Badagry Creek, South Western Nigeria. Bosede Omolara Ayo-Dada and Gbolagade Akeem Lameed	397
Effects of Covid-19 lockdown on plankton communities in the Lagos Lagoon Okonkwo, Cleopatra Ebere, Yusuf, Waheed Abiodun, Igwegbe, Adeline Nkechi, Nicholas, Eteobong	403
Mercury Accumulation in the Edible Bivalve <i>Crassostrea gasar</i> (Adanson, 1757) Collected from the Western Creeks of the Lagos Lagoon, Nigeria Lawal-Are, A.O, Moruf, R.O, Mbanefo, I.S. And Sodimu, T.J	407
Seawater flux and its Consequences on Ayetoro coastal community of Ondo State, Nigeria. Olakunle, G.W., R.O.D. Shelle and S.O. Popoola.	412
Assessment and Management of impact factors on Marine mammals during geophysical survey in Nigeria coastal waters. Olakunle, G.W., O.M. Adegbile and A.W. Yusuf.	

Selected Trace Metal Levels in Herring (<i>Clupea harengus</i> , Linnaeus 1758) and Atlantic mackerel (<i>Scomber scombrus</i> , Linnaeus 1758) from major Storage facilities in Maiduguri, Northeast, Nigeria <i>Olanrewaju, A.N., Kareem, O.K., Agbelege, O.O. and Ogunkoya, F.T.</i>	415
Effect of ammonia (NH ₃) toxicity in freshwater on the growth and haematology parametrs of <i>Clarias gariepinus</i> fingerlings <i>Kenneth-Ugochukwu, I.B., Okafor, E.</i>	418
Relationship between heavy metals uptake in crab organs and sediment of Lagos Lagoon <i>Jerome F. C.</i>	423
Toxicity of Lime Fruit Juice to an African Leech (<i>Hemiclepsis quardrata</i>) and Its Haematophagous Potential. <i>Ajibade Akinniyi Oyetunji, Adeosun Olubunmi, Obisesan Omodolapo Morohunranti, Adebukola Folake Bankole, and Rashidat Rafiu</i>	428
Nano-Clay Based Filter For Treatment of Fish Pond Wastewater <i>F. H. Kolawole, R. O. Ojutiku and A. S. Abdulkareem</i>	432
Acute Toxicity of Water Accommodated Fractions (WAFs) of Three Nigerian Crude Oils to Black Jaw Tilapia, <i>Sarotherodon melanotheron</i> <i>Omogoriola Hannah Omoloye</i>	435
Assessment of Heavy Metals in E-waste Leachates: A Case Study of E-waste Dumpsites in Lagos and Osun States, Southwest Nigeria <i>Igbo, J.K. and Nwoko, Chidinma. J.</i>	440
Bioaccumulation of some heavy metals in selected fish species in Shiroro Lake, Niger state, Nigeria. <i>Yusufu, F.O, Kolo, R.J, Ojutiku, R.O and Ibrahim, S.U</i>	445
Toxicity effects of sub-lethal concentrations of cadmium (Cd ²⁺) on haematological parameters of fingerlings and juveniles of freshwater fish, <i>Clarias gariepinus</i> (Burchell, 1822) <i>Jacob, L.T., Auta, J., Kogi, E. and Habila, J.D.</i>	451
Seasonal variations in the occurrence and distribution of pharmaceuticals in the Odo-Iya Alaro river, Lagos state Southwest Nigeria <i>Olatayo M. Ogunbanwo</i>	457
Distribution and composition of aquatic insects in relation to the physicochemical parameters of Awba reservoir, Ibadan <i>Kolawole-Daniels Aghogho, Popoola K.O.K, Nicholas Eteobong</i>	464
Evaluation of the transfer of heavy metals from pig manure to maggot, house fly larvae (<i>Musca domestica</i>) <i>Yusuf, A., Onimisi, M.M., Olorundare. T.L.</i>	467
Assessment of Bioaccumulation Rate of Heavy Metals in Drill Cuttings by Mangrove Periwinkle (<i>Pachymelania aurita</i>) of the Lagos Lagoon. <i>Igwegbe, A.N.</i>	470
Impact of some hydrological factors on biodiversity of fish in Agaie/Lapai dam reservoir of Niger State, Nigeria <i>Yakubu, U.P., Ibrahim, S.U., Yusuf, J., Yusuf, M.K.</i>	475
LIST OF AUTHORS	482

Fish Weight and Species Diversity of Traditional and Modified Malian Traps in Tagwai Dam of Niger State, Nigeria

J. Yusuf^{1*}, S.L. Lamai², M. M. Abdullahi³, A. Ibrahim⁴, S.U. Ibrahim⁵, U.P. Yakubu⁶

^{1,2,4,5,6}Federal University of Technology Minna, Niger State

³Nigeria Institute for Oceanography and Marine Research (NIOMR)

P.M.B. 12729, 3-Wilmot Point Road, Victoria Island, Lagos, Nigeria.

*Corresponding Author's E-mail: j.yusuf@futminna.edu.ng GSM: +2348136161501

Abstract

Over the last century riverine ecosystem have suffered from intense fishing, resulting to over exploitation and extinction of several species of fish. This study seeks to determine fish weight and species diversity of Tagwai Dam using traditional and modified Malian traps. The traps were designed using computer paint application and constructed with *Mimosa pigra* wooden frame. Traditional Malian trap (TMT) has cone shape, while the modified Malian traps has semi-circular (MSCMT) and rectangular shape (MRMT) respectively. The traditional Malian traps were enclosed in 3.75 cm mesh-size while the modified Malian traps were both enclosed in a netting material of 5 cm mesh-sizes (Standard). A total of 236 fishes were caught by both traps. traditional Malian trap caught fishes comprising of *Sarotherodon galileus*, *Tilapia zillii*, *Clarias gariepinus*, *Oreochromis niloticus* and *Synodontis membraneceous*. While modified semi-circular Malian trap caught fishes of similar species and modified rectangular Malian traps caught fishes which were similar species with addition of *Hydrocynus forskalii*. The total weight of the fish caught were 7,247.20 g, 4,154.20 g and 4,209.78 g for (TMT), (MSCMT), and (MRMT) respectively. There was no significant difference ($p > 0.05$) in the biomass and species of fish caught with the Traps. This shows that despite the large netting mesh size of the modified Malian traps and low number of catches, it has the potential of competing favourably with its traditional counterpart. Therefore, modified Malian traps of this nature are recommended as tools in the study of ichthyofaunal composition and sustainable management of fishery resources.

Keywords: Mesh size, Cone shaped Malian trap, Semi-circular shaped Malian trap, Rectangular shaped Malian trap, Tagwai Dam.

Introduction

Nigeria is blessed with abundant natural water bodies with diverse fish resources. Most Nigeria's populations dwelling near water bodies such as lakes, lagoons, reservoirs, rivers, swamps and coastal lagoons depend heavily on the resources of such water bodies for their main source of livelihood (Abubakar *et al.*, 2006). The fish productivity of most Nigeria inland waters is generally on the decline (Jamu and Ayinla 2003). The decline of these fishes has been attributed to several causes ranging from environmental degradation of the water bodies to inadequate management of the fisheries resources. For sustainable exploitation of these resources, a crucial management tool is to have a comprehensive understanding of the ichthyofaunal composition of the water bodies. Lawson and Olusanya (2010) reported that species diversity is the number of different species in a given area. They are commonly used in conservation studies to determine the balance of ecosystem and their species. Information generally on fish abundance studies of natural water bodies is helpful in the sustainable management of the aquatic resources. Lawson and Olusanya (2010) further noted that for the sustainability of fish resources to be attained, it is pertinent to have adequate knowledge of species diversity, composition and relative abundance of the water bodies. Also, the avoidable decline of fisheries resources in an area occasioned by overexploitation and inadequate management of inland waters could be unraveled through the availability of relevant information of various parameters of species in the river. According to Suter (2007), species diversity and relative abundance studies have been advocated for ecological risk assessment in the aquatic system. Fish abundance studies could also help identify the presence of species of importance and economic value to the livelihood of the people living in an area. However, Jamu and Ayinla (2003) reported that the yields of most inland waters are generally on the decline as result of improper management of fisheries resources, environmental degradation such as water pollution. This study is aim at determine the fish species diversity and weight of Tagwai dam reservoir with a view of providing information on the ichthyofaunal composition of the dam and the potential of modified alian traps in relation to its traditional counterpart.

Materials and methods

Study Area

The research was carried out at Tagwai Dam located in Minna, Niger State. The geographical co-ordinates of the Dam are Latitude 6°39' to 6°44' East and Longitude 9°34' to 9°37' North to South East of Minna-Suleja Road (Fig. 1). (Alkali, 1994).




Figure 1. Nigeria and Niger State inset (Tagwai Dam)
Source: (Abdullahi, 2015)

Traditional Malian trap

Wooden stick of giant sensitive plant (*Mimosa pigra*) were used as wooden frame for the construction. The diameters of the stick ranges from 3 cm to 4 cm respectively (Fig. 2). The sticks were then bent to form round shape of different diameters 130 cm, 100 cm and 70 cm respectively, and tied tightly with (rope size number 9) to avoid loosening. Wooden frames of 75 cm height were mounted and tied on the round bottom. The same procedure was repeated for the middle and the tops. Thereafter, the structures were covered with polyamide netting material (3.75 cm) mesh-size. 3 spaces for non-return valves were carved out and they were finally fixed. 6 traditional Malian Traps were constructed (plate I)




Figure 2: A sketch of a traditional Malian trap
Source: (Field Design, 2018)

Plate I: Traditional Malian traps after construction trap (TMT)
Source: (Field work, 2018)

Modified semi-circular Malian trap

Again, the giant sensitive plant (*Mimosa pigra*) was used for the construction of modified Malian trap. 100 cm length, and 50 cm width of the wooden stick were tied together to form rectangular base. See (Fig. 3). This was followed by the mounting of bent wooden frame of 50 cm height. The frame was supported with straight wooden sticks strongly




Figure 3: A sketch of a modified semi-circular Malian trap Source: (Field work, 2018)




Plate II: Modified semi-circular Malian trap after construction. Source: (Field design, 2018)

Modified Rectangular Malian Trap

The construction process was similar to figure 2, only the shape was rectangular (plate III).




Figure 4: A sketch of modified rectangular Malian trap Source: (Field work, 2018)




Plate III: Modified rectangular Malian trap Source: (Field work, 2018)

Traps Setting, Inspection, Monitoring and Collection of Catches

After conveyance, the traps were set and monitored twice in a week for a period of 26 weeks (6 Months) making the total of 48 fishing visits conducted in different locations of the Dam. At each monitoring visits, the traps were lifted out of water and the weaved top opening was loosed to collect the trapped fish into a cooler jug. Thereafter, the loosed top opening was weaved back and the traps were set again for subsequent monitoring. Both the traditional and modified traps were moved backward to the littoral part as the water volume increases to prevent their vulnerability to flood.

Determination of fish weight

The weight of the fish was determined by placing the fish on the top tray of the weighing balance (Citizen model) and the reading was observed and recorded for each of the trapped fish.

Identification of species

Monograph of Olaosebikan and Raji (2013) was used to observe the similarity in the morphological features of the trapped fishes and those in the monograph and each species was identified in the process.

Results

Number and weight of fish caught with traditional and modified Malian traps

Table 1. depicts the monthly number and weights of fish caught with traditional and modified Malian traps. The total weight of the fish caught were 7,247.20 g, 4,154.20 g and 4,209.78 g for (TMT), (MSCMT), and (MRMT) respectively. There was significant difference ($p > 0.05$) in the biomass of fish caught with the traps

Table 1: Number and Weight of the fish Caught with Traditional and Modified Malian Traps

Months	TMT	MSCT	MRT	Malian Traps		
				TMT (g)	MSCMT (g)	MRMT (g)
July	29	27	25	232.25 ± 73.13 ^a	413.68 ± 137.5 ^a	478.61 ± 160 ^a
August	33	7	5	351.62 ± 115.5 ^a	58.3 ± 8.66 ^b	69.05 ± 18.92 ^b
September	19	6	2	179.32 ± 54.2 ^a	23.9 ± 8.42 ^b	40.37 ± 6.2 ^c
October	12	3	4	86.15 ± 20.3 ^b	26.75 ± 8.43 ^b	27.9 ± 8.3 ^d
November	20	7	4	181.2 ± 54.93 ^a	25.72 ± 8.46 ^b	41.6 ± 5.77 ^c
December	18	9	7	177.33 ± 53.54 ^a	144.03 ± 41.58 ^a	44.12 ± 4.62 ^c
TOTAL	131	58	47			

Data on the same column carrying different superscript are significantly different from each other (P<0.05).

Where: ±SDM= standard deviation of the mean; TMT= traditional malian trap; MSCMT= modified semi-circular malian trap; MRMT= modified rectangular malian trap

Fish Species Diversity of Traditional and Modified Malian Traps

Table 2 expressed the family, species and number of fish caught with traditional and modified Malian traps. A total of four family, and six species were caught. The highest family of fish caught was *Cichlidae* accounting for 113 *S. galileus*, 45 *T. zillii*, and 5 *O. niloticus* amounting to total of 163, followed by *Mochokidae* with a value of 62 *S. membranaceous* recorded, next is *Claridae* accounting for the total of 6 *Clarias gariepinus* recorded and the lowest was *Characidae* accounting for 5 *H. forskalii* recorded.

Table 2: Fish species diversity of traditional and modified malian traps

Family	Species	Shape of Traps		
		TMT	MSCT	MRT
<i>Claridae</i>	<i>C.gariepinus</i>	1	3	2
<i>Alestidae</i>	<i>H. forskali</i>	0	0	5
<i>Cichlidae</i>	<i>O. niloticus</i>	3	2	0
<i>Cichlidae</i>	<i>S. galileus</i>	61	27	25
<i>Mochokidae</i>	<i>S. mebraneaceous</i>	39	13	10
<i>Cichlidae</i>	<i>T. zilli</i>	27	13	5
Total		131	58	47

Where: TMT= Traditional Malian trap; MSCMT= Modified semi-circular Malian trap; MRMT= Modified Rectangular Malian trap

Discussion

The study on fish weight and species diversity of traditional and modified Malian traps revealed the potential of the traps in catching different species of fishes with their corresponding weight. A total of 236 fishes weighing 15, 611.18 g of different size and species were caught. Ayanwale *et al.* (2013) reported that variation in mesh- size of gears influenced number of catches. This conform with the result of this study in which there was a variation in mesh size used for traditional and modified Malian traps. Akinngbagbe and Osibona (2017) reported 47.7 g as the highest mean fish weight in their study of Fish weight and species diversity of traditional and modified traps in selected fishing communities in Lagos lagoon. The reason for the differences could be as a result of differences in study location. There was no significant different (p>0.05) observed in weight of fish caught among the three different shapes of the traps in this study. This agreed with the report of Ago *et al.* (2012) who observed no significant difference (p>0.05) in the catch (number and weight) of the two gear types used in their study. Agbelege *et al.* (2004) also reported 37.68 kg as the total biomass of fish caught with no significance difference in the biomass of fish caught in the 4-v and 6-v traps. This also tally with the result of this study. Trisnani *et al.* (2016) reported (18 kg) of fish using common *Payang* fishing trap while modified *Payang* fishing trap caught (15 kg) of fish with no significant difference (p>0.05) observed in the two weights. This also conform with the result of this study.

In this research, a total of four family comprising of six species were caught. The highest family of fish discovered was *Cichlidae*. This finding is in line with Ayanwale *et al.* (2013) who reported *Cichlidae* as the Preponderance fish family comprising of *S. galileus* and *T. zillii* in Tagwai dam. Thus, they further reported *Mormyridae* as one of the

abundant species in the Dam also. This contradicts the finding of this research in which *S. mebraneceous* was the next abundant after *Cichlidae*. This could be as a result of differences in gear used, time of the study, over exploitation or migration of this aforementioned species. The finding of this study does not conform with that of Ipinjolu *et al.* (2004) who reported *O. niloticus* as the most abundant species in their study. The variation in their species result with this study could be attributed to location of the study. Mshelia *et al.* (2015) reported *Cichlids* as the dominant species in Lake Alau. Similar reports were observed by Balogun *et al.* (2000) in Zaria reservoir; Kangimi lake (Balogun and Auta, 2001) and Ero reservoir (Kester *et al.*, 2007). The finding of Trisnani *et al.* (2016) where trevally fish sp. was reported to be the dominant fish species using traditional and modified Payang fishing traps were used also disagreed with the finding of this study.

Conclusion

The study revealed that Modified Malian traps can compete favourably with its traditional counterpart in trapping different species, size and weight of fish in Tagwai Dam. The species of fish caught with the traps shows an extinction of few valuable species such as *Mormyrus* and *Alestes* species which were hitherto reported to be among the dominant species in the Dam. Therefore, information obtained on the ichthyofaunal composition in this study can be use in sustainable management of the dam.

References

- Abubakar, K.A., Haruna, A.B., and Ladu, B.M.B. (2006). An assessment of the physiochemical parameters and productivity of Lake Geriyo, Yola, Adamawa State, Nigeria. *Biological and Environmental sciences Journal for the Tropics* 3(1):18-23
- Akinnigbagbe, R.O. and Osibona, A.O. (2017). Fish weight and species diversity of Traditional and modified traps in selected fishing communities in Lagos Lagoon *International Journal of Fisheries and Aquatic Studies* 5 (4): 363-370
- Alkali, A.L. (1994). Some water quality parameters and potential fisheries yield in Tagwai Dam, Minna, Nigeria. M. Tech Thesis, Fisheries Department, Federal University of Technology, Minna, Nigeria.
- Ayanwale, A.V., Shokunbi, M.T., Olayemi, I.K., Chukwuemeka V.I., Falusi, F.M. and Erhabor, O.F. (2013). A study of the fish fauna of Tagwai lake Minna, Nigeria, in Relation to Gear Selectivity. *Pakistan Journal of Biological Sciences*, 16: 731-734.
- Balogun, J. K. and Auta, J. (2001). Fisheries resource and development potentials of Lake Kangimikaduna state. *Nigeria Journal for Bioscience*, 1, 50-56
- Balogun, J. K., Auta, J. A., Balarabe, M. L. and Bako, S.P (2000). Fisheries management and development considerations for small reservoir: A case study of Zaria Reservoir: Paper present at the Fisheries Society of Nigeria (FISON). 15th Annual Conference, Jos, Nigeria
- Ibrahim, A., Lamai. S. L., and Yisa A.T (2015). Growth and Survival of *Sarotherodon galileus* and *Tilapia zillii* Obtained from Tagwai Dam into Federal University of Technology Research Farm facilities
- Ipinjolu, J. K., Agbelege, O. O. and Hassan, W. A. (2004). The performance of Malian trap using four baits types in Lake Kainji, Nigeria, presented at *FISON Annual Conference*, Ilorin.
- Jamu, D.M. and Ayinla, A.O. (2003). Potential for the Development of Agriculture in Africa. *NAGA*, 26(3): 9-13.
- Kester, C. T., Osofero, S. A. and Daramola, J. A. (2007). Fisheries potential assessment of Ero reservoir, Ekiti state. *Nigeria. Agricultural Journal*; 2(6):721-725.
- Lawson, O.E. and Olusanya O.M. (2010). Fish Diversity in Three Tributaries of River Ore, South West, Nigeria. *World Journal of Fish and Marine Sciences*. 2010; 2(6):524-531.
- Olaosebikan, B. D. and Raji, A. (2013). *Guide to Nigeria Freshwater Fishes*. FCFFT Publication 1st edition. Ilorin: Decency Printers. 106.
- Suter, G. W. (2007). *Ecological Risk Assessment*. 2nd edition. CRC Press, Boca Raton, Florida
- Trisnani, D. H. and Aristi, D.P. (2016). *Technical and Economic Analysis of Products Processing and Health Aquatic Procedia* (7): 254 – 264u