Assessing the Effects of Potential Climate Change on Fisheries of River Zungeru, Niger State, Nigerland L. M. Obiaso

Abstract
Fishes have generally evolved physiologically to live within specific environmental variations, and existence outside the second as a first of climate change on fisheries of River Zung. Abstract
Fishes have generally evolved physiologically to live within specific environmental variations, and existence outside these can be stressful or fatal. Extreme environmental factors such as, elevated water temperature can have deleterious effects of climate change on fisheries of River Zungeru, Nicola a number of qualitative methods; focus group de Nicola and existence outside these can be stressful or fatal. Fishes have generally evolved physiologically to the can be stressful or fatal. Extreme environmental factors such as, elevated water temperature can have deleterious entirely can be stressful or fatal. Extreme environmental factors such as, elevated water temperature can have deleterious entirely can be stressful or fatal. Extreme environmental factors such as, elevated water temperature can have deleterious entirely can be stressful or fatal. Extreme environmental factors such as, elevated water temperature can have deleterious entirely can be stressful or fatal. Extreme environmental factors such as, elevated water temperature can have deleterious entirely can be stressful or fatal. Extreme environmental factors such as, elevated water temperature can have deleterious entirely can be stressful or fatal. Extreme environmental factors such as, elevated water temperature can have deleterious entirely can be stressful or fatal. Extreme environmental factors such as, elevated water temperature can have deleterious entirely can be stressful or fatal. Extreme environmental factors such as, elevated water temperature can have deleterious entirely can be stressful or fatal. Extreme environmental factors such as, elevated water temperature can have deleterious entirely can be stressful or fatal. Extreme environmental factors such as fatal entirely can be stressful or fatal entirely can be stressful o can be stressful or fatal. Extreme environments of climate change of the first stress of kiver Zungeru, effects of climate change of the fishes. This research was carried out to assess the potential effects of climate change of kiver Zungeru, kigeru and kiver zungeru. Kigeru and kiver zungeru, kigeru and kiver zungeru, kigeru and kiver zungeru, kigeru and kiver zungeru. Nigeria. Data collection included cross-sectional survey with a number or quantative metals, focus group discussions. Nigeria. Data collection included cross-sectional survey with a number or quantative metals. Nigeria. Data collection included cross-sectional survey with a number of quantative metals. Nigeria, focus group discussions of the secondary data on rainfall and temperature were gotten from Nigerian Metals and number of secondary data on rainfall and temperature were gotten from Nigerian Metals and number of secondary data on rainfall and temperature were gotten from Nigerian Metals and number of secondary data on rainfall and temperature were gotten from Nigerian Metals and number of secondary data on rainfall and temperature were gotten from Nigerian Metals and number of secondary data on rainfall and temperature were gotten from Nigerian Metals and number of secondary data on rainfall and temperature were gotten from Nigerian Metals and number of secondary data on rainfall and temperature were gotten from Nigerian Metals and number of secondary data on rainfall and temperature were gotten from Nigerian Metals and number of secondary data on rainfall and temperature were gotten from Nigerian Metals and number of secondary data on rainfall and temperature were gotten from Nigerian Metals and number of secondary data on rainfall and temperature were gotten from Nigerian Metals and number of secondary data on rainfall and temperature were gotten from Nigerian Metals and number of secondary data on rainfall and temperature were gotten from Nigerian Metals and number of secondary data on rainfall and temperature were gotten from Nigerian Metals and number of secondary data on rainfall and temperature were gotten from Nigerian Metals and number of secondary data on rainfall and temperature were gotten from Nigerian Metals and number of secondary data on rainfall and temperature were gotten from Nigerian from Nigeria. Data collection included cross-section depth interviews were administered. The secondary data on rainfall and temperature increased, while the amount and number of Nigerian Metrol, Agency (NIMET), Niger State. The study revealed that, temperature increased, while the amount and number of rainfall and temperature peaked companies. Agency (NIMET), Niger State. The study revealed that, temperature increased, which and number of the dominant fish species in the river. It also showed at the complete of damming downwell as, the effect of damming downwell as, the effect of damming downwell as, the effect of damming downwell as the effect of da Agency (NIME 1), Niger State. The dropped Increase in temperature caused fish mortality particularly during dry season with the emperature peaked rainly shocks from the large number of electric fish which was one of the dominant fish species in the river. It also shocks from the large number of electric fish which was one of the dominant fish species in the river. It also showed a downstream to the effect of damming downstream and understanding downstream. dropped. Increase in temperature cause shocks from the large number of electric fish which was one of the dominant rish species in the liver. It also showed all shocks from the large number of electric fish which was one of the dominant rish species in the liver. It also showed all shocks from the large number of electric fish which was one of the dominant rish species in the liver. It also showed all shows a showed a downstream. It also showed a downstream in number of catches and biodiversity of the River due to overfishing, as well as, the effect of damming downstream in number of catches and biodiversity of the River due to overfishing, as well as, the effect of damming downstream in number of catches and biodiversity of the River due to overfishing, as well as, the effect of damming downstream in number of catches and biodiversity of the River due to overfishing, as well as, the effect of damming downstream. shocks from the large name? In number of catches and biodiversity of the River due to overfishing, as wen as, the study showed that, most of the fisher folks were locals with limited knowledge and understanding of climate change.

Introduction

Natural climatic fluctuations, particularly those at medium (decadal) scale have always affected fisheries as well as the company of the compa Natural climatic fluctuations, particularly those at medium (decadar) scare have arrived insurers as well as management performance (Ipinjolu et al., 2014). Global warming means an increase in global mean air temperature. As the management performance (ipinjolu et al., 2014). Global warming means an increase in global mean air temperature. As the management performance (ipinjolu et al., 2014). management performance (Ipinjolu et al., 2014). Global warming means an increase in great an temperature which has wide spread effects on the life him temperature increases, a similar increase is expected in water temperature. Which significantly affects lake levels, fish catches, production and supply the appropriate temperature. physiology and behavior of most aquatic organisms including risnes. Worker, enterthe affects lake levels, fish catches, production and supply white affects lake levels, fish catches, production and supply white affects lake levels, fish catches, production and supply white affects lake levels, fish catches, production and supply white rainfall a particular region experiences, which significantly affects lake levels, that are supply with significance of fisheries is often understated, the implications of climate change for these sectors; and for coastal and supply with the ignore (Barange et al., 2010).

The potential effects of climate change on fisheries of River Zungeru, Niger State, Nigeria was assessed in this research.

Data Collection

Data Collection

Data for this study were generated through a survey on production function which controls the aquatic environment mostly area; and how climate change affect fisheries. Some of the study area; and how climate change affect fisheries. Data for this study were generated through a survey on production rather than the physical, biological and socio-economic factors of the study area; and how climate change affect fisheries. Some of the physical, biological and socio-economic factors of the study area; and how climate change affect fisheries. Some of the physical, biological and socio-economic factors of the study area; and how climate change affect fisheries. Some of the physical ph physical, biological and socio-economic factors of the study area, and for the documented in maps (Figures 1 and 2), graphs, and complemented with participatory questionnaire, focus group discussional documented in maps (Figures 1 and 2), graphs, and complements of potential effect of climate change and the study area. in-depth interview to obtain salient but important human components of potential effect of climate change and to get in-depth the specific actions of the people to avert climate change. The secondary data consisted of ready materials obtained in value and rainfall records from 1984-2010 and only and only and the specific actions of the people to avert climate change. formats such as, meteorological data from NIMET (i.e. temperature and rainfall records from 1984-2010) and relevant literality While the topographic maps of Niger State and its environs were digitalized to act as a baseline map of the study area.



Plate1: Satellite image of River Zungeru




Figure 1: Base Map of River Zungeru

Data Analysis

The data obtained on climate (i.e. temperature and rainfall records) were related statistically and presented in form of grant were analyzed and presented in form of grant presented in grant pre while the primary data were analyzed and presented as tables and themes respectively.

Results and Discussion

Table 1 is a summary of the observable changes around the study area.

PROCEEDINGS OF THE 34TH ANNUAL CONFERENCE OF FISON_

Changes	in	the	River	from	1981	-2011
---------	----	-----	-------	------	------	-------


sh and Fishing	1981	2011	Reasons for the change
etivities ecies availability	Higher	Lower	Rapid increase in number of fisher folk in the community
sh growth rate	Moderate	Higher	Increased availability of food and ability to escape catch
Jortality rate	Lower	Higher	High temperatures especially during dry season as well as, shock from electric fish which happen to be one of the dominant species in the
lumber of	Lower	Higher	High cost of living and increase demand for fish
ishermen Volume of catch	Higher	Lower	Increase in number of fisher folk in the community
Fish population	Higher	Lower	Overfishing and mortality

fish growth, the fisher folk interviewed were of the view that, the fishes grew faster and bigger now than before. This of his grown laster and bigger now than before. This was due to increase in natural food in the river as a result of high rate of fertilizer application on and the river, and that, most fishes have developed abilities of escaping catch. It also showed that, the mortality around and a strong around the of electric fish that dominated the river. The decrease in the biodiversity of the river is a factor of potential climate be of circuit de like overfishing due to the rapid increase in number of fisher folk in the community. This was ascertained the three decadal metrological data obtained which reflected increase in temperature Figures 2 and 3.

12: Fisher Folks Experiences on Climate Change

Process	1981	2011	Reasons for the change
Rate of river flow	Lower	Higher	Construction of dam upstream which spill its bank into the river.
Frequency of flood	Higher	Lower	Damming upstream which regulates the volume of water coming into the river at any point in time
Amount rainfall	Higher	Lower	Drop in number of rain days
Rainfall intensity	Moderate	Higher	Natural
length of dry season	Shorter	Longer	Drop in number of rain days
Intensity of the sun	Lower	Higher	Natural
Temperature	Lower	Higher	High intensity of the sun
Number of trees around the river	Higher	Lower	Erosion and deforestation
Turbidity level	Lower	Higher	Erosion

(Source: Authors' compilation, 2010)



Mean Temperature (°c) of the Study Area (1984-2010)




Figure 3: Annual Mean Rainfall (mm) of the Study Area (1984-2010)

The results of the interview (Table 2) showed that, the increase in the river flow and decrease in the frequency of flow and time. It also repeat the river at a regulated volume and time. It also repeat flow The results of the interview (Table 2) showed that, the interest in the attributable to the daming at Shiroro which spill its bank into the river at a regulated volume and time. It also revealed to the daming at Shiroro which spill its bank into the river at a regulated volume and time. It also revealed to the daming at Shiroro (with moderate intensity) but lower now (with high intensity) Also attributable to the daming at Shiroro which spill its bank into the river at a regulated volume and time. It also revealed to the daming at Shiroro which spill its bank into the river at a regulated volume and time. It also revealed to the daming at Shiroro which spill its bank into the river at a regulated volume and time. It also revealed to the daming at Shiroro which spill its bank into the river at a regulated volume and time. attributable to the daming at Shiroro which spill its bank into the tree amount of rainfall received was higher before (with moderate intensity) but lower now (with high intensity). Also the manual of the control of the spill its bank into the tree and the spill its bank into amount of rainfall received was higher before (with moderate interior). Also the positive due to run-off. This was in agreement with the findings of Edephene of trees around the river were more before but rewer now due to electronic same time increased the turbidity of the river due to run-off. This was in agreement with the findings of Edegbene et al. and the river due to run-off. This was in agreement with the findings of Edegbene et al. same time increased the turbidity of the river due to run-on. This was a same time increased the turbidity of the river due to run-on. This was a same time increased the turbidity of the river due to run-on. This was a same time increased the turbidity of the river due to run-on. This was a same time increased the turbidity of the river due to run-on. This was a same time increased the turbidity of the river due to run-on. This was a same time increased the turbidity of the river due to run-on. This was a same time increased the turbidity of the river due to run-on. This was a same time increased the turbidity of the river due to run-on. This was a same time increased the turbidity of the river due to run-on. This was a same time increased the turbidity of the river due to run-on. This was a same time increased the turbidity of the river due to run-on. This was a same time increased the turbidity of the river due to run-on. This was a same time increased the run-on turbidity of the river due to run-on turbidity of turbi

Conclusion

Conclusion

The study shows that, the mean air temperature around Zungeru River increased with time of while the amount of the study shows that, the mean air temperature around Zungeru River increased with time of while the amount of the study shows that, the mean air temperature around Zungeru River increased with time of while the amount of the study shows that, the mean air temperature around Zungeru River increased with time of while the amount of the study shows that the study shows the study The study shows that, the mean air temperature around Zungers to the study shows that, the mean air temperature around Zungers to the study shows that, the mean air temperature around Zungers to the study shows that, the mean air temperature around Zungers to the study shows that, the mean air temperature around Zungers to the study shows that, the mean air temperature around Zungers to the study shows that, the mean air temperature around Zungers to the study shows that, the mean air temperature around Zungers to the study shows that, the mean air temperature around Zungers to the study shows that, the mean air temperature around Zungers to the study shows that the study shows the study sho number of rain days reduced with a lot of variations. This includes the enlightened about climate change and possible ways of averting some of the change has create awareness and enhance their understanding of the causes and possible ways of averting some of the observed changes

References

Barange, M., Merino, G., Blanchard, J. L., Scholtens, J., Harle, J., Allison, E. H., Allen, J. I., Holt, J. and S. Jenning (N. Impacts of climate change on marine ecosystem production in societies dependent on fisheries. Nature Climate Change 211-216 Available https://www.nature.com/articles/nclimate2119 supplementary-information

Edegbene, A. O., Arimoro, F. O., Nwaka, K. H., Omovah, G. O., Ogidiaka. E and O. J. Abolagba (2012). The physical chemical characteristics of Atapko River, Niger Delta, Nigeria. Journal of Aquatic Sciences, 27 (2):159-172

Ipinjolu, J. K., Magawata, I. and B. A. Shinkafi (2014). Potential Impact of Climate Change on Fisheries and Aquaculus Nigeria. Journal of Fisheries and Aquaculture Science, 9(5): 338-344. Doi: 10:3923/jfas.2014.338.344.