



NIGERIAN SOCIETY FOR ANIMAL PRODUCTION

in collaboration with

UNIVERSITY OF JOS AND FEDERAL COLLEGE OF FORESTRY JOS

47th Annual Conference

////////////////////////////////////

(JOS 2022)

BOOK OF PROCEEDINGS

T H E M E ▶

SECURING ANIMAL AGRICULTURE AMIDST GLOBAL CHALLENGES

////////////////////////////////////

Date: 13th-17th March, 2022

Venue: Dome Theatre,
Federal College of Forestry, Jos

EDITORS:

**Y. P Mancha, D. J. U. Kalla, T. T. Akpensuen
T. T. Igila, J. S. Luka & U. Okpanachi**

ISSN: 0331-2064



**THE 47TH ANNUAL CONFERENCE OF THE
NIGERIAN SOCIETY FOR ANIMAL
PRODUCTION**

March 13th – 17th, 2020

**The Proceedings of the 47th Annual Conference of the
Nigerian Society for Animal Production**

**Editors: Y. P. Mancha, D. J. U. Kalla, T. T. Akpensuen, T. T. Igila, J. S. Luka &
U. Okpanachi**

**Hosted by University of Jos in Collaboration with Federal College of Forestry,
Jos**

under the Auspices of Nigerian Society for Animal Production

TABLE OF CONTENT FOR THE ADDENDUM OF THE NSAP JOS 2022 BOOK OF PROCEEDING

ANIMAL BREEDING AND GENETICS

1	COMPARISON OF GOMPERTZ, LOGISTIC AND RICHARDS GROWTH MODELS FOR DESCRIBING GROWTH CURVES OF DOMESTIC PIGEON SQUABS (<i>COLUMBA LIVIA</i>) IN MAIDUGURI, BORNO STATE, NIGERIA.	*Zannah, B. B., *Yusuf, M. J., **Lamido, M., *Taki, I. M., *Aminami, M. and Mukaddas, J.	1803
2	PHENOTYPIC CORRELATIONS AMONG EGG QUALITY TRAITS OF SHIKA BROWN LAYERS FED DIETS SUPPLEMENTED WITH LYSINE IN THE TROPICS	* ¹ Badmus K.A., ² Orunmuyi M., ³ Onimisi P.A., ³ Bawa G.S. and ³ Abubakar H.	1808
3	RELATIONSHIP BETWEEN BODY WEIGHT AND LINEAR BODY MEASUREMENTS IN MATURED MUSCOVY AND KHAKI CAMBELL DUCKS	Abdu, I ¹ ., Salisu. A, D ¹ ., Yahaya B ² ., Jafar ³ , A. M. and Shuaibu, A ⁴ .	1813

ANIMAL HEALTH AND WELFARE

4	ANTITRYPANOSOMAL ACTIVITIES OF SOME SELECTED NIGERIAN ETHNOMEDICINAL PLANTS	*Obi, C.F., Nzeakor, T.A., Okpala, M.I., Aneru, G.E., Omeje, M.O., Nzekwe, S., Ezeh, I.O., Ezeokonkwo, R. C.	1819
5	SEROPREVALENCE OF <i>Peste des Petits Ruminants</i> IN SMALL RUMINANTS IN DUTSIN-MA LOCAL GOVERNMENT, KATSINA STATE	<i>Tarnongo B, Egbo, M.L, Kalla, D.J.U</i> <i>Animal production Department,</i> <i>Abubakar Tafawa Belewa University,</i> <i>Bauchi.</i> <i>bemmarvellous74@gmail.com</i>	1824

ANIMAL PRODUCT AND VALUE ADDITION

6	PROXIMATE EVALUATION OF SPENT LAYER CHICKEN NUGGETS INCORPORATED WITH POWDER AND EXTRACT OF WATERLEAF (<i>TALINUM TRIANGULARE</i>)	Alonge ¹ , O.O., Olorunsanya ² , A.O., Abdu S. B ¹ ., Shittu Afeez ¹	1828
---	---	---	------

ANIMAL PHYSIOLOGY AND BIOTECHNOLOGY

7	HEMATOLOGY OF MARSHALL BROILER CHICKEN FED DIFFERENT LEVELS OF DIETARY ENERGY DURING THE HOT DRY PERIOD IN SEMI-ARID ENVIRONMENT	IBRAHIM, N.T ^{*1} RIBAH, I.M ^{**2} ABUBKAR, A*** ³ And OSHILAJA, A.A **** ⁴	1833
8	HAEMATOTOLOGY AND SERUM CHEMISTRY VALUES OF PRE-PUBERTAL BOARS FED AIDAN (<i>Tetrapleura tetraptera</i>) POD MEAL	Ezea, J., Ezike, J. C. and Nathaniel, J.	1838
9	HAEMATOLOGICAL PARAMETERS OF JAPANESE QUAILS (<i>Coturnix coturnix japonica</i>) FED DIETS CONTAINING RUBBER (<i>Hevea brasiliensis</i>) SEED MEAL BOILED AT VARYING DURATIONS	Allen, O. G ¹⁺ , Allen, D. M ¹ , Omage. J. J ² , Buba, W ³ , Bunjah-Umar, D.S ⁴ and Kperun, T. N ⁵	1843
10	SERUM BIOCHEMICAL INDICES OF JAPANESE QUAIL CHICKS (<i>Coturnix coturnix japonica</i>) FED DIETS CONTAINING RUBBER (<i>Hevea brasiliensis</i>) SEED MEAL BOILED AT VARYING DURATIONS	Allen, O. G ^{1*} , Omage, J. J ² , Buba, W ³ Allen, D. M ¹ , Kperun, T. N ⁴ and Bunjah-Umar, D.S ⁵	1848
11	EFFECTS OF NEGRO PEPPER, GINGER AND GARLIC AS FEED ADDITIVES ON GROWTH PERFORMANCE, SERUM BIOCHEMICAL AND HEMATOLOGICAL RESPONSES OF BROILER CHICKENS	*Shittu, M. D. *Azeez, K. E., *Azeez, Y. O., +Ojebiyi, O. O., +Ojediran, T. K. and +Williams, O	1853

LIVESTOCK ECONOMICS AND MARKETING

12	PROFITABILITY AND SOLVENCY ANALYSIS OF CATFISH (<i>Clarias gariepinus</i>) PRODUCTION IN JOS METROPOLIS, PLATEAU STATE, NIGERIA.	¹ Folorunso, S.T., ² Gomina, A., ³ Onuwa, G.C. and ¹ Ezeabikwa, F.C.	1858
----	---	---	------

WILDLIFE, MICRO LIVESTOCK, APICULTURE, FISHERIES AND AQUACULTURE

13	LENGTH-WEIGHT RELATIONSHIP AND STOMACH CONTENT ANALYSIS OF AFRICAN CATFISH <i>CLARIAS GARIEPINUS</i> (BURCHELL, 1822) IN AJIWA RESERVOIR KATSINA, NIGERIA	^{*1} Umar, B. D., ² Hassan, Z. Y. and ³ Auwalu J.	1863
14	FISH CULTURE PRACTICES AND CHALLENGES AMONG WOMEN IN TELLA LOCAL GOVERNMENT AREA, TARABA STATE, NIGERIA.	^{*1} J.B. Ogunremi, ² M.M. Onimisi ³ Y.E. Ajibade, ⁴ R.O. Echo	1868
15	THE EFFECTS OF CONCENTRATE FEED SUPPLEMENTATION ON THE THE VISCERAL ORGAN WEIGHTS OF GRASSCUTTERS FED ELEPHANT GRASS AS BASAL FEED	Gboshe, N. P.	1873
16	ACUTE TOXICITY OF CALCIUM CARBIDE WASTE TO SILVER CATFISH (<i>Chrysichthys nigrodigitatus</i>) JUVENILES	Ajibade, A. O., ¹ Adesina, B. T. ² and Obisesan, O. M. ³⁻	1877

MONOGASTRIC NUTRITION AND MANAGEMENT

17	PERFORMANCE AND FINANCIAL BENEFITS OF FINISHER BROILER CHICKENS FED SELECTED COMMERCIAL DIETS SOLD IN BAUCHI METROPOLIS	Zakariya, S. M., Doma, U. D. and Abdullahi, U. S.	1881
18	NUTRIENT DIGESTIBILITY OF BROILER STARTER CHICKS FED PROCESSED BAOBAB SEED MEAL-BASED DIETS SUPPLEMENTED WITH ENZYME	*Parmata ¹ , A., Alu ² , S.E. and Adua ² , M.M.	1885
19	EVALUATION OF GRADED LEVELS OF STARCH AS BINDER IN LIVESTOCK WAFER FORMULATION	Sadiq M.S ¹ , Dakka, J. D ¹ , Bukola O.A ¹ , Lakurbe O.A ¹ Yahaya.Y.A ² , Tijjani .A ³	1890
20	PERFORMANCE AND EGG QUALITY PARAMETERS OF LAYER HENS FED DIET WITH AND WITHOUT VITAMIN E SUPPLEMENTATION	*.Ezenwosu C, Anizoba N.W, Onodugo M.O, Nwoga C C, Osita C.O, Udeh F.U, Onyimonyi, A. E	1895
21	EFFECT OF SODIUM HYDROXIDE TREATED RICE HUSK MEAL WITH ENZYME SUPPLEMENTATION ON THE CARCASS AND ORGAN CHARACTERISTICS OF BROILER FINISHER CHICKENS	Ikpe, J. N.*, Okonkwo, V. N.**, Ahaotu, E. O.***, Oko, E. C.*, Abdullahi, J.* Mbachu, M. U.****	1899
22	PERFORMANCE OF LAYING HENS FED DIETS CONTAINING VARYING ENERGY LEVELS DURING THE LATE LAYING PHASE	M. ABUBAKAR ^{*1} , A. YARI ^{**2} , A. A. SANI ^{**3} , N.T. IBRAHIM ^{***4} And H.M. HARUNA ^{****5}	1904
23	INTERNAL ORGANS CHARACTERISTICS OF BROILER CHICKENS FED DIETS CONTAINING COCOA SEED POWDER DURING DRY SEASON	©Oloko, A.B., ©Mosobalaje, M.A., and ©Adeaga, L. S.	1910

24	CARCASS QUALITY OF NOILER COMPARED TO BROILER AND COCKEREL	* [¥] Mosobalaje, M. A., ^β Adebayo, B. F., [©] Adeogun, J. B., [®] Oloko, A. B., [¥] Ogunniyi, F. B., [¥] Akanbi, T. K. and [®] Adedoyin, A. A.	1914
25	EVALUATION OF GROWTH PERFORMANCE OF NOILERS, BROILERS AND ISA BROWN COCKERELS AS MEAT TYPE CHICKENS	* [¥] Mosobalaje, M. A., ^β Adebayo, B. F., [©] Adeogun, J. B., [®] Adeaga, L. S., [®] Oloko, A. B., [¥] Dasaolu O. O., [¥] Akinwale, D. T.	1919
26	NUTRIENT COMPOSITION OF ZOBO WASTE: A BY-PRODUCT OF ROSELLE (<i>Hibiscus sabdariffa</i>) CALYX	Sudik, S. D.; Lawan, A.; Maidala, A.; Bagudu, I. A. and Mankinde, O. J.	1923
27	GROWTH PERFORMANCE OF BROILER STARTER CHICKENS FED MAIWA MILLET (<i>PENNISETUM GLAUCUM</i> L.) AS A REPLACEMENT FOR DIETARY MAIZE	¹ Tok, C. C., ² Doma, U. D., ² Mancha, Y. P., ² Bello, K. M. and ¹ Ruma, G. A.	1927
28	EFFECTS OF DIFFERENTLY PROCESSED SICKLEPOD (<i>SENNA OBTUSIFOLIA</i>) SEED MEAL BASED DIETS ON GROWTH PERFORMANCE OF BROILER CHICKENS	Isah, A. ¹ ; Ibrahim, R. ² ; Agyo, B. ³ ; Abba, A. Y. ⁴	1931
29	GROWTH PERFORMANCE AND COST BENEFIT ANALYSIS OF BROILER STARTER CHICKENS FED DIETS CONTAINING GRADED LEVELS OF ALOE VERA (<i>Aloe barbadensis</i>) LEAF MEAL	*Olabode, A.D., ⁺ Agu, C.I., *Adetutu, O.I., *Onyishi, P.I., *Olorunfunmilola, N.A. and *Uzoma, C.	1936
30	INFLUENCE OF BLACK PLUM LEAF MEAL ON THE GROWTH PERFORMANCE AND COST BENEFIT ANALYSIS OF STARTER BROILER BIRDS	*Olabode, A.D., ⁺ Agu, C.I., *Adetutu, O.I., *Ebiaku, V., *Eli, E. and *Uzoma, C.	1941
31	PERFORMANCE OF BROILER STARTER FED DIETS CONTAINING SOYA BEAN RESIDUE SBR SUPPLEMENTED WITH ENZYME	Lawan, I. ¹ , Bawa, .G. S. ² , Oimage, J. J. ² and Afolayan, M. ³	1946

32	EFFECT OF PROBIOTICS ON GROWTH PERFORMANCE OF NOILER CHICKS	Tiough, S.M. and Ogwiji, O.	1951
33	EFFECT OF DIFFERENT PROCESSING METHODS ON CHEMICAL COMPOSITION AND ANTI-NUTRITIONAL FACTORS OF KENAF SEED MEAL (<i>HIBISCUS CANNABINUS</i>)	S. Abubakar ¹ and N.B. Rano ²	1956
34	CARCASS CHARACTERISTICS OF BROILER CHICKENS FED DIETARY LEVELS OF KAURA SORGHUM AS REPLACEMENT FOR MAIZE	² Jamilu, H., * ¹ Jama'a, N. A., ¹ Doma, U. D., ³ Zahraddeen, D., ⁴ Sani, A. and ⁵ H. S. Buhari	1961
35	BONE MINERALIZATION CHARACTERISTICS OF RABBITS FED DIETS CONTAINING VARYING LEVELS OF ACHA OFFAL SUPPLEMENTED WITH MAXIGRAIN[®] ENZYME	AZI, J. A. and ALU, S.E.	1965
36	FEED POTENTIALS OF FERMENTED SPENT SORGHUM GRAINS AND SWEET POTATO LEAVES FOR BROILER CHICKENS	Onyimba, I. A.	1970
37	EFFECT OF BACTOFORT INCLUSION IN DIET AS PROBIOTIC ON GROWTH PERFORMANCE AND ABDOMINAL FAT PERCENTAGE OF BROILER CHICKENS	Buba, W., Ojo, O.A. and Shehu, B.M.	1975
38	GROWTH PERFORMANCE OF BROILER CHICKENS FED DIETS CONTAINING FERMENTED POULTRY DROPPINGS	*Rimbut, N.N., Kudu, Y.S., Alabi, O.J., Otu, B.O., Yabagi, A., Fatahi, A.O., & Jalal, A.A	1980
39	THE INFLUENCE OF FLOOR TYPES ON SKIN LESION SCORE OF CROSSBRED WEANED PIGS	¹ Ogunbode, A.A., ² Abegunde, P.T., ³ Faleye, M.A and ⁴ Adejumo, F.A	1985
40	PERFORMANCE AND CARCASS CHARACTERISTICS OF BROILER FINISHER CHICKEN FED DIFFERENT	USMAN, D., FATIMA, M., NASIRU, H. A., MADU, I., and DAUBA, B. D.	1990

	CEREAL FIBER SOURCES.		
41	EFFECT OF DIETARY LEVELS OF WHEAT OFFAL MEAL AND ENZYME ON PERFORMANCE OF BROILER CHICKEN	¹ Okechukwu, S.O., ² Amaefule, K.U. and ¹ Adetutu O. I.	1995
42	EFFECT OF SANDPAPER (<i>Ficus exasperata</i>) LEAF MEAL ON PERFORMANCE, CARCASS AND ORGAN CHARACTERISTICS OF FINISHER BROILERS	Adetutu O. I., * Olabode A. D., Okechukwu S. O., Onyishi P. I. and Opoke M. C.	2000
43	EFFECT OF SANDPAPER (<i>Ficus exasperata</i>) LEAF MEAL ON GUT MORPHOLOGY AND NUTRIENT DIGESTIBILITY OF FINISHER BROILERS	Adetutu O. I., * Olabode A. D., Anounye A. C., Uzoma C. and Orogwu C. S.	2004
44	LIPID PROFILE OF BROILER CHICKENS FED UZIZA LEAF (<i>PIPER GUINEENSE</i>) AND UDA SEED (<i>XYLOPIA AETHIOPICA</i>) SUPPLEMENTED DIETS AS AN ALTERNATIVE TO ANTIBIOTIC-SUPPLEMENTS	Chinwe U. Nwachukwu, Michael C. Orgi and Peace A. Nlemuwa	2008
45	OXIDATIVE STATUS OF BROILER CHICKENS FED UZIZA LEAF (<i>PIPER GUINEENSE</i>) AND UDA SEED (<i>XYLOPIA AETHIOPICA</i>) SUPPLEMENTED DIETS AS AN ALTERNATIVE TO ANTIBIOTIC-SUPPLEMENTS	Chinwe U. Nwachukwu, Genevieve O. Iloghalu and Deborah N. Nwankwo	2014
46	EFFECT OF DIETARY ENZYME AND SUN DRIED CASSAVA PEELS MEAL AS REPLACEMENT FOR WHEAT BRAN ON PERFORMANCE CHARACTERISTICS OF BROILER STARTER	Raji, M.O ¹ , Ogungbenro, S.D ¹ , Adams, T.O ¹ , Ogungbenro, C.F ² , Akinade, H.O ³ , Oso, A.O ⁴ and Bamgbose, A.M ⁴	2020
47	EFFECTS OF DIETARY SILICON OXIDE LEVELS ON GROWTH PERFORMANCE AND FEET WELFARE OF	*Opoola, E., **Lawal, A. N and **Oyebanji, O. A.	2024

	BROILER CHICKS		
48	EFFECT OF GRADED LEVELS OF PROCESSED NEEM (<i>Azadirachta indica</i>) LEAF AS FEED ADDITIVE ON GROWTH PERFORMANCE AND CARCASS TRAITS OF BROILER BIRDS	Salisu, U. S., Gaddafi, S., Garba, M. G., Yahaya, M. A., Bara`u, A., Sanusi, A.Z and Sabo, M. N	2029
49	CARCASS CHARACTERISTICS AND SENSORY EVALUATION OF BROILER CHICKENS FED DIETS CONTAINING PHYTOBIOTICS	Shobiye, O.E.,*Irivboje, Y.I., Sobayo, R.A.	2034
50	NUTRIENT DIGESTIBILITY AND ECONOMIC BENEFIT ANALYSIS OF JAPANESE QUAILS (<i>Coturnix coturnix japonica</i>) FED DIETS CONTAINING GRADED LEVELS OF GINGER (<i>Zingiber officinale</i>) WASTE MEAL	Danladi, M. J.*; Ijaiya, A. T., Malik, A. A. and Asoloko, M. A.	2039
51	REPLACEMENT VALUE OF WHEAT OFFAL WITH ONION (<i>Allium cepa</i> L.) SCALES IN BROILER CHICKEN DIETS	Oshilaja, A. A., Muftau, M. A., Ribah, M. I. and Ibrahim, N. T.	2044
52	PROXIMATE AND PHYTOCHEMICAL COMPOSITIONS OF DIFFERENTLY PROCESSED SWEET POTATO (<i>Ipomoea batatas</i>) PEEL MEAL	*Yisa, A. G. and Daniel, C.	2048
53	EFFECTS OF VARYING LEVELS OF COOKED KIDNEY BEAN MEAL (<i>Phaseolus Vulgaris</i>) AS REPLACEMENT OF SOYABEAN MEAL OR GROUNDNUT CAKE ON THE PERFORMANCE OF BROILER CHICKEN	¹ *Shuaibu, M. D., ² Aremu, A.	2053
54	ECONOMICS OF PRODUCTION OF STARTER BROILER CHICKENS FED GRADED LEVELS OF MULTIGRAIN ENZYME-	Jimoh, A.	2058

	SUPPLEMENTED RICE HUSK AS A REPLACEMENT FOR MAIZE		
55	MEASUREMENT OF GROWTH PERFORMANCE INDICES IN FINISHER BROILERS FED DIETS CONTAINING EQUI-PROTEIN REPLACEMENT OF SOYABEAN MEAL WITH <i>MORINGA OLEIFERA</i> LEAF MEAL	Sule, K. ^{1*} , Onibi, G.E. ² Folorunso, O.R. ¹ and Olowu O.P.A. ¹	2063
56	PERFORMANCE AND ECONOMICS OF FINISHER BROILER CHICKENS FED DIFFERENT COMMERCIAL FEEDS	¹ Sani, A., ¹ Shuaibu, U. S., ² Jamaa, N. A., ¹ Mijinyawa, A., ⁴ Garba, A., ³ Jamilu, H., ¹ Abubakar, A., ¹ Nuhu, S. H., ¹ Oyeniran, J. O. and ² Doma, U. D.	2068
57	GROWTH PERFORMANCE OF STARTER BROILER CHICKENS FED DIFFERENT COMMERCIAL DIETS	¹ Sani, A., ¹ Mijinyawa, A., ² Jamaa, N. A., ¹ Abubakar, A., ³ Jamilu, H., ¹ Nuhu, S. H., ¹ Oyeniran, J. O. and ² Doma, U. D.	2072
58	CARCASS AND ORGAN CHARACTERISTICS OF BROILER CHICKENS FED DIFFERENT COMMERCIAL FEEDS.	¹ Sani, A., ¹ Shuaibu, U. S., ⁵ Kawu, Y. U., ² Jamaa, N. A., ¹ Mijinyawa, A., ⁴ Garba, A., ³ Jamilu, H., ¹ Abubakar, A., ¹ Nuhu, S. H., ¹ Oyeniran, J. O. and ² Doma, U. D.	2076
59	POTENTIAL OF MIXED FORAGES (<i>CALOPONUM MUCOIDS</i> AND <i>PANICUM MAXIMUM</i>) ON THE INTERNAL ORGAN CHARACTERISTICS OF PIGS	Uzegbu, H. O., Ewa, U. E. and Ukonu, E. C.	2080
60	EVALUATION OF THE POTENTIAL OF <i>PANICUM MAXIMUM</i> FORAGE ON GUT MICROBIAL POPULATION OF PIGS	Uzegbu, H. O. and Ukonu, E.C.	2084
61	EVALUATION OF THE POTENTIAL OF <i>PANICUM MAXIMUM</i> FORAGE ON GUT MICROBIAL POPULATION OF PIGS	Uzegbu, H. O. and Ukonu, E.C.	2089
62	GROWTH PERFORMANCE AND CARCASS EVALUATION OF BROILER CHICKENS FED FIVE DIFFERENT COMMERCIAL DIETS	Bello, A. R., Doma, U. D. and Kalla, D. J. U.	2094

63	EFFECT OF DIETARY UTILIZATION OF ONION AND GARLIC ON PERFORMANCE AND ORGANS WEIGHT OF BROILER CHICKS AT FINISHER PHASE	*Abdullahi, A.Y. [±] , *Bala, A. *Zango, M.H., *Madaki S., *Nasir, M., *Khaleel, A.G. [∞] Dan Abba, Y. U., [∞] Bichi, A. A., ⁺ Abubakar A.	2098
64	GROWTH PERFORMANCE OF BROILER CHICKENS FED VARIOUS DIETARY SOURCES OF ENERGY AND PROTEIN	¹ Al-Mustapha, A.J., ² Doma, U.D., ² Abubakar, M.M., ³ Jafaru, S. and ⁴ Magaji, I.M.	2103
65	EFFECT OF FEEDING TIME ON THE GROWTH AND MORPHOLOGICAL TRAITS OF ISA BROWN PULLETS	*Abe, O.S. ⁺ , Ayankoso, M.T. ⁺ and Oseyili B.G. [*]	2108
66	EFFECTS OF VARYING DURATION OF DIFFERENT PROCESSING METHODS ON NUTRIENT QUALITIES AND TOXICITY (ANTI-NUTRITIONAL FACTORS) OF NEGRO COFFEE (<i>SENNA OCCIDENTALIS</i>) SEEDS MEAL	Ibrahim*, M. J., Tsado, D.N., Adama, T.Z., Adama, J.Y., Muhammad, M.L. and Yohana, A.	2113
67	DIETARY EFFECT OF THE SUPPLEMENTATION OF PROBIOTICS (<i>Lactobacillus fermentum</i>) ON CARCASS CHARACTERISTICS OF BROILER CHICKENS	Akande, K.E., Alabi, O.J. and Jatau, L.	2117
68	BODY WEIGHTS OF DOMESTIC PIGEON SQUABS (<i>COLUMBA LIVIA</i>) BY SEX AND PARITY IN MAIDUGURI, BORNO STATE, NIGERIA.	Zannah, B. B., Aliyu, J., Yusuf, M. J., Taki, I. M., Aminami, M. and Alade, N. K.	2121
69	L-THREONINE SUPPLEMENTATION IN LOW CRUDE PROTEIN DIET OF BROILER STARTER CHICKEN	Aguihe P.C. ^{*+} , Ilaboya I.I. [¥] , Samuel K.U. [€] , Joshua D.A. [§] and Abdulwahab S. [*] .	2126
70	RESPONSE OF BROILER CHICKENS FED LOW CRUDE PROTEIN, LOCAL GROUNDNUT CAKE BASED DIETS WITH THREONINE SUPPLEMENTATION	Aguihe P.C. ^{*+} , Kehinde A.S, Samuel K.U. [€] , Abdulwahab S. [*] and D.A. Joshua [¥]	2131
71	COMPARISON OF CHEMICAL COMPOSITION	Ganiyu, A.O. and Belewu, M. A.	2136



NSAP

47th Annual Conference
(JOS 2022)

CONFERENCE PROCEEDINGS

THEMES
SECURING ANIMAL AGRICULTURE AMIDST GLOBAL CHALLENGES

DIETARY EFFECT OF THE SUPPLEMENTATION OF PROBIOTICS (*Lactobacillus fermentum*) ON CARCASS CHARACTERISTICS OF BROILER CHICKENS

Akande, K.E., Alabi, O.J. and Jatau, L.

Department of Animal Production, Federal University of Technology Minna, Niger State, Nigeria.

Email:kemi777akande@gmail.com

ABSTRACT

One hundred day old Cobb 500 broiler chickens were randomly assigned to five treatments in a completely randomized design to study the effect of dietary probiotics (*Lactobacillus fermentum*) on carcass characteristics. Treatment 1 served as the control without probiotics (0.000g) while treatments 2, 3, 4, and 5 were diets supplemented with 1.875, 3.750, 5.625 and 7.500 g of probiotics per kg of feed respectively. At the end of the experiment, four birds per treatment were randomly selected for carcass evaluation. The final live body weight was recorded before slaughter. After evisceration, the carcass weight was recorded, and the weights of primal parts, internal organs and abdominal fat were all taken and expressed as a percentage of carcass weight. Data collected were subjected to one-way analysis of variance. The results obtained showed that final live weight, carcass weight, breast weight, thigh weight and abdominal fat were significantly affected ($P < 0.05$) by dietary treatments. It was observed that birds on T5 fed probiotics of 7.500 g/kg feed had a live weight of 3050.00g which was similar to T4 but significantly higher than all the other treatments. Treatment 4 had the highest carcass weight which was followed by treatment 5. Treatment 5 birds recorded the highest ($P > 0.05$) breast and thigh weight when compared to birds in other treatment groups. The birds in the control group (T1) recorded the highest ($P > 0.05$) abdominal fat than other treatments. This invariably implies that the supplementation of this probiotic could reduce abdominal fat deposits in birds.

Keywords: Probiotics, *Lactobacillus fermentum*, carcass characteristics, Cobb 500 broilers, abdominal fat

INTRODUCTION

Research studies have shown that antibiotics improve the performance of broiler chickens however, their usage has been banned in some countries, because of their harmful side effects on broiler birds and their residual effect on meat (Mehdi *et al.*, 2018). The need, therefore, arises to find a safe alternative additive for use in poultry production. Using probiotics is believed to be safer and promotes health. The utilization of probiotics in poultry production may address the rising public health issues on meat quality and antimicrobial resistance development to some level, by substituting the use of sub-therapeutic antibiotics for growth promotion with probiotics (Jha *et al.*, 2020). Probiotics are beneficial live microbes (bacteria, yeast or fungi) that supplement the gut flora enhancing a healthy digestive system when administered in appropriate dosage (FAO/WHO, 2006). According to Siragusa (2012) probiotics for livestock are referred to as direct-fed microbial. Probiotics can be given to poultry through feed or water or can be administered to a developing embryo through in ovo feeding technology (Pender *et al.*, 2017). However, there are few research studies on the dietary effect of probiotics on the carcass characteristics of broiler chickens. Hence, this study is aimed at evaluating the effect of dietary probiotics (*Lactobacillus fermentum*) on carcass characteristics of broiler chickens.

MATERIALS AND METHODS

This experiment was carried out at the Federal University of Technology Minna, Bosso Campus, Niger State, at the Department of Animal Production Teaching and Research Farm (Poultry Unit). A total of one hundred Cobb-500 broilers, at day old were used for this study. The probiotic used was *Lactobacillus fermentum*. A completely randomized design (CRD) was used in the experiment. The birds were weighed, and then randomly allotted into five treatments. Treatment 1 served as the control



NSAP

47th Annual Conference
(JOS 2022)

CONFERENCE PROCEEDINGS

THEMES
SECURING ANIMAL AGRICULTURE AMIDST GLOBAL CHALLENGES

without probiotics supplement (0.000g) while treatments 2, 3, 4, and 5 were diets supplemented with 1.875, 3.750, 5.625 and 7.500g of probiotics (*Lactobacillus fermentum*) per 1kg of feed respectively. The birds were raised in a deep litter system. Before the arrival of the birds, the pen was washed and disinfected, wood shavings was evenly spread on the floor. Upon the arrival of the chicks, they were weighed to obtain their initial weights and after which they were randomly distributed to the various treatment groups. The chicks were administered anti-stress (vitalyte) through drinking water according to the manufacturer's instruction. The birds were vaccinated against gumboro disease at the end of the first week. This was repeated at 2 weeks of age. In the fourth and eighth week, Newcastle vaccine (NCV lasota) was given to the birds. The experiment was conducted for eight (8) weeks. Feed intake was measured daily while the weighing of birds was done on weekly basis. Iso-caloric and iso-nitrogenous single-phase diets were formulated for the broilers. Feed and water were provided *ad libitum*.

Evaluation of carcass characteristics

At the end of the experiment, four birds per treatment were randomly selected for carcass evaluation. The final live weight was recorded before slaughter. Birds were weighed, starved overnight to clear the gut but allowed access to fresh clean water before slaughter. The birds were slaughtered by cutting the jugular vein with a sharp knife. After evisceration, the carcass weight and the weights of internal organs (liver, gizzard, lungs, heart, spleen) and abdominal fat were all taken and expressed as a percentage of carcass weight. Primal parts like the wing, thigh, breast and drumstick were weighed and expressed as a percentage of the carcass weight. All data obtained were subjected to One-Way Analysis of Variance (ANOVA) and significant treatment means where it occurred were separated by Duncan multiple range test using (SPSS, 2017) version 16.0.

RESULTS AND DISCUSSION

Table 1 shows the result of the effect of dietary *Lactobacillus fermentum* probiotics supplementation on carcass characteristics of Cobb 500 broiler chickens. The results from this research showed that there were significant differences ($P < 0.05$) in the final live weight, carcass weight, breast meat weight, thigh weight and abdominal fat of broilers. This might be as a result of the improved gut microflora as highlighted by Markowiak and Ślizewska, (2017) and Jha *et al.* (2020). However, there was no significant differences ($P > 0.05$) observed in the weight of drumstick, wing, heart, liver, lungs, gizzard and spleen. The birds on T4 and T5 had similar final live weight which was higher than ($P > 0.05$) the other treatments. Birds on T4 had the best ($P > 0.05$) carcass weight which was followed by T5. Treatment 5 birds recorded the highest ($P > 0.05$) breast and thigh weight when compared to birds in other treatment groups. The supplementation of the probiotic level of 7.500g/kg of feed favoured both breast and thigh weight. Tang *et al.* (2021) reported a significant increase in breast muscles of broiler chickens with probiotic supplementation which is consistent with the result of increased breast weight of birds on *Lactobacillus fermentum* probiotics treatment in this research. The birds in the control group (T1) recorded the highest ($P > 0.05$) abdominal fat than other treatments. This invariably implies that the supplementation of this probiotic was capable of reducing abdominal fat deposits in birds, thereby improving meat quality and higher feed and economic efficiency. The significant decrease in abdominal fat of broilers fed diets with probiotics obtained in this study is in line with the report of Tang *et al.* (2021) which stated that the supplementation of probiotics (*Bacillus subtilis*) led to a significant decrease in abdominal fat of broilers when compared with broilers without probiotics. Additionally, this result also agrees with the findings of Hidayat *et al.* (2016) who reported that the increase in the level of use of probiotics (*Bacillus spp*) tends to decrease the percentage of abdominal fat. Excessive abdominal fat in chicken meat is highly undesirable, most consumers prefer chicken meat with low-fat content for economic and health reasons.

**NSAP****47th Annual Conference**
(JOS 2022)**CONFERENCE PROCEEDINGS**THEMES
SECURING ANIMAL AGRICULTURE AMIDST GLOBAL CHALLENGES

Table 1 Dietary effect

of Lactobacillus fermentum probiotics on carcass characteristics of broiler chickens

Parameter	T1	T2	T3	T4	T5	SEM	P-value
Final live weight (g)	2963.00 ^b	2935.50 ^c	2892.00 ^d	3049.50 ^{ab}	3050.00 ^a	21.789	0.01
Carcass weight	2214.00 ^c	2200.00 ^d	2171.00 ^e	2323.00 ^a	2295.50 ^b	19.471	0.00
Breast weight (%)	37.60 ^c	37.86 ^b	36.82 ^e	37.25 ^d	39.49 ^a	0.317	0.01
Thigh weight (%)	16.23 ^d	17.52 ^b	16.77 ^c	17.49 ^b	18.56 ^a	0.282	0.02
Drumstick (%)	12.78	11.53	12.14	11.96	11.94	0.174	0.25
Wings (%)	10.00	10.05	10.06	9.85	9.43	0.121	0.52
Heart (%)	0.47	0.42	0.36	0.41	0.38	0.017	0.33
Liver (%)	1.65	1.75	1.86	1.46	1.74	0.061	0.32
Lungs (%)	0.13	0.17	0.12	0.11	0.11	0.010	0.29
Gizzard (%)	2.38	2.52	2.33	2.21	2.47	0.061	0.66
Spleen (%)	0.40	0.40	0.34	0.44	0.52	0.025	0.25
Abdominal fat (%)	1.52 ^a	0.37 ^d	0.41 ^c	0.82 ^b	0.82 ^b	0.148	0.02

a, b, c, d, e Means with different superscripts on the same row are significantly different (P>0.05).

T1: Without *Lactobacillus fermentum* probiotics 0.00g/kg of feed (control)

T2: *Lactobacillus fermentum* probiotics of 1.875g/kg of feed

T3: *Lactobacillus fermentum* probiotics of 3.750g/kg of feed

T4: *Lactobacillus fermentum* probiotics of 5.625g/kg of feed

T5: *Lactobacillus fermentum* probiotics of 7.500g/kg of feed

SEM: Standard Error of Mean

CONCLUSION AND RECOMMENDATION

Results from this study have shown that *Lactobacillus fermentum* probiotics can serve as a growth promoter in the production of broiler chickens. The supplementation of the probiotic level of 7.500g/kg of feed is therefore recommended for improved live weight, carcass weight, breast weight and thigh weight. Further research study can be carried out using these probiotics as a dietary supplement for other poultry species.

REFERENCES

- FAO/WHO (2006). Probiotics in food: Health and nutritional properties and guidelines for evaluation. *Food and Agriculture Organisation of the United Nations; Rome, Italy: World Health Organisation; Geneva, Switzerland: FAO Food and Nutrition and Paper.*
- Hidayat, M. N., Malaka, R., Agustina, L., & Pakiding, W. (2016). Abdominal fat percentage and carcass quality of broiler given probiotics *Bacillus spp.* *Scientific Research Journal* 4(10) 33-37.
- Jha, R., Das, R., Oak, S., & Mishra, P. (2020). Probiotics (direct-fed microbials) in poultry nutrition and their effects on nutrient utilization, growth and laying performance, and gut health: a systematic review. *Animals*, 10(10), 1863, 1-23.
- Markowiak, P. & Ślizewska K. (2017). Effects of probiotics, prebiotics, and synbiotics on human health. *Nutrients*, 9(9), 1021-1051.



NSAP

47th Annual Conference
(JOS 2022)

CONFERENCE PROCEEDINGS

THEMES
SECURING ANIMAL
AGRICULTURE AMIDST
GLOBAL CHALLENGES

Mehdi, Y., Létourneau-Montminy, M.P., Gaucher, M.L., Chorfi, Y., Suresh, G., Rouissi, T., Brar, S.K., Côté, C., Ramirez, A.A. & Godbout, S. (2018). Use of antibiotics in broiler production: Global impacts and alternatives. *Animal Nutrition*, 4(2), 170-178.

Pender, C. M., Kim, S., Potter, T. D., Ritzi, M. M., Young, M., & Dalloul, R. A. (2017). In ovo supplementation of probiotics and its effects on performance and immune-related gene expression in broiler chicks. *Poultry Science*, 96(5), 1052-1062.

Siragusa, G. R. (2012). Modern probiology direct-fed microbials and the avian gut microbiota. In 23rd *Annual Australian Poultry Science Symposium*, 120-130.

Tang, X., Liu, X., & Liu, H. (2021). Effects of dietary probiotic (*Bacillus subtilis*) supplementation on carcass traits, meat quality, amino acid, and fatty acid profile of broiler chickens. *Frontiers in Veterinary Science*, 8.1-10