

Proceedings of the

2nd

International Conference of Agriculture and Agricultural Technology

ICAAT 2022



Theme:

Climate-Smart Agriculture in the Post
COVID Era:
A Gate Way to Food Security in Africa

Held at
Caverton Hall
Federal University of Technology Minna, Nigeria

Published by
School of Agriculture and Agricultural technology
Federal University of Technology
P. O. M. 65, Minna, Nigeria
Email: icaat@futminna.edu.ng
Tel: 08028573305

Edited by

Dr. (Mrs) Adeniran, O.A.
Dr. Muhammad, H.U.
Dr Rajan Sharma

Prof. Alabi, O.J.
Dr. (Mrs) Akande, K.E.
Dr. Otu, B.O.

Dr. Mrs Carolyn Cherotich
Mr. Ibrahim, A.
Mr Adesina, O.A



Contents

1	SUSTAINABLE CROP PRODUCTION PRACTICES FOR CLIMATE RESILIENCE FOOD AND NUTRITION SECURITY	3
2	ADOPTION INDEX OF MAIZE PRODUCTION TECHNOLOGIES AND CORRELATION MATRIX IN SMALLHOLDER SYSTEMS	9
3	PROXIMATE AND SENSORY ASSESSMENT OF AADUN PRODUCED FROM MAIZE AND GROUNDNUT PASTE	15
4	RURAL WOMEN AND AGRO – PROCESSING: A CASE STUDY OF RURAL WOMEN PARTICIPATION IN GROUNDNUT PROCESSING IN KATSINA STATE	22
5	PROFITABILITY OF DRIP IRRIGATED MAIZE PRODUCTION FOR IMPROVED FOOD SECURITY IN MAIDUGURI SEMI-ARID REGION OF BORNO STATE, NIGERIA	28
6	PROFITABILITY ANALYSIS OF POULTRY EGG PRODUCTION IN IBADAN METROPOLIS, OYO STATE, NIGERIA	37
7	ADOPTION OF MODERN BEEKEEPING TECHNOLOGIES IN SELECTED LOCAL GOVERNMENT AREAS OF BENUE STATE, NIGERIA	44
8	ADOPTION OF RECOMMENDED COCOYAM PRODUCTION TECHNOLOGIES AMONG FARMERS IN ENUGU STATE, NIGERIA	53
9	INNOVATIVE APPLICATION IN MANAGEMENT OF PROBLEMATIC SOIL (ACIDIC SOIL) UNDER MAIZE PRODUCTION IN NIGER STATE, NIGERIA	62
10	EFFECTS OF RISK MANAGEMENT STRATEGIES ON	69
11	POVERTY STATUS OF RICE FARMERS IN NIGER STATE, NIGERIA	69
12	EFFECTS OF FARMER-HERDER'S CONFLICT ON THE FARMERS PRODUCTIVITY IN ADAMAWA STATE, NIGERIA	75
13	EFFECTS OF ADOPTION OF IMPROVED BEEHIVE TECHNOLOGIES ON INCOME AND WELFARE STATUS OF BEEKEEPERS IN EKITI STATE, NIGERIA	81
14	ASSESSMENT OF OUT-MIGRATION AMONG ARABLE CROP FARMERS IN KOGI STATE, NIGERIA: GENDER DYNAMIC APPROACH	88
15	THE EFFECT OF AGRITECH STARTUPS ON PARTICIPATION AND POVERTY STATUS OF ADOPTERS IN OGUN STATE, NIGERIA	97
16	OUTCOMES OF WORK-FAMILY CONFLICT AMONG UNIVERSITY EMPLOYEES: CORRELATES OF TIME, STRAIN AND JOB SATISFACTION	104
17	MICROBIOLOGICAL QUALITY OF OVEN ROASTED PLANTAIN (MUSA PARASIDIACA)	109

	SAVANNAH	620
95	EFFECT OF FOLIAR FERTILIZER APPLICATION ON PHYSIOLOGICAL CHARACTERISTICS AND HERBAGE YIELD OF AMARANTHUS AND CORCHORUS	626
96	EFFECT OF FOLIAR FERTILIZER APPLICATION ON PHYSIOLOGICAL CHARACTERISTICS AND HERBAGE YIELD OF AMARANTHUS AND CORCHORUS	631
97	RESPONSE OF SESAME (<i>SESAMUM INDICUM L.</i>) TO DIFFERENT NUTRIENT SOURCES IN ANYIGBA, KOGI STATE NIGERIA	636
98	EFFECTS OF INTEGRATED NUTRIENT MANAGEMENT ON WEED INFESTATION AND YIELD OF SOYBEAN IN SOUTHERN GUINEA SAVANNA ZONE OF NIGERIA	648
99	EFFECT OF MILLED GROUNDNUT SHELL AS A NUTRIENT SOURCE ON THE NUTRITIONAL QUALITY OF PEPPER IN COMPARISON WITH OTHER NUTRIENT SOURCES	656
100	REACTION OF RICE VARIETIES TO RICE BLAST - AN INSIGHT INTO UNDERSTANDING OF RICE RESILIENCE TO CLIMATE INDUCED RICE DISEASES	664
101	ALLELOPATHIC EFFECT OF MINTWEED (<i>HYPTIS SUAVEOLENS (L.) POIT</i>) GREEN AND BROWN LEAVES AQUEOUS EXTRACT ON SEED GERMINATION AND SEEDLING GROWTH OF COWPEA	676
102	PROXIMATE COMPOSITION OF SOME NEW SWEET POTATO VARIETES GROWN AT GIDAN KWANO, MINNA, NIGER STATE, NIGERIA	682
103	EVALUATION OF <i>SACCHARUM OFFICINARUM</i> GENOTYPES FOR JUICE QUALITY, CANE AND SUGAR YIELD	690
104	IMPACTS OF SUGARCANE GENOTYPES AND WEED MANAGEMENT PRACTICES ON WEED DRY MATTER AND SUGARCANE PRODUCTIVITY	696
105	ISIGHTS INTO PREVALENC AND DISTRIBUTION OF VIRUSES INFECTING MELON IN GEORGIA, UNITED STATE	703
106	GROWTH AND YIELD RESPONSES OF MAIZE TO PARTIAL SUBSTITUTION OF INORGANIC NITROGEN WITH FARM YARD MANURE AT GIDAN-KWANO	704
107	EFFECT OF BIOCHAR AND NANOPARTICLES ON THE MICROBIAL DEGRADATION OF CRUDE OIL CONTAMINATED SOIL	713
108	INDIGENOUS KNOWLEDGE AND COPING CAPACITY OF FARMERS FOR CLIMATE CHANGE ADAPTATION: A CASE STUDY OF AGAIE LOCAL GOVERNMENT AREA, NIGER STATE. NIGERIA	721
109	ASSESSMENT OF FERTILITY STATUS OF SOME SELECTED FADAMA SOILS IN BIDA LOCAL	

	GOVERNMENT AREA OF NIGER STATE, NIGERIA	725
110	EFFECTS OF DIFFERENT NITROGEN SOURCES ON THE GROWTH OF MAIZE AND SOYBEAN IN AN INTERCROPPED SYSTEM	731
111	EVALUATION OF IRRIGATION SUITABILITY FOR SURFACE IRRIGATION IN BAKAJEBA, NIGER STATE, NIGERIA	738
112	EFFECTS OF BIOCHAR AND NITROGEN FERTILIZER ON SOIL ORGANIC CARBON, NUTRIENT RETENTION AND MAIZE PERFORMANCE	744
113	IMPLICATION OF LAND SURFACE TEMPERATURE AS A DRIVER OF LAND COVER CHANGES ON THE URBAN DYNAMICS OF ABUJA NIGERIA	751
114	PHOSPHATE MOBILIZATION BY ADDITIONS OF THREE ANIMAL MANURE SOURCES IN SOILS OF KARU, NASARAWA STATE	758
115	RESPONSE OF GROWTH AND YIELD OF MAIZE (<i>ZEA MAYS</i> L.) TO LIME, INORGANIC AND ORGANIC FERTILIZER IN MOKWA, NIGER STATE OF NIGERIA	763
116	UNDERSTANDING THE INTENTION TO USE GOOD AGRICULTURAL PRACTICES ON VEGETABLE FARMS – A COMPARATIVE STUDY OF FARMERS IN PUNJAB, INDIA AND NAKURU, KENYA	771
117	SENSORY AND PROXIMATE COMPOSITION OF 'BISCUIT' PRODUCED FROM SOME LESS VALUED DRIED FISH POWDER	778
118	COMBINATION OF ORGANIC MANURE, INORGANIC FERTILIZER AND BIO-FERTILIZER ON SOYBEAN [(<i>GLYCINE MAX</i> L. (MERRIL)] GROWTH	785
119	GENDER ANALYSIS OF FARMING HOUSEHOLDS' ACCESS TO LIVELIHOOD RESOURCES IN SELECTED LOCAL GOVERNMENT AREAS OF NIGER STATE, NIGERIA	793
120	EFFECTS OF PHOTOPERIOD AND FEEDING RATE ON THE GROWTH PERFORMANCE AND FEED UTILIZATION OF <i>CLARIAS GARIEPINUS</i>	802

ALLELOPATHIC EFFECT OF MINTWEED (*Hyptis suaveolens* (L.) Poit)
GREEN AND BROWN LEAVES AQUEOUS EXTRACT ON SEED
GERMINATION AND SEEDLING GROWTH OF COWPEA

Aguwa, N., Daniya, E. and Mamudu, A. Y.

Department of Crop Production, School of Agriculture and Agricultural Technology,
Federal University of Technology, P.M.B 65, Minna, Niger State, Nigeria.

*Corresponding Author's Email: nnamdiaguwa4eck@gmail.com

Mobile: 07063056463

ABSTRACT

Hyptis suaveolens is an invasive weed that has negative effects on native plant communities and impedes the growth of vulnerable or threatened species growing near it. This study revealed the allelopathic effect of the aqueous extract of the green and brown leaves of *Hyptis suaveolens* on seed germination and seedling growth of cowpea at different concentration (0 %, 2.5 %, 5 %, 7.5 % and 10 %) in a laboratory – based experiment. The treatments were arranged as a 2 x 5 factorial in a Completely Randomized Design (CRD) in four replications. Results revealed that irrespective of the leaf type, there was no significant difference in the allelopathic potential on the seed germination and seedling growth of cowpea. In contrast, the concentration of the extracts had a significant allelopathic effect on cowpea seed germination and seedling growth, such that the inhibition was concentration dependent. The inhibition of the seed germination and seedling growth increased with the increase in the concentration from 0 to 10 %. The highest concentration (10%) of *H. suaveolens* leaf extract significantly reduced the germination percentage, germination index, mean germination time, radicle and plumule lengths by 61.5 %, 63.9 %, 61.7 %, 54.2 % and 62.5 %, respectively. It was apparent that the 10% concentration had more inhibitory (-61.5%) effect on the germination and seedling growth of cowpea relative to the control. This weed must therefore be controlled in the cowpea field at their initial growth.

Keywords: Allelopathy, Germination, Seedling growth, Concentration and Cowpea

INTRODUCTION

Hyptis suaveolens (L.) Poit, which is commonly known as Mintweed, is an important invader of the tropical and sub-tropical regions of the world (Afreen *et al.*, 2017). *Hyptis* possesses several characteristics common to other invasive species, such as allelochemicals that help in its survival and spread. This species has wide ecological amplitude, high plasticity and reproductive capacity because it is able to grow on a variety of soil types, land uses and land cover type (Padalia *et al.*, 2013). *Hyptis* possesses faster growth rate, massive seed production, high proliferation rate, dual mode of reproduction (from perennating root-stocks as well as seed). Higher resistance against pathogens due to allelochemicals and essential oil (Raizada, 2006).

The growth and establishment of other plant species near the clumps of *H. suaveolens* is quite restricted (Raizada *et al.*, 2006) but the specific reasons that lead to the dominance of *H. suaveolens* still remain unclear. One of the plausible reasons for such interference could be due to the presence of allelochemicals in this plant. Allelochemical activities of aqueous methanol extract of *H. suaveolens* on the germination and seedling growth of several weeds and crop species have been reported by Momimul-Islam *et al.* (2013) and Daniya *et al.* (2014).

Allelopathy refers to the effect of one plant species on another through the release of chemical compounds known as allelochemicals into the environment (Weih *et al.*, 2008), which may either be promoting (synergistic) or inhibiting (antagonistic) depending on the compounds released and target plants. The potential use of allelopathy as a natural means for weed suppression in Agro ecosystems has attracted the interest of researchers for a long time (Yongqing *et al.*, 2014).

Cowpea (*Vigna unguiculata* (L.) Walp) is an important pulse crop widely cultivated in the tropics. The pods are highly nutritive and are a good source of digestible protein, dietary fiber and vitamin A and C. In addition to these the pods also contain Ca, P, Na, K, Mg, Fe, Zn, Mn and Cu (Gerrano *et al.*, 2017). However, it is sensitive to weed competition especially at the early stage of crop development and the period from 14 to 40 days after sowing (DAS) has been considered critical for the weed competition (Osipitan *et al.*, 2016). Weed infestation has been reported to be more severe during the rainy season and causes severe yield reduction (Gupta *et al.*, 2016). The impact of weed interference on cowpea yield depends on the duration and stage at which the crop-weed interference takes place. The season-long competition has resulted in 53 to 76 percent yield reduction in cowpea (Gupta *et al.*, 2016).

The aim of the study was to assess the allelopathic effect of aqueous extracts of green and brown leaves of *H. suaveolens* at varying concentrations on germination and seedling growth of cowpea.

MATERIALS AND METHODS

A set of experiments was carried out to assess the allelopathic effect of *H. suaveolens* on germination and seedling growth of cowpea in a laboratory at Department of Crop Production, Federal University of Technology, Minna, Niger State. Green and Brown leaves of *H. suaveolens* were collected at flowering stage from farmlands within the Federal University of Technology Gidan Kwano Campus, Minna, Niger State during the rainy season of 2022. The two leaves were

separated, air dried under the shade for four weeks and later oven dried at 70°C to a constant weight. The powder of each plant part was soaked in distilled water for 24 hours in a ratio of 1:20 (w/v) at room temperature. The water extracts of each plant part were filtered through a Whatman No. 1 filter paper to obtain the pure aqueous extracts without impurities. The extracts prepared in this manner were used as stock solutions with concentration of 10 % (w/v), and further dilutions to 2.5 % (w/v) of extracts of green and brown leaves of *H. suaveolens* were made by adding distilled water to the stock solution for carrying out the experiment at varying concentrations: - 0%, 2.5% (1:40 w/v), 5% (1:20 w/v), 7.5% (1: 15 w/v) and 10% (1: 10 w/v) against cowpea in a laboratory condition. Ten cowpea seeds were placed on a filter paper in 9 cm diameter petri dishes, separately. In each petri dish, 10 ml of green or brown leave aqueous extracts at 2.5%, 5%, 7.5% and 10% and distilled water was poured as per the treatments. Distilled water was used as control for comparison. The temperature during the growing period of cowpea ranged from 29.0 to 31.8°C. The experiment was observed for 16 days. Petri dishes were kept moistened during the whole period of study. The number of seeds germinated was counted on a daily basis. Germination percentage, germination index, mean germination time and germination energy were determined using the formulae shown below: -

(a) Germination percentage (GP)

$$GP = \frac{\text{Germinated seeds}}{\text{Total seeds}} \times 100 \text{ as described by the Association of Official Seed Analysts (AOSA, 1990).}$$

(b) Mean germination time (MGT)

$$MGT = \frac{\sum Dn}{N} \text{ Ellis Roberts as described by Safder et al. (2021)}$$

Where n = number of germinated/emerged seeds on day 'D' and 'D' equal days of count

(c) Germination Index (GI)

$$GI = \frac{\text{No of germinated seeds}}{\text{Days of first count}} + \frac{\text{No of germinated seeds}}{\text{Days of first count}} \text{ as described by AOSA (1990)}$$

(d) Radicle and Plumule length: This were measured using a centimetre rule.

(e) Percentage Inhibition/Stimulation: -

$$\text{Inhibition (-) or Stimulation (+)} = \frac{\text{Germinated seeds in extracts}}{\text{Germinated seeds in control}} - \text{Germinated seed in control} \times$$

100

The data collected were subjected to the analysis of variance (ANOVA) using the Statistical Analysis System (SAS) software package version 9.0. Differences between the treatments means were separated by the Least Significant Difference (LSD) at 5% level of probability.

RESULTS AND DISCUSSION

Germination Indices of Cowpea

The allelopathic effect of green and brown leaves extracts of *H. suaveolens* on germination indices of cowpea are shown in Table 1. The germination percentage, germination index and mean germination time was not significantly affected by either green or brown leaves of *H. suaveolens* (Table 1). In contrast, the germination indices were significantly reduced by the concentrations of the aqueous extract of the leaves (Table 1). Furthermore, the germination percentage, germination index and mean germination time were significantly reduced as the concentration of the extract was increased from 0% to 10%. The highest concentration of 10% consistently and significantly reduced the germination percentage, germination index and mean germination time compared to the control. The allelopathic inhibition of germination in cowpea could be attributed to the presence of allelochemicals at varying concentrations in the water extracts of the *H. suaveolens* leaves. Oraon and Mondal (2021) demonstrated that *H. suaveolens* inhibited the seed germination, though in rice. Furthermore, Oraon and Mondal (2021) substantiated that allelopathic activity depends on the concentration and varies from species to species.

Seedling Growth of Cowpea

The allelopathic effect of extracts of *H. suaveolens* on seedling growth of cowpea is shown in Table 2. The radicle and plumule lengths were not significantly affected by either green or brown leaves of *H. suaveolens*. In contrast, the radicle and plumule length were significantly reduced by the concentrations of the aqueous extract of the *H. suaveolens* (Table 2). Both radicle and plumule lengths were significantly reduced as the concentration of the extract was increased from 0% to 10%. Maximum reduction in radicle and plumule lengths were recorded with the highest concentration (10%) in this study. Increasing the concentrations of aqueous leaf extracts increased the inhibitory effect of the radicle and plumule lengths of cowpea. Oraon and Mondal (2021) also reported that the germination and seedling growth of several crops are affected by the allelopathic influence of *H. suaveolens*.

CONCLUSION

It can be concluded in this study that irrespective of the type of leaves of *H. suaveolens*, the aqueous extract of the leaves had a strong allelopathic potential to fully inhibit the germination and seedling growth of cowpea. The 10 % concentration of aqueous extract of the *H. suaveolens* leaves strongly inhibited the germination and seedling growth of cowpea.

REFERENCES

- Afreen, T., Srivastava, P., Singh, H. & Singh, J. S. (2017). Effect of invasion by *Hyptis suaveolens* on plant diversity and selected soil properties of a constructed tropical grassland, *Journal of Plant Ecology* 5, 751-760.
- Association of Official Seed Analyst (AOSA). (1990). Rules for testing seeds. *Journal of Seed Technology*, 12, 1 – 112.
- Daniya, E., Ibrahim, H. & Abdullahi, F. (2014). Allelopathic potential of mint weed (*Hyptis suaveolens* Poit.) on germination behaviour of some okra (*Abelmoschus esculentus* (L.) Moench) varieties; *Journal of Applied Agricultural Research* 6(11), 245-251.
- Gerrano, A.S., Van Rensburg, W.S.J & Adebola, P. (2017). Nutritional composition of immature pods in selected cowpea (*Vigna unguiculata* (L.) Walp) genotypes in South Africa. *Australian Journal of Crop Science* 11 (02), 134-141.
- Gupta, K. C., Gupta, A. K. & Saxena, R. (2016). Weed management in cowpea (*Vigna unguiculata* (L.) Walp.) under rainfed conditions. *International Journal of Agricultural Science* 12 (2), 238-240.
- Mominul-Islam, A. K. M. & Kato-Noguchi, H. (2013). “Plant growth inhibitory activity of medicinal plant *Hyptis suaveolens*: could allelopathy be a cause?” *Emirates Journal of Food and Agriculture*, 25(9), 692–701.
- Oraon, S. & Mondal, S. (2021). Allelopathic effect of lamiaceae weeds on seed germination and early growth of aromatic rice (*Oryza sativa*). *Acta Agrobotanica* 74, article 741.
- Osipitan, O. A., Adigun, J. A & Kolawole, R. O. (2016). Row spacing determines critical period of weed control in crop: Cowpea (*Vigna unguiculata*) as a case study. *Azarian Journal of Agriculture* 3 (5), 90-96.
- Padalia, H., Kudrat, M & Sharma K. P. (2013). Mapping sub-pixel occurrence of an alien invasive *Hyptis suaveolens* (L.) Poit. Using spectral unmixing technique. *International Journal of Remote Sensing* 34, 325 – 40.
- Safder, V. & Nix, H. A. (2021). Land use and productive potentials enhancing sustainable production. *International Journal of Research in Agriculture and Forestry*, 16, 17 – 21.
- Weih, M., Didon, U. M. E., Ronnberg-Wastljung, A. C & Bjorkman C. (2008). Integrated agricultural research and crop breeding: allelopathic weed control in cereals and long-term productivity in perennial biomass crops. *Agricultural System* 97, 99–107.
- Yongqing, Ma., Meng, Zhang., Yaolin, Li., Junfeng, Shui & Yeng, Jun Zhou. (2014). Allelopathy of rice (*Oryza sativa*. L.) exudates and its relations with *Orobanche cumana* wallr. and *Orobanche minor* sm. Germination, *Journal of Plant Interactions* 9(1), 722-730.

Table 1: Allelopathic effect of water extract of green and brown leaves of *Hyptis suaveolens* at varying concentration on germination indices of cowpea

Treatment	Germination Percentage	Germination Index	Mean Germination Time
-----------	------------------------	-------------------	-----------------------

Leaf type (L)			
Green	54.0a	2.4a	5.4a
Brown	63.0a	2.7a	6.3a
LSD (0.05)	11.6	0.5	1.2
Concentration(C)			
(%)			
0	81.3a	3.6a	8.1a
2.5	70.0ab	3.1ab	7.0ab
5.0	58.8bc	2.6bc	5.9bc
7.5	51.3c	2.2cd	5.1c
10	31.3d	1.3d	3.1d
LSD (0.05)	18.3	0.8	1.8
Interaction			
L x C	NS	NS	NS

Means with different letters from each other differ significantly at $p \leq 0.05$: NS = Not significant.

Table 2: Allelopathic effect of water extract of green and brown leaves of *Hyptis suaveolens* at varying concentration on radicle length, plumule length, stimulation and inhibition of cowpea

Treatment	Radicle length (cm)	Plumule length (cm)	Stimulation and inhibition (%)
Leaf (L)			
Green	2.8a	6.7a	-28.8a
Brown	3.1a	7.6a	-25.6a
LSD (0.05)	0.5	1.0	16.0
Concentration (C)			
0	3.5a	9.6a	0.0a
2.5	3.5a	8.0b	-12.7ab
5.0	3.0a	7.3b	-26.6b
7.5	3.0a	7.0b	-35.2b
10	1.6b	3.6c	-61.5c
LSD (0.05)	0.8	1.5	25.27
Interaction			
L x C	NS	NS	NS

Means with different letters from each other differ significantly at $p \leq 0.05$: NS = Not significant: negative values represent inhibition