

## EXTRACTION AND PARTIAL PURIFICATION OF PHYTOCYSTATIN FROM *Calotropis procera*

\*<sup>1,2</sup>AbdulKadir, A., <sup>1</sup>Umar, I. A. S., <sup>1</sup>Ibrahim., E., <sup>1</sup>Onyike., <sup>2</sup>Kabiru, A.Y., <sup>1</sup>Pala, Y. Y., and <sup>3</sup>Gabi, B.

<sup>1</sup> Department of Biochemistry, Ahmadu Bello University, Zaria.

<sup>2</sup> Department of Biochemistry, Federal University of Technology, Minna, Niger state, Nigeria.

<sup>3</sup> Department of Applied Sciences, Kaduna Polytechnic, Kaduna, Nigeria.

### ABSTRACT

Phytocystatins are cysteine protease inhibitors (CPIs) that are known for their several applications in biotechnology and medicine. Various extraction media which include, Sodium Chloride, Sodium Hydroxide, Hydrochloric acid, Sodium phosphate buffer and distilled Water were used to evaluate Leave, root, flower, latex and stem bark of *Calotropis procera* for cysteine protease inhibitory activity against Papain enzyme. Sodium phosphate buffer extract from latex of *C. procera* with higher CPI activity were concentrated by cold acetone precipitation and further subjected to purification using ammonium sulphate fractionation and Phenyl sepharose column. The fifty percent inhibitory concentration (IC<sub>50</sub>) for crude latex extract was 1.35mg/ml. Electrophoretic analysis of partially purified protein showed a band with an estimated molecular weight of 10.23 kDa. Cysteine protease inhibitor investigated in this work could serve as template in biotechnology of traditional medicine and transgenic crops to arrest the negative pathogenic expressions of cysteine protease.

**Keywords:** Phytocystatins, cysteine, protease, inhibitory activity

### INTRODUCTION

Phytocystatins are plant proteins that specifically inhibit cysteine proteinase and belong to papain family. Unlike cystatins of animal origin (except stefins), plant derived cystatins have no di-sulfide bond and putative glycosylation site. They are plant proteins cysteine protease inhibitors that constitute a powerful regulatory system for endogenous proteases which may otherwise cause excessive proteolysis and cell damages (Karthik *et al.*, 2014). Cysteine proteases are expressed throughout the animal and plant kingdoms as well as in microorganism. These enzymes have been implicated in

many adaptation mechanisms for survival in the host, including modulation of the host immune system, invasion and destruction of host tissues, parasite dissemination, and acquisition of essential nutrients that assure survival and proliferation to sustain the infection (Pirta *et al.*, 2016).

These enzymes have drawn special attention as drug target for cure of several diseases such as osteoporosis, arthritis, cancer and many parasitic infections (Rosenthal *et al.*, 1999). The interaction between cysteine proteases and inhibitor is a necessary in averting unwanted potentially destructive

proteolysis. These plant derived inhibitors are gaining consideration as potential candidate which can be exploited in the chemotherapy and in the engineering of transgenic crops.

*Calotropis procera* is a plant of *Asclepiadaceae* family and are widely distributed in West Africa. The plant has received special attention because of its use in folk medicine (Lawal *et al.*, 2015). Previous reports have confirmed that this plant possesses various pharmacological properties that may be used for the treatment of inflammatory disorders such as arthritis, cancer and sepsis (Kumar *et al.*, 2014). Furthermore, numerous bioactive proteins were also isolated from this plant and their therapeutic effects documented (Al-snafi, 2015).

#### MATERIALS AND METHODS

The plant parts was collected between periods of November to January in Samaru, Zaria (Latitude 11.15°N and Longitude 7.65°E). The plants were identified in the Department of Biological Sciences, Ahmadu Bello University, Zaria with voucher number 900219.

##### Extraction of Phytocystain

Fresh parts (25g each of); leaves, flowers, latex, root and stem barks of *C.procera* were blended with 100 mL each of sodium chloride 15% (w/v), sodium hydroxide 0.2% (w/v), hydrochloric acid 0.05M, phosphate buffer 0.1 M (pH 7) and distilled water as described by (Wu and Whitaker, 1990). The clear supernatant obtained after centrifugation (10,000 rpm, 15 min, 4°C) represent the crude extract, and was assayed for CPI activity and protein content. The phosphate buffer extract with highest activity was used for further studies.

##### Acetone Precipitation

Four volumes of cold acetone was added with stirring to the crude extract. After 20 min stirring, the precipitate was collected by suction filtration, washed with cold acetone. The precipitate was air dried and stored at 4°C under desiccation. This was suspended in 100mM phosphate buffer, pH 7.6 and was designated as the crude protein extract (Rao *et al.*, 1983).

##### Inhibitory Effect of Acetone Precipitate on Papain Activity

Different concentrations of protein precipitate (ranging from 0.1-3.0mg/ml) were incubated with 1mg/ml papain. Activity of inhibitor was assayed as described below.

##### Protease Inhibitor Assay

Protease inhibitor activity was assayed according to the method of Wannapa *et al.*, (2006). Assays were performed by adding 290µL of 50mM Tris-HCl (pH 7.6) and 200µL of 1.25mM BAPNA solution to the previously preincubated cysteine protease (1mg/ml papain prepared in 100mM phosphate buffer, pH 6.8, containing 0.3mM EDTA and 2 mM cysteine-HCl) with inhibitor extract for 10min, at 25°C. After 30 min at 37°C, the reaction was stopped by adding 150µL of acetic acid (30%). The resulting color was measured spectrophotometrically at 410nm. Protein Estimation

Protein content was determined according to the method of Bradford (1976) using Bovine Serum Albumin (BSA) as the standard.

##### Purification of Crude Extracts from *C.procera*

Crude protein extract was precipitated at different percentages of ammonium sulphate saturation (10-80%).The



precipitated proteins were separated by centrifugation at 10,000 rpm at 4°C for 20 min. Active protein pellets was dialyzed against the buffer. The dialyzed sample was used for Hydrophobic interaction chromatography on a Phenyl Sepharose column (Gupta *et al.*, 2005). The column (1 cm×6.5 cm) was equilibrated with 100mM sodium phosphate buffer, pH7.2 containing 1M ammonium sulphate. The bound proteins were then eluted from the column using 0.1M sodium phosphate buffer, pH 7.2 containing a decreasing step gradient of ammonium sulphate (0.5 M, 0.2 M, 0.1 M). The final fraction was eluted with 0.1M sodium phosphate buffer, pH 7.2 containing no ammonium sulphate.

#### Electrophoresis on SDS-PAGE

This was carried out as described by Laemmli (1970) to determine the Molecular weight.

### RESULTS

The Latex of *plant* showed higher protein contents compared to other parts of the plants (Figure1). The results for inhibitory activity of the different parts of the plant extracts with papain enzyme

(Figures 2) revealed that 0.1M phosphate buffer soluble protein extracts from latex possessed higher inhibitory activity. The Phosphate buffer extracts had higher specific activity (Figure 3) when compared to other parts of the plant and solvents used.

Figure 4 showed dose dependent inhibitory activity of different concentrations of acetone precipitated protein for latex of the plant. The fifty percent inhibitory concentration (IC<sub>50</sub>) for crude latex extract was 1.35mg/ml. Crude acetone precipitated proteins showed maximal specific protease inhibition between 20-60% saturation of ammonium sulphate with maximum activity recorded at 50% saturation (Figure 5).

The activity yield and protein purification fold of cysteine protease inhibitor extracted from the *C.procera* latex is summarized in Tables 2. The final purification fold of 3.95 and activity yield of, 48.72 percent respectively was observed for CPI. SDS-PAGE analysis of partially purified protein peak showed prominent band with an estimated molecular weight of 10.23 kDa (Figure).




Figure 1: Protein Concentration in Extracts of *C. procera* Parts using Different Solvents.




Figure 2: Protease Inhibitor Activity in Extracts of *C. procera* Part using Different Solvents. Values are Mean  $\pm$  SD. (n=3)




Figure 3: Specific Protease Inhibitory Activity in Extracts of *C. procera* using Different Solvents. Values are Mean  $\pm$  SD (n=3)





Figure 4: Protease Inhibitory Activity of the Acetone Precipitate Extracts of *C. procera* Latex. Values are Mean  $\pm$  SD (n=3).





Ammonium sulphate Fractionation Profile of Protease From Extracts of *C. procer* Latex




Figure 6: An Elution Profile for the Hydrophobic Interaction Chromatography of *C. procer* Latex Inhibitor on phenyl-Sepharose Column.

|  | Specific Protease inhibitor activity (Units)/mg protein | Purification Fold | Yield (%) |
|--|---|-------------------|-----------|
| Crude extract  | 19.29   | 1                 | 100       |
| Acetone precipitation  | 39.66   | 2.05              | 73.40     |
| (NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub> fraction (20-60) | 35.84   | 1.85              | 58.29     |
| Hydrophobic interaction chromatography                           | 76.33   | 3.95              | 48.72     |




Figure 7: Electrophoregram of Partially Purified Inhibitor

### DISCUSSION

The central role of cysteine proteases in the pathogenesis and survival parasites in the host erythrocytes makes them attractive drug targets (Rosenthal, 2011). Inhibition of these proteases by natural proteins/peptides that will not have off target binding could be useful in the

treatment of these parasitic diseases (Karthik *et al.*, 2014; Rosenthal, 2011).

In the present investigation, significantly higher specific cysteine protease inhibition was exhibited by phosphate buffer extract compared to other solvents, thus, phosphate buffer was selected as the effective extraction medium for maximal extraction of protease inhibitor without any loss in activity. Saline extract, which contain proteins from the latex of *C.procera*, was reported to have anticancer activities (Oliveira *et al.*, 2010). Other researchers used 0.01MNaOH (Marconi *et al.*, 1993), distilled water (Jung *et al.*, 2012) and 0.1 M phosphate buffer, pH 7.5 (Bijina *et al.*, 2011) as extraction media for various therapeutic proteins.

The crude acetone protein precipitate exhibited dose dependent inhibitory activity with papain enzyme with an IC<sub>50</sub> value of 1350 µg/ml *C.procera*. These findings are more than hundred fold those reported by Abd El-latif (2015) with an IC<sub>50</sub> value of 21.04 µg/ml on papain by inhibitor extracted from *Callosobruchus maculatus*.

Phenyl-Sepharose as an adsorbent for hydrophobic interaction chromatography has been reported to be effective for the purification of some proteases and with purification factors in the range of 2.9 - 60 and activity yields in excess of 88% were obtained (Gupta *et al.*, 2005). The increase in activity of up to 48.7% in this investigation is an indicative of exclusions of endogenous protease and an improved stability of the inhibitors. Karlsson *et al.* (1985) successfully separated calpain and their inhibitors in one single step on phenyl-sepharose column.

SDS-PAGE analysis of the inhibitors under reducing conditions gave an estimated molecular weight of 10.23kDa for *C.procera* CPI. Other researcher had

earlier reported small peptide and proteins of molecular weight between 5-15kDa, with cysteine protease inhibitory activity (Soares *et al.* 2015).

## CONCLUSION

Cysteine protease inhibitors were isolated and purified from *C.procera* latex, with in *vitro* inhibitory activity against papain enzyme. This inhibitor could serve as potential candidate in biotechnology of transgenic crops and chemotherapy.

## ACKNOWLEDGEMENTS

We thank University Board of Research (UBR) of Federal University of Technology, Minna for funding part of this work.

## REFERENCES

- Al-snafi, A. E. (2015). The constituents and Pharmacological properties of *Calotropis procera* an overview. *International Journal of Pharmacy Review and Research*, 5 (3), 259-275.
- Bijina, B., Sreeja, C., Krishna, S. M., Basheer, K. K., Elyas, A. H. and Bahkali, C. M. (2011). Protease Inhibitor from *Moringa oleifera* Leaves: Isolation, Purification and Characterization, *Process Biochemistry*, 46, 2291-2300.
- Bradford, M. M. (1976). A Rapid and Sensitive Method for the Quantitation of Micrograms Quantities for Proteins Utilizing the Principle of Protein Dye Binding. *Analytical Biochemistry*, 72,248-254.



- Gupta, A., Roy, I., Patel, R. K., Singh, S. P., Khare, S. K., and Gupta, M. N. (2005). One-step Purification and Characterization of an Alkaline Protease from Haloalkaliphilic *Bacillus* sp., *Journal of Chromatography, A*, 1075,103-108.
- Jung, S. A., Kim, K., Kim, M., Kim, D., Sun Woo, C., Kim, H., Jeong, D., Jeong, H., Kim, T., Cho, Y. and Ahn, D. (2012). Trypsin Inhibitory Activity of Water Extracts from *Ecklonia cava* as Affected by Temperature and pH. *Journal of the Korean Society of Food Science and Nutrition*, 41 (6), 840-845.
- Karlsson, J., Gustavsson, S., Hall, C. and Nilsson, E. (1985). A Simple One-Step Procedure for the Separation of Calpain I, Calpain II and Calpastatin, *Biochemical Journal*, 231, 201-204.
- Karthik, L., Kumar, G., Keswani, T., Bhattacharyya, A. and Chandar, S.S. (2014). Protease Inhibitors from *Marine Actinobacteria* as a Potential Source for Antimalarial Compound. *Plos One*, 9(3), 1-13.
- Krishnan, V. G. M. and Murugan, K. (2015). Purification, characterization and kinetics of protease inhibitor from fruits of *Solanum aculeatissimum* Jacq. *Food Science and Human Wellness*, 4(3), 97-107.
- Kumar, V. L., Priyanka, C., Renato M. O. and Marcio, V. R. (2014). *Calotropis procera* Latex Proteins Ameliorate Functional Limitations Associated with Adjuvant Induced Inflammation in Rat. *Musculoskeletal Biology*, 10, 1-10.
- Lawal, B., Shittu, O. K., Kabiru, A. Y., Jigam, A. A., Umar, M. B., Berinyuy, E. B. and Alozieuwa, B. U. (2015). Potential Antimalarial from African Natural Products: A review. *Journal of Intercultural Ethno pharmacology*, 4(4), 318-343.
- Marconi, E., Nq NQ, and Carnovale, E. (1993). Protease Inhibitors and Lectins in Cowpea. *Food Chemistry*, 47, 37-40.
- Oliveira, C., Luz, L., Paiva, P., Coelho, L., Marangoni, S. and Macedo, M. (2010). Evaluation of Seed Coagulant *Moringa oleifera* lectin (cMoL) as a Bio insecticidal Tool with Potential for the Control of Insects. *Process Biochemistry*, 46,498-504.
- Pirta, C., Sharma, N. N. and Banyal, H. S. (2016). A 43 kDa Recombinant Plasmepsin Elicits Immune Response in Mice against *Plasmodium berghei* Malaria. *Acta Parasitologica*, 61(1), 102-107.
- Rao, N., Mallikarjuna, H., Nayana, R. and Pattabiraman, T. N. (1983). Enzyme Inhibitors from Plants. Isolation and Characterization of a Protease Inhibitor from Arrow Root (*Maranta arundinaceae*) Tuber. *Journal of Bioscience*, 5 (1), 21-33.

Biotechnology Society of Nigeria 2017 Book of Proceedings

Rosenthal, P. J., Kim, K., McKerrow, J. H. and Leech, J. H. (1999). Identification of Three stage-specific proteinases from *Plasmodium falciparum*. *Journal of Experimental. Medicine*, 166, 816-821.

Shamsi, T. N., Parveen, R. and Fatima, S. (2016). Characterization, Biomedical and Agricultural Applications of Protease Inhibitors: A review. *International Journal of Biological Macromolecules*, 91, 1120-1133.

Wannapa, S., Gregory, P., William, F., Siems, C. A., Ryan, R. W. and Dhirayos, W. (2006). Isolation and Characterization of Is inhibitors of the Potato Protease Inhibitor I Family from the Latex of the Rubber Trees, *Hevea brasiliensis*, *Photochemistry*, 67, 1644-1650.

Wu, C. and Whitaker, J. R. (1990). Purification and Partial Characterization of four Trypsin/Chymotrypsin Inhibitors from Red Kidney Beans (*Phaseolus vulgaris* var. *Linden*), *Journal Agricultural Food Chemistry*, 38, 1523-1529.