




FUDMA Journal OF AGRICULTURE And AGRICULTURAL Technology



About the Journal

FUDMA Journal of Agriculture and Agricultural Technology

Current Issue

Vol. 8 No. 1 (2022): FUDMA Journal of Agriculture and Agricultural Technology

Published: 2022-09-15

Articles

EFFECT OF FERMENTED CINNAMON (*Cinnomomum zeylanicum*) JUICE EXTRACT ON RED SOKOTO BUCKS THERMOREGULATORY INDICES, HAEMATOLOGY AND SERUM BIOCHEMISTRY

M. G GARBA, S GADDAFI, A. H SAIDU

1-7



PROFITABILITY ANALYSIS OF DRY SEASON FLUTED PUMPKIN PRODUCTION AMONG SMALLHOLDER FARMERS IN OKIGWE, SOUTHEASTERN NIGERIA

Oliver Utobo, C I EZEANO, E.E UMEBALI, C.C OKEKE, M.O NWIBO

9-17



ANALYSIS OF AGRO-WASTE UTILISATION AS SOURCE OF ORGANIC MANURE BY VEGETABLE FARMERS IN KUMBOTSO LOCAL GOVERNMENT AREA OF KANO STATE, NIGERIA.

MUDASHIRU AKINYEMI, G. E DAIKWO

18-25



COMPUTATIONAL ANALYSIS OF FUNCTIONAL EFFECTS OF SINGLE NUCLEOTIDE POLYMORPHISM ON OVINE INTERFERON TAU

S.I MUSA, K.O YAHAYA, A YAHAYA

26-29



EFFECT OF WEED CONTROL TREATMENTS, SOWING DATE AND SOWING METHOD ON GROWTH AND YIELD OF FINGER MILLET (*Eleusine coracana* (L.) Gaertn) IN SUDAN SAVANNA OF NIGERIA

T.T BELLO, M. A MAHADI, A LADO

30-37



EFFECT OF COMPOSTED POULTRY MANURE AND BIOCHAR ON BIOACCUMULATION OF LEAD/ZINC IN OKRA (*ABELMOSCHUS ESCULENTUS*. L.) IN AMAGU MINING SOILS

I.G OKORO, U.J NWOKEH, C.U ORODEJI

38-48



ASSESSMENT OF SMALL-HOLDER RICE FARMERS' ADOPTION LEVEL OF CLIMATE-SMART AGRICULTURAL PRACTICES IN KWARA STATE, NIGERIA

Ibrahim. A Oba, H. A ABDULLAHI, A. A JAMILU

49-55



ANALYSIS OF PROFITABILITY AND TECHNICAL EFFICIENCY OF BROILER PRODUCTION IN NORTH-CENTRAL, NIGERIA.

H. S. UMAR, E. G. LUKE, E. S ALU, EMMANUEL PETER

56-63



EFFECT OF DIETARY SUPPLEMENTATION OF BAKER'S YEAST (*Saccharomyces cerevisiae*) ON GROWTH PERFORMANCE AND CARCASS CHARACTERISTICS OF WEANER RABBITS (*Oryctolagus cuniculus*)

Hadiza Ibrahim, J.M GAJERE, A ZAHARADEEN, S. U ABDULMALIK

64-69



NUTRITIVE VALUE OF UNTREATED AND MOLASSES-UREA TREATED TYPHA (*Typha domingensis*) SILAGE

Ahmad Rufai Musa, Y GARBA

70-76



EFFECT OF GARLIC (*Allium sativum*) SUPPLEMENTATION ON GROWTH PERFORMANCE AND CARCASS CHARACTERISTICS OF BROILER CHICKENS

Hadiza Ibrahim, Z ALIYU, T. B OLUWAWUNI, G. AWOLOLA, S.U ABDULMALIK

77-82



pdf

THE EFFECT OF DIFFERENT STORAGE MEDIA AND DURATION ON SEED EMERGENCE AND GROWTH OF MORINGA (*Moringa oleifera* LAM) IN DELTA STATE, NIGERIA

K ATUSA

83-86



pdf

PASTORALISTS' KNOWLEDGE ON PREVALENT CATTLE DISEASES AND PARASITES MANAGEMENT PRACTICES IN NORTH-EAST, NIGERIA.

Bashir Mohammed Bawuro, Ndaghu Augustine Anthony , Anonguku Iorfa

87-98



pdf

SOCIO-ECONOMIC ANALYSIS OF THE ADOPTION OF SELECTED IMPROVED SOYBEAN (*Glycine max*) VARIETIES IN TOFA LOCAL GOVERNMENT AREA OF KANO STATE, NIGERIA

I TAFIDA, B NAZIFI, A. S ADAM

99-107



pdf

APPLICATION OF PARTICIPATORY LAND USE PLANNING TOOL IN SOME COMMUNITY DEVELOPMENT AREAS OF JIGAWA STATE, NIGERIA

Garba Umar Garba Umar, M.J. MAGAJI, A. M. SAMNDI, I. B. BUJI

108-122



pdf

PREVALENCE OF GASTROINTESTINAL HELMINTHS OF AFRICAN CATFISH (*Clarias gariepinus* BURCHELL 1822) IN ZOBE RESERVOIR, KATSINA STATE, NIGERIA

MUSTAPHA AHMADU SADAUKI, AKEEM BABATUNDE DAUDA, MOHAMMAD AWWAL YUSUF

123-130



pdf

RESOURCE-USE EFFICIENCY OF RICE PRODUCTION IN KURA LOCAL GOVERNMENT AREA OF KANO STATE, NIGERIA

GAMA Emmanuel Nkwi, UMAR MUKHTAR, DJOMO CHOUMBOU RAOUL FANI
131-141



ANALYSIS OF POST-HARVEST LOSSES AND TECHNICAL EFFICIENCY OF RICE FARMERS IN MIDDLE RIMA VALLEY IRRIGATION SCHEME (MRVIS), SOKOTO STATE, NIGERIA

A SANI, Umar Ibrahim, S IBRAHIM
142-150



ASSESSMENT OF SOIL MICROBIAL DIVERSITY OF DIFFERENT RUBBER PLANTATION AGES

S. O AGHEDO, A.T. ADEKUNLE, R.O OKUNDIA, U ESEIMUDE, J.A. OMOROGBE
151-157



BIODEGRADED MANGO PEEL BASED DIETS ALTERS LIPOGENIC GENES AND mRNA EXPRESSION IN GROWING RABBITS.

J. A AGBANA, O. C AMANA, D. A. AJAGBE, U OKPANACHI
158-163



HAEMATOLOGICAL AND SERUM BIOCHEMICAL PARAMETERS OF BROILER CHICKENS FED DIFFERENT CEREAL GRAIN BY-PRODUCTS

U. J. BAH, A.B. BABURO, I. M. DIKKO, U. S. BAH
164-170



HAEMATOLOGICAL AND BIOCHEMICAL CHARACTERISTICS OF FINISHER BROILER CHICKENS FED ROASTED BAMBARA NUT (*Vigna subterranea* L.)-BASED DIETS

Nuhu Bello Rano, R. M BASHIR
171-176



FACTORS INFLUENCING ADOPTION OF IMPROVED CASSAVA- GARRI PROCESSING TECHNOLOGIES AMONG RURAL WOMEN IN VANDEIKYA LOCAL GOVERNMENT AREA OF BENUE STATE, NIGERIA

S. K. VIHI, E. A. CHOMINI, L. G TOR, B. JESSE, A. A. DALLA, E. A. BASSEY, G.T. OWA
177-188



COMPARATIVE ANALYSIS OF GOAT AND SHEEP MARKETING IN WUDIL LOCAL GOVERNMENT AREA OF KANO STATE, NIGERIA

Abraham Adaighofua Adomi, U. W UMAR

189-197



RESPONSE OF GROWER RABBITS FED DIETS CONTAINING ROASTED KARAYA GUM TREE (*Sterculia setigera*) SEED MEAL

A. M. YUSUF, A. AREMU, A.T. IJAIYA, A.A. MALIK, B.J. LEKENE, A. SULYMAN

198-202



PERFORMANCE ANALYSIS OF SWEET POTATOES PRODUCTION IN OGBARU L.G.A OF ANAMBRA STATE, NIGERIA

ONYINYECHUKWU HOPE ONUBOGU, EBELE. G. OSEGBUE, IFEANYI. E ODOGWU

203-211



PROXIMATE COMPOSITION, MINERAL CONTENTS AND FATTY ACIDS PROFILE OF SELECTED CROP RESIDUES ACROSS SAVANNAH AGRO-ECOLOGICAL ZONES OF NORTHERN NIGERIA

H. K. YUSUF, Y. GARBA, M.A. ADAM

212-222



EFFECTIVENESS OF AFRICAN LOCUST BEAN WASTEWATER AS A RESOURCE FOR PRESERVING AGRICULTURAL PRODUCE (GRAIN AS A CASE STUDY)

M.A. AMUSAT, F. M. BOLARIN, C. U. ONYEMIZE, O.O. POPOOLA, O.T. DADA-JOEL, A. IBRAHIM, Y.S. ADEMILUYI, M.Y. KASALI

223-226



MORPHOLOGY AND MORPHOMETRY OF THE FORAMEN MAGNUM IN SELECTED NIGERIAN LOCAL BREED DOGS

A. DANMAIGORO, M.A. MAHMUD

227-231



BARRIERS TO THE ADOPTION OF CLIMATE SMART AGRICULTURAL PRACTICES IN THE DRYLAND OF NORTHERN NIGERIA

K SALISU

232-243



EFFECT OF YELLOW VEIN MOSAIC VIRUS DISEASE OF OKRA ON GROWTH AND PROXIMATE ANALYSIS OF SOME OKRA CULTIVARS IN DUTSIN-MA

A. A. AHMED, M. K. LADAN

244-250



ANALYSIS OF UTILIZATION OF INFORMATION AND COMMUNICATION TECHNOLOGY (ICT) AMONG CASSAVA FARMERS IN ANIOCHA SOUTH LOCAL GOVERNMENT AREA OF DELTA STATE, NIGERIA

O OMOREGIE, G.O. AZIKEN

251-257



POTENTIAL OF MAIZE (*Zea mays*) YIELD IN THE SAVANNA OF KANO STATE, SEMI-ARID REGION OF NIGERIA

A.U. ALIYU, M.A. YUSUF, L.F. BUBA

258-264



SURVEY OF CHARACTERISTICS AND CHALLENGES OF LOCAL MILK PRODUCERS IN DAURA LOCAL GOVERNMENT AREA OF KATSINA STATE

A ARUWAYO, S. S. ADEOLA, M. MAMMAN

265-272



MANAGEMENT OF EGG BOUND CONDITION IN NON-DESCRIPT NIGERIAN INDIGENOUS CHICKEN-A CASE REPORT

I.A. IDRIS

273-275



ASSESSMENT OF THE EFFECTS OF INTERNATIONAL FUND FOR AGRICULTURAL DEVELOPMENT'S INTERVENTION ON LIVELIHOOD OF INTERNALLY DISPLACED MILLET FARMERS IN BORNO STATE, NIGERIA

Mary Samuel, Hussein Ahmed Abdullahi, Sani Isiyaku

276-282



pdf

EFFECTS OF FEEDING UREA ENSILED MILLET STOVER BASED DIETS WITH CONCENTRATES ON GROWTH PERFORMANCE AND ECONOMY OF SUPPLEMENTATION IN GROWING YANKASA RAMS

A YUSUF, M. ABUBAKAR, B.F. MUHAMMAD

283-291



pdf

BIOACTIVE POTENCY OF MEDICINAL PLANT EXTRACTS AND SYNTHETIC CHEMICAL ON POST-HARVEST ROT PATHOGENS OF WHITE YAM (*Dioscorea rotundata* Poir) TUBER CULTIVARS

V. I. Gwa, A.A. ABDULLAHI

292-297



pdf

INFLUENCE OF WEIGHT AND SEX ON INTESTINAL BACTERIA OF WILD *Clarias gariepinus* (Burchell, 1822) IN RIVER RIMA SOKOTO STATE, NIGERIA.

M.A. YUSUF, A. M SALIM, I.O. OBAROH

298-303



pdf

EFFECT OF INTERVAL OF FEEDING ROUGHAGE AND CONCENTRATE ON THE HAEMATOLOGICAL AND BIOCHEMICAL PARAMETERS OF SOKOTO RED BUCKLING

A ARUWAYO, R.A. ADELEKE, M SANI

304-312



pdf

VARIABILITY IN SOME SELECTED RAINFALL CHARACTERISTICS AND YAM PRODUCTION IN MAKURDI AND GBOKO, BENUE STATE, NIGERIA.

Shaibu James Idakpo, Fanen Terdoo, Bello Yusuf

313-322



pdf

ECONOMICS ANALYSIS OF FEEDING BRACHIARIA DECUMBENS OR DIGITARIA SMUTSII HAY TO BALAMI, UDA AND YANKASA RAMS

Dr. Immanuel I. Madziga

323-328



pdf

EFFECTS OF ANCHOR BORROWERS PROGRAMME (ABP) CREDIT ON THE PRODUCTIVITY OF BENEFICIARY RICE FARMERS IN KEBBI STATE, NIGERIA

J SALISU, C. O. ADEBAYO, A.J. JIRGI, A.O. OJO

329-338



pdf

SUITABILITY EVALUATION OF RAINFED RICE (ORYZA SATIVA) IN FLOOD PLAIN OF SOUTHERN GUINEA SAVANNAH ZONE OF NIGERIA

J Usman, M USMAN, O.J. OGBU, E.D. AZAGAKU

339-350



pdf

BIOLOGY AND MORPHOMETRICS OF THE FALL ARMYWORM, Spodoptera frugiperda J. E. SMITH (LEPIDOPTERA: NOCTUIDAE) IN IBADAN, SOUTHWEST NIGERIA

O.A. OJUMoola, A.A. OMOLOYE

351-363



pdf

LIVELIHOOD STATUS OF RICE CONTRACT FARMERS IN BENUE STATE OF NIGERIA

K.M. YISA, J. H. TSADO, I. MOHAMMED, H.U. MOHAMMAD, M LAWAL, P. N. KOLO, J.J. PELEMO, O.A. ADEYEMI

364-371



pdf

TECHNICAL EFFICIENCY IN COWPEA (Vigna unguiculata) PRODUCTION IN KANKE, PLATEAU STATE, NIGERIA

Godfrey Onuwa

372-382



pdf

CONSTRAINTS TO YAM (DIOSCOREA SPP) PRICE FORMATION IN THE URBAN AND RURAL MARKETS OF THE CENTRAL AGRICULTURAL ZONE OF NASARAWA STATE, NIGERIA

U. K. IROEGBUTE, I MOHAMMED, S.A. JIBRIL

383-392



POST-HARVEST LOSS AND FOOD SECURITY: A CASE STUDY OF MAJOR FOOD CROPS IN KATSINA STATE, NIGERIA

H.I IBRAHIM, H.Y. IBRAHIM, S. S. ADEOLA, E. A. OJOKO

393-403



[View All Issues >](#)

Platform &
workflow by
OJS / PKP



<https://doi.org/10.33003/jaat.2022.0801.084>

RESPONSE OF GROWER RABBITS FED DIETS CONTAINING ROASTED KARAYA GUM TREE (*Sterculia setigera*) SEED MEAL

*¹Yusuf, A. M., ²Aremu, A., ²Ijaiya, A.T., ²Malik, A.A., ¹Lekene, B. J. and ¹Sulyman, A.
¹Federal College of Wildlife Management of Forestry Research Institute of Nigeria, P.M.B. 268,
New Bussa, Niger State, Nigeria
²Federal University of Technology, Minna, Niger State, Nigeria
*Corresponding Author: yusufaliyu6518@gmail.com Tel: +2348059427514

ABSTRACT

This study was carried out to determine the response of weaned rabbits to roasted *Karaya* gum tree seed meal in a twelve weeks experiment. Three diets were formulated with control diet (I) having 0 % *Karaya* gum tree seed meal (KGSM) while diets II and III had 5 % and 10 % KGSM respectively. Thirty-six heterogenous mixed sexes of weaned rabbits with average initial weight of 525 g were allocated to the three diets in completely randomized design with twelve rabbits per treatment. At the end of the twelve weeks, digestibility trial was conducted. Feed intake and weight gains were determined. Feed intake and weight gains were significantly ($P < 0.05$) higher in diet containing 5 % KGSM. Feed conversion ratio was lower ($P < 0.05$) in treatment containing 5 % KGSM than control and 10 % KGSM diets. Nutrient digestibility was significantly ($P < 0.05$) higher in diet containing 0 % roasted KGSM (control) than treatments with 5 % and 10 % KGSM. However, the values for nutrient digestibility of diet containing 0 % roasted KGSM (control) and treatment with 5 % KGSM were numerically within close range indicating 5 % level of inclusion is comparable to 0 % inclusion. Nutrient digestibility decreased with increase (10 %) in roasted KGSM in the diet. This experiment had revealed that up to 5 % level of inclusion of roasted *Karaya* gum tree seed meal can be incorporated in the diet of rabbit without negative effects on growth performance and nutrient digestibility.

Keywords: Roasted; Karaya; seed; Performance; Digestibility.

INTRODUCTION

Price of most conventional feed ingredients is high sequel to stiff competition for them directly by man and industrial usage (Ojebiyi *et al.*, 2008). While extensive livestock production system lack adequate nutrients for satisfactory productivity, intensive rearing systems are faced with escalating price of conventional feed ingredients thus raising the cost of production (Tewe, 2004). Presently, Maize that is a major conventional feed ingredient is facing competition between man, livestock and industries and therefore, is no longer economically viable using it in livestock feed. Escalating cost of conventional ingredients has been the prime stimulant for continue search for alternatives (Onimisi *et al.*, 2008). The search for alternatives that would reduce cost of production is therefore necessary.

Use of unconventional feedstuffs in animal feed has been recommended (Rabiu *et al.*, 2021) to reduce competition and cost of production. Such unconventional ingredients include seeds of tropical trees that are not competed for. However, such unconventional ingredients have to be available, cheap, easy to obtain, nutritionally adequate and safe for inclusion in the diet without detrimental effects on the health of animal or man that would consume the

animal. Utilisation of some of these tropical seeds is subjected to some inherent constrains of particular importance are problems associated with anti-nutrient factors (Wafer *et al.*, 2021). Presence of anti-nutritional factors such as phytate, oxalate, hydrocyanide and nitrate in some seeds hinder animals from benefiting from them nutritionally (El-Mahmood *et al.*, 2008).

The unconventional feed ingredient focused in this study is *Karaya* gum tree seed. *Karaya* gum tree (*Sterculia setigera*) is a wild plant that is wide spread in tropical Africa (Agishi, 2004). It produces seeds in pods which are less utilized by man and animals. Research on the seeds of this plant could help to reveal the potentials of the seeds as feed ingredient for incorporation in livestock feed. Hamidu (2012) had reported *Karaya* gum tree to contain active metabolites such as tannin, flavonoids, saponin, phenolics and glycosides. Different processing methods to detoxify anti-nutrients include roasting or toasting, soaking in water, fermentation, addition of alkaline salt, extruding and blanching (Ayanwale, 2004). The study adopted roasting as a method of processing to detoxify the anti-nutrients of the seed. This research work therefore was carried out to determine the growth performance and nutrient

digestibility of grower rabbits fed diets containing roasted *Karaya* gum tree seed meal.

MATERIALS AND METHODS

This experiment was conducted at the Rabbitry unit of Federal College of Wildlife Management, New Bussa, Niger State, Nigeria, with Latitude N 9°49'36" and Longitude E 4°34'15" (GPS, 2021). Matured seeds of *Karaya* gum tree were collected within New Bussa and its environs. The collected seeds were cleaned to be free of unwanted particles and properly air-dried.

Table 1: Composition of the Experimental Diets

Composition (%)	Diets		
	I (0 % KGSM)	II (5 % KGSM)	III (10 % KGSM)
Maize	42.00	37.00	32.00
Roasted KGSM	0.00	5.00	10.00
Soya bean Cake	14.50	13.50	12.50
Full fat soya	15.00	14.00	14.00
Rice offal	24.00	26.00	27.00
Bone meal	3.50	3.50	3.50
Vitamin / premix	0.30	0.30	0.30
Lysine	0.20	0.20	0.20
Methionine	0.20	0.20	0.20
Salt	0.30	0.30	0.30
Total	100.00	100.00	100.00
Calculated composition:			
% Crude protein	17.19	17.16	17.12
% Crude fiber	12.34	12.40	12.52
M.E. (Kcal/kg)	2607	2608	2605

KGSM = *Karaya* gum tree seed meal, M.E = Metabolisable energy

Thirty-six heterogenous mixed sexes of weaned rabbits with average initial weight of 525 g were used for the experiment. Each treatment had triplicates with four animals per replicate, housed in wooden/wire hutches of 60 cm × 60 cm × 50 cm (length × breadth × height) in a well-ventilated block (7 m × 7 m) pen. The animals were allotted to the three treatment diets in a completely randomized design experiment. Two weeks adjustment period was given for adaptation followed by twelve weeks experimental period. Feed and water were served *ad-lib*. Feed intake and weight gains were determined over the twelve weeks period followed by one week digestibility period. Two animals from each replicate were randomly selected and transferred to metabolism cages for digestibility trial. They were allowed two days adjustment period and faecal samples were collected for five days. The faecal samples

The seeds were coarsely milled to allow heat to get to particles appropriately, and roasted in open frying pan heated by naked fire from dried wood at about 75° C for thirty (30) minutes with constant stirring. Thereafter, they were milled with hammer mill for inclusion in the diets. Three experimental diets were formulated in which diet I served as control with 0 % *Karaya* gum tree seed meal (KGSM) while diets II and III contained 5 % and 10 % roasted seed meal respectively (Table 1).

collected were properly dried to constant weight, weighed and stored in plastic bags for proximate composition. Raw and roasted *Karaya* gum tree seed meal, feed and faecal samples were analysed for nutrient composition according to Association of Official Analytical Chemists (AOAC, 2000) method. All data collected were subjected to analysis using statistical analysis system (SAS) package while means were separated using Duncan's Multiple Range Test as contained in the same package.

RESULTS AND DISCUSSION

The ingredients composition of the experimental diets is presented in Table 1 while nutrient composition of raw and roasted seed meal and the experimental diets is presented in Table 2.

Table 2: Proximate Composition of Diets, Raw and Roasted *Karaya* Gum Tree Seed Meal

Parameters (%)	Raw KGSM	Roasted KGSM	Diets		
			I (0 % KGSM)	II (5 % KGSM)	III (10 % KGSM)
Dry matter	93.17	94.84	93.89	93.30	94.36
Crude fibre	6.76	6.35	11.96	12.12	12.68
Crude protein	18.72	18.53	16.75	16.94	16.66
Ash	2.56	6.58	11.77	11.64	12.83
Ether extract	18.90	16.83	6.56	7.16	7.30
Nitrogen Free Extract	46.23	46.55	46.85	45.44	44.89

KGSM = *Karaya* gum tree seed meal

The results had shown that roasting had effect on the nutrient composition of the seeds. There was slight increase in the dry matter content of the roasted seeds. This could be due to the exposure of the crushed seed particles to heat which would have decreased the moisture content by evaporation thereby increasing the dry matter. Similarly, ash and nitrogen free extract also increased which could also be due to increase in dry matter content. Ether extract, crude fibre and crude protein decreased due to effect of heat. Heat would have liberated some oil from the seed to reduce the ether extract content. The crude fibre and the crude protein of the diets were similar and were within the range recommended for growing rabbit.

The results of the performance of rabbits fed experimental diets are presented in Table 3.

Table 3: Performance of Rabbits Fed Diets Containing Roasted *Karaya* Gum Tree Seed Meal

Parameters	Diets			SEM	LOS
	I (0 % KGSM)	II (5 % KGSM)	III (10 % KGSM)		
Initial body weight (g)	525.00	525.00	525.00	0.124	NS
Final body weight (g)	2065.00 ^b	2130.00 ^a	2030.00 ^b	13.666	*
Total weight gain (g)	1540.00 ^b	1605.00 ^a	1505.00 ^b	13.650	*
Average daily gain (g)	18.33 ^b	19.11 ^a	17.91 ^c	0.163	*
Daily feed intake (g)	62.11 ^b	63.40 ^b	68.00 ^a	0.492	*
Total feed intake (g)	5217.10 ^c	5325.60 ^b	5712.0 ^a	232.826	*
Feed conversion ratio	3.39 ^{ab}	3.32 ^a	3.80 ^b	0.042	*

^{a, b, c} Means on the same row with different superscripts were significantly (P<0.05) different

LOS = Level of significance, NS = Not significant, * = Significant, SEM = Standard error of mean, KGSM = *Karaya* gum tree seed meal

Significantly (P<0.05) higher daily feed intake and higher total feed intake were recorded in diets containing roasted *Karaya* gum tree seed meal. This could be due to the aroma of the seeds which might have attracted the rabbits to take more feed than those on diet without roasted *Karaya* gum tree seed meal. Increased roasted *Karaya* gum tree seed meal in the diet led to increase in aroma and palatability and consequently feed intake. This is in line with the observation of Yusuf *et al.* (2021). However, average daily gain and total weight gain were significantly

(P<0.05) higher in treatment fed diet with 5 % roasted *Karaya* gum tree seed meal. The values were significantly (P< 0.05) different between the treatments. Similarly, feed conversion ratio followed the same trend with weight gains. The daily feed intake and daily weight gain were better than those recorded by Adeyina and Apata (2008), Carew *et al.* (2008) and Onimisi *et al.* (2008) on effects of dietary level of cocoa bean shell on nutrient digestibility, organ weight and serum metabolites of weaner rabbits, nutrient composition and feed value of *Tephrosia bruceolata* seed meal for rabbits. A preliminary study, growth performance, and

carcass characteristics of young rabbits fed steamed castor bean cake based diets. Significantly ($P<0.05$) higher daily feed intake, higher weight gain and better feed conversion ratio is an indication that 5 % roasted

Karaya gum tree seed meal had no detrimental effect on these parameters. The results of the nutrient digestibility of rabbits fed experimental diets are presented in Table 4.

Table 4: Nutrient Digestibility of Rabbits Fed Diets Containing Roasted *Karaya* Gum Tree Seed Meal

Parameters (%)	Diets			SEM	LOS
	I (0 % KGSM)	II (5 % KGSM)	III (10 % KGSM)		
Dry matter	79.50 ^b	79.74 ^a	69.18 ^c	0.48	*
Crude fibre	79.93 ^a	79.74 ^b	69.19 ^c	0.02	*
Crude protein	79.93 ^a	79.76 ^b	69.18 ^c	0.19	*
Ash	79.93 ^a	79.95 ^a	69.17 ^b	0.46	*
Ether extract	79.94 ^a	79.75 ^b	69.18 ^c	0.08	*
Nitrogen free extract	79.93 ^a	79.74 ^b	69.18 ^c	0.31	*

^{a, b, c} Means on the same row with different superscripts were significantly ($P<0.05$) different

LOS = Level of significance, * = Significant, SEM = Standard error of mean, KGSM = *Karaya* gum tree seed meal

Table 4 shows the nutrient digestibility of rabbits fed diets containing roasted *Karaya* gum tree seed meal. From the results, all the nutrients in the feeds were relatively (more than 60 %) digested. Digestibility of the nutrients was higher in diets containing 0 % roasted KGSM with the exception of dry matter. The values for diets I and II were numerically within close range but were significantly ($P< 0.05$) different. Digestibility decreased with increase in the KGSM in the diet. Inclusion of roasted KGSM beyond 5 % (10 %) had led to decrease in nutrient digestibility which could mean negative effect on nutrient digestibility. This could be probably as a result of residual effect of anti-nutritional factors of the seed and possibly from the amino acid imbalance. It could also be due to high lignin and cellulose fraction of the fibre of roasted KGSM. This agreed with Agbo (2003) who stated that the lignin and cellulose fraction of the dietary fibre of cooked cowpea made the total cell wall, hemicellulose and dry matter highly digestible compare to uncooked cowpea.

CONCLUSION AND RECOMMENDATIONS

From the results obtained in this study, it is evident that up to 5 % roasted *Karaya* gum tree seed meal can be incorporated in the diets of weaned rabbits without any deleterious effect on performance and nutrient digestibility. This will reduce dependency and competition on conventional feed ingredients. However, beyond 5 % level of inclusion may have negative effect on growth performance as well as nutrient digestibility of rabbits. Further studies are recommended on effects on haematological and serum biochemical indices as well as on economic advantage

of using the seed. Further studies are also recommended on effects of the seed on other animals.

REFERENCES

- Adeyina, A.O. and Apata, D.F. (2008). Effects of dietary level of cocoa bean shell (CBS) on nutrients digestibility, organ weight and serum metabolites of weaner rabbits. *Proceedings of the 33th Annual Conference of Nigerian Society for Animal Production (NSAP)* held on 16th-20th March, 2008 at Olabisi Onabanjo University, Yewa Campus, Ayetoro, Ogun State, Nigeria. Pp 178-181.
- Agbo, J. Y. (2003). Nitrogen balance studies in rats fed diets of cowpea (*Vigna unguiculata*) and rice (*Oryza sativa*). *Proceedings of the 8th Annual Conference of Animal Science Association of (ASAN)* held on 15th-18th September, 2003 at Federal University of Technology, Minna, Niger State, Nigeria. Pp 178-182.
- Agishi. E.C. (2004). Etulo, Idoma, Iyede, Tiv and Hausa names of plants. AGITAB publications limited. Makurdi, Nigeria. P 188.
- A.O.A.C. (Association of Official Analytical Chemists) (2000). Official methods of analysis. 15th Edition. Washington, D.C.
- Ayanwale, B.A. (2004). Effect of feeding different alkaline treated soya beans on protein and energy utilisation of broilers. *Pertanika Journal of Tropical Agricultural Science* 27(2): 107-112.

- Carew, S.N., Ayoade, J.A., Okwori, A.I., and Sha'ahu, D.T. (2008). The nutrient composition and feed value of *Tephrosia bracteolata* seed meal for rabbits: A preliminary study. *Proceedings of the 33rd Annual Conference of Nigerian Society for Animal Production (NSAP)* held on 16th-20th March, 2008 at Olabisi Onabanjo University, Yewa Campus, Ayetoro, Ogun State, Nigeria. Pp 164-167.
- EI-Mahmood, A. B., Doughari, B. H. & Cljanji, F. J. (2008). Invitro antibacterial activities of crude extract of *Nauclea latifolia* and *Daniella oliveri*. *Science Research Essay* 3(3): 102-105.
- Hamidu, A. A. (2012). Phytochemical constituents of the leaves of *Sterculia setigera*. *Journal of Pharmacology* 2(1): 62 – 64.
- Ojebiyi, O. O., Farinu, G. O., Togun, V. A., Oladunjoye, I. O., Aderinola, O. A., Babarinde, F. O., Asogba, O. S. and Adegbenro, O. S. (2008). Feeding value of processed cassava peel blood meal mixture on growth performance of cross-bred growing male rabbits. *Proceedings of the 13th Annual Conference of Animal Science Association of Nigeria (ASAN)* held on 13th-19th September, 2008 at Ahmadu Bello University, Zaria, Kaduna State, Nigeria. Pp 401-405.
- Oninisi, P.A., Omage, J. J., Hussaini, N. S. and Ademu, L. (2008). Evaluation of steamed Castor-bean meal on performance of young rabbits: effects on digestibility and histopathology. *Proceedings of the 13th Annual Conference of Animal Science Association of Nigeria (ASAN)* held on 13th-19th September, 2008 at Ahmadu Bello University, Zaria, Kaduna State, Nigeria. Pp 413-417.
- Rabiu, M., Muhammad, U. R., Idris, D.Y., Alimi, H. M. & Madu, A. (2021). Growth performance of broiler chickens fed diets containing levels of boiled moringa (*Moringa oleifera*) seed meal. *Proceedings of 45th Annual Conference of Nigerian Society for Animal Production (NSAP)* held at Federal University, Dutsin-Ma, Katsina State, Nigeria. Pp 36-38.
- Tewe, O. O. (2004). Cassava for livestock feed in sub-Saharan Africa. Publishing Management Service Information Division, Food and Agricultural Organisation, Rome, Italy.
- Wafer, R. J., Abdullahi, U. & Sini, T. (2021). Growth performance of grower rabbits fed graded levels of *Jatropha* seed meal as replacement for soyabean meal. *Proceedings of the 46th Annual Conference of Nigerian Society for Animal Production (NSAP)* held on 15th –18th March, 2021 at Federal University, Dutsin-Ma, Katsina State, Nigeria. Pp 16-19.
- Yusuf, A. M., Aremu, A. Ijaiya, A. T. Malik, A. A. and Adelokun, K. M. (2021). Nutritional evaluation of boiled *Karaya* gum tree (*Sterculia setigera*) seed meal on growth performance and nutrient digestibility of rabbits. *Proceedings of the 46th Annual Conference of Nigerian Society for Animal Production (NSAP)* held on 14th-18th March, 2021 at Federal University, Dutsin-Ma, Katsina State, Nigeria. Pp 187-190.