



NIGERIAN SOCIETY FOR ANIMAL PRODUCTION
(NSAP)



Proceedings of



ANNUAL
CONFERENCE
14TH-17TH MARCH, 2010

P. 3

THEME:
Fast-tracking
ANIMAL AGRICULTURE
in a Challenged Economy

Held at

UNIVERSITY OF IBADAN, NIGERIA

Edited by:

O. J. BABAYEMI
O. A. ABU
E. O. EWUOLA

ISSN:1596-5570



NIGERIAN SOCIETY FOR ANIMAL PRODUCTION

NIGERIAN SOCIETY FOR ANIMAL PRODUCTION

Proceedings of

PROCEEDINGS 35TH ANNUAL CONFERENCE

35th

Editors

O. J. Babayemi, O. A. Abu and E. O. Ewuola

Conference

14th-17th March 2010

**DEPARTMENT OF ANIMAL SCIENCE
FACULTY OF AGRICULTURE AND FORESTRY
UNIVERSITY OF IBADAN**

TECHNOLOGICAL AND AGRICULTURAL IN A CHANGING TECHNOLOGY

14th-17th

University of Ibadan, Nigeria

Edited by

O. J. Babayemi, O. A. Abu and E. O. Ewuola

ISBN 978-90521-5250-5

Copyright © 2010
All rights reserved

Printed in Nigeria
by the University of Ibadan Press

Growth Performance And Nutrient Utilization of <i>Clarias gariepinus</i> Fingerlings Fed With Diets In Which Fish Meal Protein Was Replaced With Soybean Meal Protein.	F.M. Ayanwamide, O.O. Okunlola and J.A. Oshinowo	Emmanuel Alayande College of Education,	263
Diurnal variations in erythrocyte and haemoglobin concentration of African Giant Rat (<i>Cricetomy gambianus</i>)	T. T. Lawal	UI	265
Gross anatomy of male reproductive organs of the domesticated grasscutter (<i>Thryonomys swinderianus</i> , Temminick)	Olukole, S.G. Oyeyemi, M. O. and Oke, B. O.	UI	268
Nutritional Constituents of <i>Tilapia zilli</i> produced under different types of solar Tent driers.	Ipinmoroti, M.. O, Iyiola Adams O and Adeyemi Olusegun	UNI OSUN	272
Evaluation of processed mango seed kernel meal on nutrients digestibility of growing rabbits	R.O. Olabanji, O.A. Amao, M. D. Shittu , O.A.Aderinola and G.O. Tona,	LAUTECH	275
Performance and nutrient digestibility of weaned rabbits fed graded levels of <i>Piliostigma thonningii</i> leaf meal-based diets	G. O. Tona, J. A. Akinlade, R.O. Olabanji, S. U. Onyia and A. B. Adekiitan	LAUTECH	278
Performance of weaned rabbits fed graded levels of fossil shell flour-based diets	Anurudu, N. F. and Familade F.O.	UI	282
Antimicrobial efficacy of a new generation water sanitizer (isochlorr) in aquaculture	Orgem, C. M., Adesiyan, F.A. and Olufemi, B.E.	MAKURDI	284
MONOGASTRIC ANIMAL PRODUCTION, NUTRITION AND MANAGEMENT			287
Cost benefit ratio of varying levels of energy and protein diets under single versus double phase feeding.	A. Aremu, T.Z. Adama. E.L. Shiawoya and B.A. Ayanwale	FUT, Minna	288
Growth performance and carcass characteristics of broilers fed three commercially produced diets	A.D. Olabode and M. U. Onyekwere	Federal College of Agriculture, Ishiagu, Ebonyi State.	292
Substitution of crystalline lysine with solar-dried-blood meal: Economic implications in starter - finisher broiler diets	E.K. Ndelekwute, U. H. O. Uzegb, U. O. Inyang, I.R. Igwe, R. J. Nosike, S.E., Ogbe, D.O. Agbara and P. C. Ndukwu	UYO	295
Effect of <i>Tetrapleura tetraptera</i> under different feeding regimes on growth performance and gut microbes of broiler chicken	B.O. Nweze and A.E. Nwankwagu	Ebonyi	299
Vermitechnology as a potential source of earthworm meal for replacement of fishmeal in poultry diets: A review	A.A. Malik, A. Aremu and A.H. Dikko	FUT, Minna	302
Evaluation of nutritive and anti-nutritive components of raw and processed	C. O. Obun,, S M Yahaya, A Kibon, , A. S. Kehinde, ,	Federal College of Wildlife	305

Nitrogen Retention And Haematological Indices Of West African Dwarf Rams Fed Sundried And Fermented Rumen Digesta And Cage Hen Droppings Diets	A.N. Fajemisin, A.A. Fadiyimu and J.A. Alokun	FUTA	604
Growth and performance characteristics of West African dwarf goats fed trichoderma treated <i>Jatropha curcas</i> seed cake.	Okukpe K.M, Belewu M.A, Badmos A.H.A	UNILORIN	608
Nutrient Digestibility By Kano Brown Goats Fed Graded Levels Of Rice Milling Waste As A Replacement For Wheat Offal	Nayawo, A. A. Maigandi, S. A. and Muhammad, I. R.	Sa'adatu Rimi College of Education	612
Total protein and cholesterol concentrations in the reproductive tract of WAD rams during sperm maturation in the southern guinea savanna.	T. Ahemen,* I.I. Bitto and F. O. I. Anugwa	UMUDIKE	616
The performance of West African Dwarf (Wad) goats fed tridax and siam weed in ficus based diet	Okukpe K.M., Adeloye A.A and Olaniran T.O.	UNILORIN	618
Grazing Behaviour Of Cows And Ewes In Sown Pasture	O.S. Iyasere., J.O. Daramola., O.A. Osinowo., B.R. Adio., M.I. Salami., L.O. Obanla., O.A. Ogunsola., O.E. Oke., J. Ochefu and O.M. Onagbesan.	UNAAB	622
Effect Of Combined Levels Of <i>Panicum maximum</i> (Jacq) And <i>Gliricidia sepium</i> (Jacq) Wale On The Antiquality Factors Digestibility In West African Dwarf Does Fed Cassava Offal Basal Diets	A.M. Ogungbesani: O.O. Emolorunda': O.A. Jiatadu, A.Makanjii, A.A. Lamidi and E.Mbomie	OOU	625
In Vitro Degradation And Methane Gas Production Of Selected Browse Plants Consumed By Small Ruminants In Western Nigeria	Isah, O. A. and Gazaly, M. B.	UNAAB	628
LIVESTOCK ECONOMICS AND EXTENSION			632
A Survey of Prices of Ruminant Animals in Four Market Locations in Niger State.	O.O. Egbewande, M. K. Yusuf, H. Ibrahim and A. L. Boku,	IBB, Lapai	633
Factors Affecting Honey Production Under Improved Method of Beekeeping in Kano State, Nigeria.	A. U. Shu'aib, , S.Isyaku, A. A. Nayawo and I. A. Kurawa	Sa'adatu Rimi College of Education, Kano.	636
Effect of Sex on Carcass Characteristics of Intensively Reared Guinea Pig (<i>Cavia porcellus</i>)	S.S.A. Egena, A.H. Dikko, D.N. Tsado, A.A. Malik and H. Ibrahim	FUT, Minna	640
Determinants of access to credit among poultry farmers in Nigeria: Evidence from Ogun State	D. A. Babalola and G. O. Tayo	Babcock	643
Effects of incubation positioning and	E.J Okpala,, J.A. Edache,	NVRI, Vom	647

VERMITECHNOLOGY AS A POTENTIAL SOURCE OF EARTHWORM MEAL FOR REPLACEMENT OF FISHMEAL IN POULTRY DIETS: A REVIEW

A.A. Malik¹, A. Aremu¹ and A.H. Dikko¹

¹Department of Animal Production, Federal University of Technology, Minna, Niger State.

Corresponding author's e-mail: delemalik@yahoo.com

ABSTRACT

Earthworms are regarded as ecosystem engineers for their ability to modify soils and plant communities. They are a good source of protein and are therefore used as live fish food and bait. Recently, they are being fed to livestock and fish as high quality protein supplements to replace fishmeal in fish and poultry diets because of the high cost of fishmeal, and for their high protein contents (above 60%). Large scale production of earthworm meal can be achieved through vermiculture and vermicomposting technologies: which are inexpensive and locally-available, using epigeic and endogeic species of earthworms that are prolific breeders and have high biomass production. The dynamic and growing poultry industry in Nigeria require a cheap, non-conventional but high quality animal protein ingredient that is readily available in large commercial quantities throughout the year to replace fishmeal. And earthworm meal produced through vermiculture has the greatest potential of meeting this need, thus saving the nation millions of naira expended yearly on the importation of fishmeal from all parts of the world.

INTRODUCTION

Earthworms are soil invertebrates that belong to the phylum *Annelida* and are found throughout the world. They are commonly referred to as ecosystem engineers for their ability to modify soils and plant communities (Hale *et al.*, 2005). They improve the physical structure of the soil, improve soil fertility and improve plant growth and health. Also, large population of earthworm suppresses weed growth, clean up dangerous chemicals in the environment (such as hexachlorocyclohexane) and improves water absorption capacity of the soil, thus preventing soil erosion (Barley, 1961). In aquaculture, earthworm has been found to be a good source of protein and is therefore used as live fish food and bait. Ita *et al.* (1984) reported that the most preferred bait by anglers for sport fishing in the Kainji Lake was earthworms while Tomlin (1983) observed that in North America, there was a strong demand from about 50 million anglers for baitworms. Recently, because of the increased price of fishmeal (which is the sole animal protein source in poultry diets), earthworm is being fed to livestock and fish as a high quality protein supplement (Ravindran *et al.*, 1993). To ensure regular supply of this non-conventional animal protein source throughout the year, large scale cultivation and culturing of earthworms through vermiculture and vermicomposting technologies are being developed, for partial or total replacement of fishmeal in poultry diets.

Why Earthworm Meal?

Earthworm meal prepared from different earthworm species have been reported to contain high protein.

Sogbesan *et al.* (2007) reported a crude protein value of 63.04% for *Hyperiodrilus euryaulos* as compared to 71.64% for Clupeid fishmeal (Table 1). Guerrero (1983) reported 69.8% CP for *Perionyx excavatus* while Medina *et al.* (2003) reported 61.8% for *Eisenia foetida*. Also, the essential amino acids profile of earthworm meal compares favourably with that of fishmeal; with earthworm meal having a higher percentage of methionine than fishmeal (Table 2).

Vermiculture and vermicomposting technologies

Vermicomposting is the biological non-thermophilic decomposition or degradation of organic matter when earthworms feed on organic waste materials. It is used as a popular waste management option in the Americas, Europe and the Indian sub-continent, producing organic fertilizers called vermicomposts as well as earthworm biomass (Mainoo, 2007). The vermicomposts contain not only earthworm castings, but also bedding materials and organic wastes at various stages of decomposition. Hence, vermicomposts often contain 5 to 11 times more nitrogen, phosphorus and potassium than the surrounding soils and is superior to the original manure as a fertilizer. This is because secretions in the intestinal tracts of earthworms, along with soil passing through their gut, make nutrients more concentrated and available for plant uptake, including micronutrients (Dickerson, 2009). Many earthworm species used for vermicomposting around the world are epigeic (i.e. they live in the upper litter layer of the soil, typically ingesting litter materials extensively). Examples include the

following: *Eisenia foetida*, *Eisenia andrei*, *Eisenia hortensis*, *Eisenia veneta*, *Perionyx excavatus*, *Eudrilus eugeniae*, *Amyntas cortices* and *Amyntas gracilis*. Of all these species however, only *Eudrilus eugeniae* (Kinberg) and *Perionyx excavatus* (Perrier) are the most commonly used earthworms for vermicomposting in tropical and sub-tropical countries (Giraddi., *et al.*, 2002). For example, in a 20 day trial in Accra, Ghana, Mainoo (2007) determined that *Eudrilus eugeniae* decomposed 99% of pineapple fibres and 87% of pineapple peels supplied indicating that this earthworm species is a good vermicomposter.

Vermiculture, on the other hand, is the household rearing of earthworms, using locally-available and inexpensive technology, to produce them in large commercial quantities. Species of earthworm most suitable for vermiculture are endogeic earthworms (i.e. those found in the first 10 to 20cm layer of the soil). In Nigeria, the most culturable species is the semi-arid zone earthworm, *Hyperiodrilus euryaulos* (Sogbesan *et al.*, 2007). But anecic earthworms are deep burrowing earthworms that come to the surface only to feed. A good example is the *Lumbricus terrestris* (the North American Night Crawler). It is not good for vermiculture.

During vermiculture, the earthworms are fed fermented, sun-dried and grinded poultry droppings and vegetables (*Amaranthus* species) at 10% of their body weights. This is because the growth rate of earthworms have been shown to strongly correlate with the particle size of organic materials ingested; and finer materials are more preferred and readily ingested and assimilated than coarse ones (Curry and Schmidt, 2007).

ECONOMIC POTENTIAL OF EARTHWORM MEAL AS A FISHMEAL REPLACER

Earthworm meal can be used to replace fishmeal completely in fish feed without any adverse effect on the growth performance of the fish species (Ogbe *et al.*, 2004). Earthworm can be cultured in wooden boxes initially lined with banana leaves and old newspapers and covered with substrate of up to 5cm level before the worms are introduced (Sogbesan *et al.*, 2007). Worm bins can also be made of plastic or from recycled containers like old bathtubs, barrels or trunks. And earthworms have high reproductive rate and biomass production. Sogbesan *et al.* (2007) reported highest biomass production of 25.7g earthworm/week in cellulose substrate, 22.9g earthworm/week in dry neem leaves and soil substrate and lowest value of 17.8g earthworm/week in soil substrate for *Hyperiodrilus euryaulos* (Clausen); while Giraddi *et al.* (2008) reported mean fecundity of 6.75 cocoons/week and 2.63 cocoons/week for *E. eugeniae* and *P. excavatus* respectively. Hence, earthworms can be regarded as prolific breeders.

CONCLUSION

The dynamic and growing poultry industry in Nigeria requires a non-conventional feed ingredient that is cheap, of high protein quality and readily available in large commercial quantities throughout the year to replace fishmeal in poultry diets. Earthworm meal produced through vermiculture and vermicomposting technologies has the greatest potential of meeting this need, thus saving the nation millions of hard-earned foreign exchange expended each year on the importation of fishmeal from all parts of the world.

Table 1: Proximate and Mineral Composition (DM %) of tested Animal Protein Sources

Composition	Earthworm Meal	Clupeid Fishmeal
Crude protein %	63.04	71.46
Ether extract %	5.90	7.97
Crude fibre %	1.90	1.18
Ash %	8.90	18.22
Nitrogen Free Extract %	13.76	3.17
Moisture %	8.6	8.89
Dry matter %	91.40	90.21
Gross energy kJ/100g	1968.24	2074.09
Sodium (g/100g)	0.43	0.91
Calcium (g/100g)	0.53	3.53
Potassium (g/100g)	0.62	0.96
Phosphorus (g/100g)	0.94	2.40
Magnesium (g/100g)	NA	0.08

Source: Sogbesan *et al.* (2007)

Table 2: Essential amino acids composition (g/16Ng dry matter) and amino acids indices of earthworm meal and Clupeid fishmeal

Amino acid composition	Earthworm Meal	Clupeid Fishmeal
Arginine	2.83	5.34
Histidine	1.47	4.19
Isoleucine	2.04	2.62
Leucine	4.11	8.31
Lysine	6.35	10.96
Methionine	5.30	2.26
Phenylalanine	6.26	5.52
Threonine	4.43	5.28
Valine	4.43	5.88
Tryptophan	0.88	0.97
Total essential amino acids	37.99	51.33
Crude protein %	63.04	71.64
EAAI (%)	71.50	96.70
Chemical score/protein score (%)	36.10	48.80
TEAA: Crude protein	0.60	0.72

Source: Sogbesan *et al.* (2007)

REFERENCES

- Barley, K.P. (1961). *Advances in Agronomy*, Vol. 13, pp. 262-264.
- Curry, P.J. and Schmidt O. (2007). The feeding ecology of earthworm – A review. *Pedobiologia* 50:463-477.
- Dickerson, G.W. (2009). Vermicompost. www.earthwormvietnam.com, downloaded in January, 2009.
- Guerrero, R.D. (1983). The culture and use of *Perionyx excavatus* as protein resource in the Philippines. In: J.E. Satchell (Ed.), *Earthworm Ecology*, Chapman and Hall, London: 309-319.
- Giraddi, R.S., Tippannavar, P.S. and Kulkarni, K.A. (2002). Utilization of peregrine earthworm, *Eudrilus eugeniae* (Kinberg), for bioconversion of agriculture, animal and agro-industrial wastes into manure. *Proc. 7th Int.Symp. Earthworm Ecol.*, Cardiff Univ., UK, p.248.
- Medina, A.L., Cova, J.A., Vidna, R.A., Pujic, P., Carlos, M.P. and Toress, V. (2003). Immunology and chemical analysis of proteins from *Eisenia foetida* earthworm. *Food and Agricultural Immunology*, 15 (3-4): 251-263.
- Ogbe, F.G., Tiamiyu, L.O. and Eze, P.N. (2004). Growth performance of *Clarias gariepinus* fingerlings fed earthworm meal (*Lumbricus terrestris*) as replacement for fishmeal. *Conf. Proc. Fisheries Soc. Nig.*, pp. 214-218 (2004).
- Ravindran B., Ravindran, V. and Blair, R. (1993). Feed resources for poultry production in Asia and the Pacific: Animal protein sources. *World's Poultry Sci. J.*, 49: 219-235.
- Sogbesan, O.A., Ugwumba, A.A.A. and Madu, C.T. (2007). Productivity potentials and nutritional values of Semi-Arid Zone earthworm (*Hyperiodrilus euryaulos*; Clausen, 1967) cultured in organic wastes as fish meal supplement. *Pakistan J. Bio. Sci.*, 10 (17): 2992-2997.
- Tomlin, A.D. (1983). The earthworm bait market in North America. In: *Earthworm Ecology*. Ed. Satchell, J., Chapman & Hall, London
- Giraddi, R.S., Gundannavar, K.P., Tippannavar, P.S. and Sunitha, N.D. (2008). Reproductive Potential of Vermicomposting Earthworms, *Eudrilus eugeniae* (Kinberg) and *Perionyx excavatus* (Perrier), as influenced by seasonal factors. *Karnataka J. Agric. Sci.*, 21(1): 38-40.
- Hale, C.M., Frelich, L.E., Reich, P.B. and Pastor, J. (2005). Effects of European earthworm invasion on soil characteristics in Northern hardwood forests of Minnesota, USA. *Ecosystems* 8, 911-927.
- Ita, E.O., Omorinkoba, W.S. and Bankole, N.O. (1984). Preliminary report of Sport Fishing records in KLRI Reservoirs. *Kainji Lake Research Institute Annual Report, 1984*.
- Mainoo, N.K. (2007). Feasibility of low cost vermicompost production in Accra, Ghana. *M.Sc. Research Thesis of McGill University, Montreal, Canada*.