


Mr. M. A. A. A.



**NIGERIAN SOCIETY
FOR ANIMAL PRODUCTION (NSAP)**



**43RD |
ANNUAL
CONFERENCE
OWERRI 2018**



THEME:

**Exploring Science and Technology
INNOVATIONS FOR SUSTAINABLE
Livestock Development**

18th – 22nd March 2018

New SAAT Auditorium
Federal University of Technology Owerri

Book of Proceedings

Editors: I. C. Okoli, I. P. Ogbuwu, O. O Emenalom and B. O. Esonu

APR -29 Inadvertent Slaughter of Female Goats and Foetal Wastages at Rumuokoro Abattoir in Port Harcourt, Rivers State	913
<i>P.K. Ajuogu, A.O. Aniebo, S.N. Wekhe, N.S. Herbert and O.U. Anyaso</i>	
APR 030 Influence of Multi-enzyme Supplementation on Broilers Fed Normal Energy and Protein Level Diet in South-East Tropical Zone of Nigeria	917
<i>R.A. Amaefule, G. Daniel- Igwe, K.U. Amaefule and M.A. Oguike</i>	
APR -31 Carcass Yield of Broiler Chickens Fed Turmeric (<i>Curcuma Longa</i>) Powder and Cayenne Pepper (<i>Capsicum Frutescens</i>) Powder	920
<i>K.A. Sanwo, A.V. Adegoke, I. Abanikanda, L.T. Egbeyale J. A.Abiona, R.A. Sobayo, O.A. Oso</i>	
APR -32 Assessment of Physicochemical, Microbial and Sensory Characteristics of Fresh White and Red Sorghum Extract Preserved West African Soft Cheese Retailed in Ogbomoso, South-Western Nigeria	923
<i>G.O. Tona, G.A. Ibhaze², A.O. Ajala, A.O. Babatunde¹ and T. Adelokun</i>	
APR -33 Economic Characteristics and Sensory Properties of Broiler Chicken Administered Roselle (<i>Hibiscus sabdariffa</i>), Garlic (<i>Allium sativum</i>) and Ginger (<i>Zingiber officinale</i>) Extracts	927
<i>A.A. Malik, E.Z. Jiya, B.A. Ayanwale and D.L. Kasimi</i>	
APR -34 Growth Performance and Carcass Characteristics of Broilers Fed Varying Levels of <i>Ocimum gratissimum</i> Leaf Meal	930
<i>C.O. Osita[*], C.U. Emezue, A.O. Ani, M.C. Ogwuegbu and I.E. Ezemagu</i>	
APR-35 Carcass Measurements and Weights of West African Dwarf Goat as Affected by Dressing Methods	935
<i>I.P. Okoh and E.E. Adomeh</i>	

Animal Health and Disease Control (AHC)

AHC -01 Blood Parameters and Prevalence of Caprine Babesiosis at Azare Abattoir of Bauchi State, Nigeria	939
<i>A. Maidala, A.M. Dunya, I.C. Mohammed O.J. Makinde, I.O. Adejumo and I.J. Dantata</i>	
AHC -02 Pathological Changes in the Liver of Albino Rats Fed Raw or Processed Sickle pod (<i>Senna obtusifolia</i>) Seed Meal Based-Diets	942
<i>C. Augustine</i>	
AHC -03 Poultry Production Methods and Common Diseases Associated With Poultry in Kogi Central	945
<i>M. Abubakar, I. Sani A. Nuratu, O.K. Yusuf and A.M. Umar</i>	
AHC -04 Ameliorative Effects of Kaempferol on Erythrocyte Osmotic Fragility induced by Experimental <i>Trypanosoma brucei</i> Infection in Mice	949
<i>Y. Muhammad, M.M. Suleiman, I.D. Jatau and A.M. Umar</i>	
AHC -05 A Preliminary Study of Prevalence of Laminitis in Nigerian Trade Horses at Obollo-Afor, Enugu State, Nigeria	952
<i>K.C. Ogbanya, C.A. Eze, J.I. Ihedioha, T.O. Nnaji and P.C. Ugwu</i>	
AHC -06 Blood Changes Observed in Laminitic Nigerian Horses	954
<i>K.C. Ogbanya, C.A. Eze, J.I. Ihedioha and P.C. Ugwu</i>	
AHC -07 Comparison of Point-of-Care Glucometers for Blood Glucose Determination in Cows, Goats and Fish	957
<i>C.O. Okorie-Kanu, C.E. Igbokwe, O.J. Okorie-Kanu, P.E. Aba and N.D. Nwagbara</i>	
AHC -08 Methanol Extract of <i>Spondias Mombin</i> Leaf Protects the Liver against Acetaminophen-Induced Hepatotoxicity in Albino Rats	960
<i>P.E. Aba, V.U. Omoja, N.E. Obioma and C.O. Okorie-Kanu</i>	
AHC -09 Participatory Epidemiology and Sero-Surveillance for Avian Influenza in Local Chicken at Live bird Market and House Holds in Enugu State, Nigeria	963
<i>S.C. Okoli, J.A. Nwanta and E.V. Ezenduka</i>	
AHC -10 Knowledge Attitude and Practice of Meat Inspection among Stakeholders in Owerri, South-Eastern Nigeria	967
<i>C. Unamba-Oparah, I.C. Unamba-Oparah, V.U. Odoemelam and B.O. Obasi</i>	
AHC -11 Influence of Body Weight and Methods of Castration on the Growth Performance and Nutrient Digestibility of Savanna Brown Goats	971
<i>B. Abdulkareem, A.I. Osuolale, M.A. Olaoye, D.N. Tsado, A.A. Malik and O.O.A. Fasanya</i>	
AHC -12 Socio-Cultural Characteristics and Drug Use Habit of Poultry Farmers in Akwa Ibom	974

State	
I.O. Basse, O.H. Ogegbuna, C.C. Achonwa, C.C Ugwu and I.C. Okoli	
AHC -13 Hepatoprotective Activity of Methanol Tuber Extract of <i>Cyperus esculentus</i> Linnon CCL₄-Induced Hepatotoxicity in Albino Rats (<i>Rattus norvegicus</i>)	977
T.E. Ihedioha, R.I. Odo, C.E. Chiwetalu and J.I. Ihedioha	
AHC -14 Assessment of Degree of Spoilage of Meat in a Daily Simulated Market Style of Southwest Nigeria	981
O.D. Ilori, O.O. Adekolurejo and T.A.M. Awoniyi	
AHC -15 Current Anticoccidials Used in Nigeria; Benefits and Challenges: Review	984
M.O. Otu, I.A. Lawal, B.D. George, M.S. Abubakar, I.A. Adeyinka, A.A. Sekoni, B.I. Nwagu, F.O. Abeke	
AHC -16 Biosecurity Assessment of Selected Poultry Farms in South-West, Nigeria	988
B.O. Oyebanji, I.T. Akintoke and T.F. Ayodele	
AHC -17 Prospective Use of Anthelmintic Plants for Sustainable Worm Control in Nigeria	992
M.I. Udobi, C.O. Nwosu and A. Onyeabor	
AHC -18 Vaccination: The Future of Helminth Parasite Control	996
M.I. Udobi, A. Onyeabor and E. Uwalaka	
AHC -19 Influence of Body Weight and Methods of Castration on the Growth Performance and Nutrient Digestibility of Savanna Brown Goats	1000
B. Abdulkareem, A.I. Osuolale, M.A. Olaoye, D.N. Tsado, A.A. Malik and O.O.A. Fasanya	
AHC -20 Abundance of Ectoparasites on Ruminant Animals in Langtang South Local Government Area of Plateau State	1004
J.A. Yohana, I.Z. Abu and S.E. Alu	
AHC -21 Assessment of Antimicrobial Properties of Neem (<i>Azadirachta indica</i>) And Moringa (<i>Moringa oleifera</i>) Leaf Meal in Broiler Chicken Production	1008
M.O. Oladele-Bukola, I.O. Olatoye, B.A. Makanjuola, O.O. Olaseinde, O.A. Morakinyo and S.O. Omotoso	
AHC -22 Prevalence of Parasites among Dogs Undergoing Treatment at Polo Veterinary Clinic Jos, North Central Nigeria	1011
D.D. Pam, V.A. Pam, L.R. Felix ¹ , V.A. Adejoh, A. Ombugadu, S. Terhemem	
AHC -23 Prevalence of Human Parasites Associated with Vegetables Sold in Maikatako Market of Bokkos L.G.A Plateau State, Nigeria	1015
V.A. Pam, A.V. Adejoh, A. Ombugadu and D. Kumbak	
AHC -24 Studies on Parasitic Contamination of Soil and Local Drinking Water Source in Doma Local Government Area of Nasarawa State, Nigeria	1019
V.A. Pam, A.A. Idris, A. Ombugadu, A.V. Adejoh, D.D. Pam S. K. Dogo and D. Kumbak	
AHC -25 Fatal Haemorrhagic Enteritis due to Helminthosis in a One-Year- Old Male-Camel-Calf (<i>Camelus dromedarius</i>): A Case Report	1023
M. Abdullahi, Z.M Wunti, M. Hussaini, I. Haruna and A.K Mohammed	
AHC-26 Performance and Histological Responses of Albino Rats Fed <i>Moringa oleifera</i> Seed Meal	1025
E.E. Akangbe and O.A. Abu	
Microlivestock Production (MLP)	
MLP -01 Replacing Soybean with <i>Leucaena leucocephala</i> Leaf in the Diets of Growing Rabbits	1029
O.A. Abu, M. Orunmuyi, O.M. Jesuyon, F.A. Opawoye and G.O. Balogun	
MLP -02 Growth Response and Nutrient Digestibility of Japanese Quails (<i>Coturnix coturnix japonica</i>) Fed Sun-Dried and Honey-Flavored Cassava Peel Meal Diets	1033
H.I. Abdullahi, A.T. Ijaiya, A.A. Malik and S.S.A. Egena	
MLP -03 Effect of Yam-Peel Meal Diets on Performance and Nutrients Digestibility of Grower Rabbits	1036
M.S. Tamburawa, M.A. Suraj, A.G. Khaleel and S. Madaki	
MLP -04 Effect of Dietary Garlic Powder on the Haematology and Cholesterol Level of Japanese Quail	1039
F.U. Udeh, C.C. Onah, C.M. Nwakor, M.O. Onodugo, V.C. Udeh and D. Ozor	

MLP -02

Growth Response and Nutrient Digestibility of Japanese Quails (*Coturnix coturnix japonica*) Fed Sun-Dried and Honey-Flavored Cassava Peel Meal Diets

H.I. Abdullahi, A.T. Ijaiya, A.A. Malik and S.S.A. Egena

Department of Animal Production, Federal University of Technology, Minna, Niger State

Corresponding author: H.I. Abdullahi; E-mail: abdulhusain01@yahoo.com; Phone: 08060498831

Abstract

A six-weeks feeding trial was conducted to determine the effect of feeding varying levels of sun-dried and honey-flavored cassava peel meal, as a replacement of maize, on the growth performance of Japanese quails (*Coturnix coturnix japonica*) using 200 two-weeks-old chicks. The wet cassava peels were collected fresh, cleaned and sun-dried for seven days, after which they were coarsely milled and flavoured with honey. The birds were randomly assigned to four dietary treatments with 45 birds per treatment, each replicated thrice designated T₁, T₂, T₃ and T₄ containing 0, 50, 75 and 100% of sun-dried and honey-flavoured cassava peel meal (HSCPM) as replacement for dietary maize respectively. The parameters measured were initial body weight, final body weight, average feed intake, average weight gained, and feed conversion ratio. At the end of the 5th week, a nutrient digestibility trial was carried out. Results show that there were no significant ($p > 0.05$) differences among the treatment means for initial body weight, average weight gain and feed conversion ratio. Significant ($p < 0.05$) differences were however recorded for average feed intake and final body weight. There were significant ($p < 0.05$) differences in the nutrient digestibility of the birds across the treatment means. Hence, it was concluded that HSCPM can be used to replace up to 75 % maize in the diet of growing Japanese quails with no detrimental effect on growth performance and nutrient digestibility.

Keywords: Honey-flavored, cassava peel meal, growth response, nutrient digestibility, Japanese quails,

Introduction

The animal protein intake level of humans in most developing countries including Nigeria is very low due to the high cost of protein (Abeke *et al.*, 2003). There has been call for substantial increase in the intake of proteins of animal origin in the developing countries (FAO, 1985) as they are richer in the essential amino acids. This low level of animal protein intake has generated concerns as it affects both the physical and mental development of man. In Nigeria, the domestic fowl is the major protein source providing both chicken meat and eggs. In order to widen the scope of meat and egg supply and alleviate the problem of pressure on egg and meat supply from chicken, other under-utilized species are now under focus. Efforts are being directed towards boosting the livestock industry through the introduction of other livestock species which are prolific, have short generation interval and rapid growth (Owen and Amakiri, 2010). Some species having these qualities include Japanese quail, duck, guinea fowl and ostrich (Edache *et al.*, 2007). According to Odunsi *et al.* (2007), Japanese quail are prolific, have short generation interval and rapid growth rate. They are also suitable for diabetic and hypertensive patients because of its low cholesterol level (Agwunobi and Ina-Ibor, 2007). Faniyi (2002) identified feed as constituting up to 70-80 % of total cost of poultry production of which maize constitute the major cost. Many energy supplying feed ingredients have been investigated as possible replacement for maize in the diet of poultry.

One of such alternatives is cassava and its derivatives. A good derivative is cassava peel, a by-product of cassava production, which is found abundantly in Nigeria. It is a cheaper and unconventional alternative feed resource for livestock. Omoikhoje *et al.* (2008) have reported on the extensive use of cassava peel meal as a cheaper substitute for maize in the diet of monogastric animals.

The objective of the study is to determine the growth response and nutrient digestibility of Japanese quails fed sun-dried and honey-flavoured cassava peel meal diets.

Materials and Methods

This study was carried out at the Poultry Unit of the Department of Animal Production Teaching and Research Farm, Federal University of Technology, Minna, Niger State, Nigeria. Minna lies between latitude 9° 28' and 9° 37' North and on longitude 6° 23' and 6° 33' East. The research was carried out in 6 weeks. Fresh cassava peels were obtained from cassava processing plants within Minna metropolis were cleaned up and spread on a clear polythene sheet to sun-dry for 7 days. The peels were then coarsely milled with an attrition machine and flavoured with honey at 6 % inclusion level to form the honey-flavoured cassava peel meal (HSCPM). Honey inclusion at 6% was the amount in percentage of honey that was used to flavour the cassava peel. HSCPM with others ingredients were used to prepare the experimental diets with HSCPM replacing maize at 0, 50, 75 and 100% respectively (Table 1). Two hundred (200) two-weeks-old Japanese quails sourced from National Veterinary Research Institute Vom, Plateau State, Nigeria, were used for the experiment which were randomly allotted into four treatments

with 45 birds per treatment each replicated thrice (15 birds per replicate) in a completely randomized design. The cages were equipped with feeders and drinkers. Lighting and heat were provided using 100 watt bulbs.

Table 1: Composition of experimental diets fed to growing Japanese quails (%)

	T ₁	T ₂	T ₃	T ₄
Ingredients (%)				
Maize	44.40	21.23	10.39	0.00
HSCPM	0.00	21.23	31.16	40.67
Groundnut	38.90	40.84	41.75	42.63
Maize offal	10.00	10.00	10.00	10.00
Fish meal	2.00	2.00	2.00	2.00
Bone meal	2.50	2.50	2.50	2.50
Limestone	1.50	1.50	1.50	1.50
Methionine	0.10	0.10	0.10	0.10
Lysine	0.10	0.10	0.10	0.10
*Premix	0.25	0.25	0.25	0.25
Salt	0.25	0.25	0.25	0.25
Total	100.00	100.00	100.00	100.00
Calculated analysis				
ME(Kcal/Kg)	2664.71	2396.71	2271.35	2151.47
Crude protein	24.00	24.00	24.00	23.15
Crude fibre	3.76	5.23	5.91	6.57
Lysine	0.98	0.95	0.94	0.93
Methionine	0.41	0.37	0.35	0.33
Calcium	1.62	1.62	1.62	1.62
Phosphorus	0.82	0.77	0.74	0.72

*Premix supplied per 0.25kg: vit. A (7,500.00iu), vit. D (500,000iu), vit. E (1,000iu), vit. B₁ (375mg), vit B₂ (125mg), vit. B₃ (500mg), vit. B₆ (150mg), vit. B₁₂ (2.5mg), vit. K (15mg), vit. C (10mg) and folic acid (150mg), Ca (12.5mg), Cu (8.0mg), Fe (32mg), I (0.8mg), Se (100mg), Mg (0.25mg), Chlorine (250mg), panthotenic acid (14.4mg). HSCPM=Sun-dried Cassava Peel Meal; T1=100% maize, 0% HSCPM; T2=50% maize, 50% HSCPM; T3=75% maize, 25% HSCPM; T4=100% HSCPM, 0% maize.

Routine management operations were carried out. Data was collected on average feed intake, average weight gained and feed conversion ratio. At the end of the 5th week of the experiment, five birds were randomly selected from each replicate and moved into metabolism cages for nutrient digestibility studies. They were adjusted in the cages for three days, followed by faecal collection for four days using the total collection method. The experimental diets and the collected faecal samples were analyzed for their proximate composition using the procedures of AOAC (1990) to determine the apparent nutrient digestibility of the experimental diets by the growing quails. Data collected were subjected to analysis of variance using SPSS 17.0 version.

Results and discussion

The results of growth performance and nutrient digestibility are presented in Table 2. The final body weight and the daily feed intake were significantly ($p < 0.05$) affected by the inclusion levels of HSCPM in the diets of the birds. The increased feed intake observed in the birds fed HSCPM might be as a result of the high fibre content and low energy value of the cassava peel meal which is in agreement with the findings of Salami and Odunsi (2003) who reported that increase in crude fibre decreases the metabolizable energy and leads to increase in the feed intake. It can also be attributed to the honey inclusion in their diets which improved the nutritive value, palatability, growth rate and feed efficiency (Busserolles *et al.*, 2002). Quails fed 75% HSCPM had significantly ($p < 0.05$) better digestibility which is in disagreement with earlier findings of Onyimonyi and Ugwu (2007) who reported that birds could tolerate up to 50% cassava peel as replacement for maize beyond which decrease in weight and poor digestibility sets in. The tolerance level might be due to flavouring of the cassava peel meal with honey which tends to improve the nutritive value, palatability and feed efficiency of the diets. It can therefore be concluded that HSCPM can be used to replace up to 75 % maize in the diet of growing Japanese quails with no detrimental effect on growth performance and nutrient digestibility.

Table 2: Growth performance and nutrient digestibility of Japanese quails fed dietary levels of sun-dried and honey-flavoured cassava peel meal

Parameters	T ₁	T ₂	T ₃	T ₄	SEM	LS
Initial body weight (g/b)	4.44	4.54	4.29	4.29	0.05	NS
Final body weight (g/b)	13.65 ^a	13.81 ^a	12.22 ^b	12.54 ^b	0.21	*
Daily weight gain (g/b)	3.07	3.07	2.64	2.75	0.42	NS
Daily feed intake (g/b)	13.86 ^c	18.80 ^a	17.98 ^b	18.08 ^b	0.16	*
Feed conversion ratio	0.61	0.94	1.20	1.32	0.20	NS
Nutrient digestibility coefficient (%)						
Dry matter	76.43 ^a	69.13 ^b	77.45 ^a	70.73 ^b	0.94	*
Crude protein	74.03 ^{ab}	69.78 ^c	77.53 ^a	73.09 ^{bc}	0.84	*
Crude fibre	83.84 ^c	68.46 ^b	83.72 ^a	63.46 ^c	1.88	*
Ether extract	84.14 ^c	85.71 ^{bc}	91.15 ^a	86.67 ^b	0.59	*
Ash	54.17 ^c	64.73 ^b	73.31 ^a	70.66 ^{ab}	1.80	*
Nitrogen free extract	64.37 ^b	65.26 ^b	72.90 ^a	66.26 ^b	0.94	*

^{abc}Means in the same row with different superscripts were significantly ($P<0.05$) different. SEM= standard error of means; LS= Level of significance; *= significantly different ($P<0.05$); g/b= gram per bird, HSCPM= Honey-flavoured sun-dried cassava peel meal; T1=100% maize, 0% HSCPM; T2=50% maize, 50% HSCPM; T3=25% maize, 75% HSCPM; T4=0% maize, 100% HSCPM.

References

- Abeke, F.O., Ogunduge, S.O., Sekoni, A.A., Adeyinka, A., Abubakar, B.Y., Oni, O.O. and Nwagu, B.I. (2003). Response of laying hens to dietary levels of heat-treatment sheep manure. *Tropical Journal for Animal Science*, 2: 111-116.
- Agwunobi, L.N. and Ina-Ibor, O.B. (2007). Processed rubber feed meal in the diets of growing Japanese quails (*Coturnix coturnix japonica*). *Proc. of the 32nd Ann. Conference of the Nigeria Society for Animal Prod., Calabar, Nigeria*. Pp: 380-382.
- AOAC (1990). Official methods of analysis, 15th edition. Association of Official Analytical Chemists, Washington DC.
- Busserolles, T., Gueux, E., Rock, E., Mazur, A. and Rayssiguire, A. (2002). Substituting honey for refined carbohydrates protects rats from hyper triglyceridemic and pro oxidative effect on fructose. *Journal of Nutrition*, 132: 3379-3382.
- Edache, J.A., Musa, U., Karsin, P.D., Esilonu, J.O., Yisa, A., Okpala, E.J and Zwandor, N.J. (2007). The feeding value of cassava meal diets for growing Japanese quails (*Coturnix coturnix japonica*). *Nigerian Journal of Animal Production*, 34(1): 77-82.
- Faniyi, G.F. (2002). Replacement of wheat offal with untreated citrus pulp in broiler chick diet. *Tropical Animal Production Investment*, 5: 95-100.
- FAO (1985). Food and Agricultural Organization Year book. 1986. vol. 139. Food and Agricultural Organization, Rome.
- Odunsi, A.A., Rotimi, A.A., and Amao, E.A. (2007). Effect of different vegetable protein sources on growth and laying performance of Japanese quails (*Coturnix coturnix japonica*) in a derived savannah zone of Nigeria. *World Applied Sciences Journal*, 3(5): 567-571.
- Omoikhoje, S.O., Bamgbose, A. M. and Aruna, M. B. (2008). Replacement value of unpeeled cassava root meal (UCRM) for maize in weaner rabbit diets. *Nigeria Journal Animal Production*, 35, 63-68.
- Owen, O.J. and Amakiri, A.O. (2010). Japanese quail (*coturnix coturnix japonica*); its potentials, opportunities and challenges. *Proceedings of the 31st Annual conference of the Nigerian Society for Animal production*. 14-17th March 2010. University of Ibadan, Nigeria. Pp: 333-335.
- Onyimonyi, A.E. and Ugwu, S.O.C. (2007). Bioeconomic indices of broiler chicks fed varying ratios of cassava peel bovine blood. *International Journal of Poultry Science*, 6(5): 318-321.
- Salami, R.I. and Odunsi, A.A. (2003). Evaluation of processed cassava peel meals as substitutes for maize in the diets of layers. *International Journal of Poultry Science*, 2(2): 112-116.