2nd

International Conference of Agriculture and Agricultural Technology

Theme:

Climate-Smart Agriculture in the Post COVID Era:

A Gate Way to Food Security in Africa



Held at Caverton Hall Federal University of Technology Minna, Nigeria

Published by

School of Agriculture and Agricultural technology
Federal University of Technology
P. O. M. 65, Minna, Nigeria
Email: icaat@futminna.edu.ng

Edited by

r. Muhammad, H. U. Dr. C. M. Yakubu Dr. Rajan Sharma ♣ Prof. Alabi, O. J.
♣ Dr. Otu, B. O.
♣ Dr. (Mrs) Akande, K. E.
♣ Dr. (Mrs) Adediran, O. A.

♣ Dr. Mrs Carolyne Cherotich
♣ Mr. Ibrahim, A.
♣ Mr Adesina, O. A













PROCEEDINGS

of the

2nd International Conference of

School of Agriculture and Agricultural Technology (SAAT)

Held at

Caverton Hall
Federal University of Technology
Minna

"ICAAT 2022"

11th -13th December 2022

School of Agriculture and Agricultural technology Federal University of Technology P. O. M. 65, Minna, Nigeria Email: icaat@futminna.edu.ng

DEAN

Prof. J.N. Nmadu

HEADS OF DEPARTMENTS

Prof. A. J. Jirgi Agricultural Economics and Farm Management Prof. J. H. Tsado Agricultural Extension and Rural Development

Prof. D.N. Tsado Animal Production

Dr. E. Daniya Crop Production Technology
Dr. H. Ibrahim Horticultural Department
Prof. C. E. Chinma Food Science and Technology
Dr. B. A. Lawal Soil Science and Land Management

Prof. G.G. Bake Water Resources, Aquaculture and Fishery Technology

MANUSCRIPT REVIEWERS

Prof. E. Z. Jiya

Prof. M. A. Ojo

Prof. R. O. Ojutiku

Dr. M. A. Ndanitsa

Prof. A. M. Orire

Prof. O. J. Alabi

Dr. E. S. Yisa

Dr. O. J. Ajayi

Dr. (Mrs.) A. O. Ojo

Dr. (Mrs.) K.E. Akande

Dr. C. M. Yakubu

Dr. S. James

Dr. (Mrs.) M. O. Ojo

Dr. (Mrs.) D. T. Ezekiel-Adewoyin

Dr. C. O. Adebayo

Dr. B. A Lawal

Dr. K. D. Tolorunse

Dr. U.H. Muhammad

Dr. (Mrs.) O. A. Adediran

Mr. H. A. Ibrahim

Dr. I. T. Salihu

Dr. B. O. Otu

Mr. O. A. Adesina

Mrs. F. A. Femi

Dr S.A Ayodele

Dr. S. O Medayese

Engr Prof J. Musa

Dr. Rajan Sharma

Dr S. Y. Mohammed

Dr Mrs Carolyne Cherotich

LIST OF EDITORS

Name	Address	
Prof O. J. Alabi	Animal Production Department, Federal University of Technology, Minna, Nigeria	
Dr Mrs O. A. Adediran	Horticultural Department, Federal University of Technology, Minna, Nigeria	
Dr Mrs K. E. Akande	Animal Production Department, Federal University of Technology, Minna, Nigeria	
Dr B. O. Otu	Animal Production Department, Federal University of Technology, Minna, Nigeria	
Dr U. H. Muhammad	Agricultural Extension Department, Federal University of Technology, Minna, Nigeria	
Dr C. M. Yakubu	Department of Food Science and Technology, Federal University of Technology, Minna, Nigeria	
Dr Rajan Sharma	Department of Food Science and Technology, Punjab Agricultural University, Ludhiana, India	
Dr Carolyne Cherotich	Department of Agricultural Biosystems, Economics and Horticulture, School of Agricultural Sciences & Natural Resources, University of Kabianga, Kericho, Kenya.	
Mr A. Ibrahim	Department of Water Resources, Aquaculture and Fisheries Technology, Federal University of Technology, Minna, Nigeria	
Mr O. Adesina	Horticultural Department, Federal University of Technology, Minna, Nigeria	

LOCAL ORGANISING COMMITTEE (LOC)

Prof. E. Z. Jiya	Chairman
Prof. A. T. Yisa	Member
Prof. I. S. Umar	Member
Prof. M. A. Ojo	Member
Prof. O. J. Alabi	Member
Dr. U. H. Muhammad	Member
Dr B. O. Otu	Member
Dr. Mrs A. O. Ojo	Member
Dr. Mrs. S. U. Ibrahim	Member
Mrs F.A. Femi	Member
Dr. Mrs K. E. Akande	Member
Mr O. A. Adesina	Member
Mr A. Ibrahim	Member
Dr Mrs. O. A. Adediran	Secretary

SCIENTIFIC/TECHNICAL SUBCOMMITTEE

Prof. O. J. Alabi Chairman Prof. M. A. Ojo Member Dr. (Mrs) K. E. Akande Member Dr. B. O. Otu Member Dr. U. H. Muhammad Member Mr O. A. Adesina Member Mr A. Ibrahim Member Mrs. F. A. Femi Secretary

ICAAT 2022 FUNDRAISING SUBCOMMITTEE

Prof. I. S. Umar

Prof. E. Z. Jiya

Member

Dr. (Mrs) S. U. Ibrahim

Member

Dr. U.H. Muhammad

Member

Dr. A. T. Yisa

Member

Prof. O. J. Alabi

Member

Dr. (Mrs) A. O. Ojo

Member

ENTERTAINMENT SUBCOMMITTEE

Dr. (Mrs) A. O. Ojo

Chairperson

Dr. (Mrs) S. U. Ibrahim

Dr. (Mrs) O. A. Adediran

Dr. A. O. Uzoma

Dr. A. T. Yisa

Chairperson

Member

Member

Member

PUBLICITY AND PROTOCOL SUBCOMMITTEE

Prof. M. A. Ojo
Prof. E. Z. Jiya
Member
Dr. B. O. Otu
Member
Prof. O. J. Alabi
Mrs. F. A. Femi
Member

LOGISTICS SUBCOMMITTEE

Prof. A. T. Yisa

Chairman

Dr. A. O. Uzoma

Member

Dr. (Mrs) O. A. Adediran

Mrs. F. A. Femi

Member

TABLE OF CONTENTS

PRO	GRAMME OF EVENTS	VII
	TAINABLE CROP PRODUCTION PRACTICES FOR CLIMATE RESILIENCE FOOD AND RITION SECURITY	3
AGR	ICULTURALECONOMICS EXTENSION AND RURAL SOCIOLOGY	9
1	ADOPTION INDEX OF MAIZE PRODUCTION TECHNOLOGIES AND CORRELATION MATR IN SMALLHOLDER SYSTEMS	IX 10
2	RURAL WOMEN AND AGRO – PROCESSING: A CASE STUDY OF RURAL WOMEN PARTICIPATION IN GROUNDNUT PROCESSING IN KATSINA STATE	16
3	PROFITABILITY OF DRIP IRRIGATED MAIZE PRODUCTION FOR IMPROVED FOOD SECURITY IN MAIDUGURI SEMI-ARID REGION OF BORNO STATE, NIGERIA	22
4	PROFITABILITY ANALYSIS OF POULTRY EGG PRODUCTION IN IBADAN METROPOLIS, O'STATE NIGERIA	YO 30
5	ADOPTION OF MODERN BEEKEEPING TECHOLOGIES IN SELECTED LOCAL GOVERNMENT AREAS OF BENUE STATE NIGERIA	NT 37
6	ADOPTION OF RECOMMENDED COCOYAM PRODUCTION TECHNOLOGIES AMONG FARMERS IN ENUGU STATE NIGERIA	45
7	GENDER ANALYSIS OF FARMING HOUSEHOLDS' ACCESS TO LIVELIHOOD RESOURCES I SELECTED LOCAL GOVERNMENT AREAS OF NIGER STATE, NIGERIA	N 55
8	INNOVATIVE APPLICATION IN MANAGEMENT OF PROBLEMATIC SOIL (ACIDIC SOIL) UNDER MAIZE PRODUCTION IN NIGER STATE, NIGERIA	66
9	EFFECTS OF RISK MANAGEMENT STRATEGIES ON POVERTY STATUS OF RICE FARMERS NIGER STATE, NIGERIA	5 IN 73
10	EFFECTS OF FARMER-HERDER'S CONFLICT ON THE FARMERS PRODUCTIVITY IN ADAMAWA STATE, NIGERIA	79
11	EFFECTS OF ADOPTION OF IMPROVED BEEHIVE TECHNOLOGIES ON INCOME AND WELFARE STATUS OF BEEKEEPERS IN EKITI STATE, NIGERIA	86
12	ASSESSMENT OF OUT-MIGRATION AMONG ARABLE CROP FARMERS IN KOGI STATE, NIGERIA: GENDER DYNAMIC APPROACH	93

13	THE EFFECT OF AGRITECH STARTUPS ON PARTICIPATION AND POVERTY STATUS OF ADOPTERS IN OGUN STATE, NIGERIA	102
14	MICROBIOLOGICAL QUALITY OF OVEN ROASTED PLANTAIN (MUSA PARASIDIACA)	114
15	A REVIEW OF FOOD SECURITY AND POVERTY STATUS OF WOMEN FARMERS UNDER IFAD-VCDP IN NIGER STATE, NIGERIA	119
16	COMPARATIVE ANALYSIS OF THE EFFECT OF ANCHOR BORROWERS PROGRAMME (ABP) ON FOOD SECURITY STATUS OF RICE FARMERS IN EBONYI AND KEBBI STATES, NIGERIA	126
17	EFFECTS OF LAND DEGRADATION ON CEREAL CROP PRODUCTION IN RURAL AREAS ON NIGER STATE, NIGERIA	OF 133
18	FUNCTIONAL PROPERTIES OF CASSAVA SEED PROTEIN CON	140
19	EFFECTS OF INSURGENCY ON CROP FARMING ACTIVITIES OF RURAL WOMEN IN ADAMAWA STATE, NIGERIA	145
20	ADOPTION OF BIO-FORTIFIED FOOD CROP IN NIGERIA: A REVIEW	151
21	DETERMINANTS OF THE ADOPTION OF IMPROVED BEEHIVE TECHNOLOGIES IN BENUSTATE, NIGERIA	JE 155
22	FUNGI ASSOCIATED WITH MILLET GROWN IN ZONE A AGRO-GEOGRAPHICAL ZONE ON NIGER STATE NORTH CENTRAL NIGERIA	OF 169
23	GENDER ANALYSIS OF FARMING HOUSEHOLDS' ACCESS TO LIVELIHOOD RESOURCES SELECTED LOCAL GOVERNMENT AREAS OF NIGER STATE, NIGERIA	IN 176
24	FARMERS' PERCEPTION OF EFFECTIVENESS OF EXTENSION AGENTS IN OSUN STATE, NIGERIA	183
25	SOCIOECONOMIC FACTORS INFLUENCING THE KNOWLEDGE AND ATTITUDE OF MAI FARMERS ON THE SAFE USE OF AGROCHEMICAL IN AGAIE AND BIDA, NIGER STATE, NIGERIA	ZE 190
26	A REVIEW ONEFFECTS OF BIOFORTIFIED PRO-VITAMIN A MAIZEADOPTION ON FARMER'SLIVELIHOODSTATUS IN NIGER STATE NIGERIA	197
27	YOUTH LED ENTERPRISES UNDER INPUT SUBSIDIES ON RICE PRODUCTION IN NIGER BENUE STATES, NIGERIA- A REVIEW	AND 205

28	PROGRAMME AND ITS EFFECTS ON RICE FARMER'S PRODUCTIVITY IN NIGER STATE, NIGERIA	211
29	A REVIEW ON THE EFFECTS OF ADOPTION OF CLIMATE-SMARTS AGRICULTURAL (C.S. PRACTICES ON THE PRODUCTIVITY OF RICE FARMERS IN KWARA AND NIGER STATE,	•
30	NIGERIA ANALYSIS OF DETERMINANTS OF MARKET PARTICIPATION AMONG LOCAL RICE	222
31	FARMERS IN SELECTED LOCAL GOVERNMENT AREAS OF NIGER STATE, NIGERIA LIVELIHOOD DIVERSIFICATION STRATEGIES AND ITS EFFECTS ON FOOD SECURITY	227
21	STATUS OF RURAL HOUSEHOLDS IN KADUNA AND KANO STATES, NIGERIA: A REVIEW	/ 237
32	DETERMINANTS OF FINANCIAL INCLUSION AMONG WOMEN UNDER RURAL FINANCINSTITUTION BUILDING PROGRAMME IN NORTHERN NIGERIA	E 242
33	IDENTIFYING STAKEHOLDERS' INTEREST IN SALINE QUINOA FARMING ALONG VALUE CHAIN IN MOROCCO	249
34	EFFECT OF CLIMATE-SMART AGRICULTURAL PRACTICES ON FOOD SECURITY OF RUR FARMING HOUSEHOLD IN SOUTHWEST, NIGERIA	AL 258
35	EFFECTS OF LAND TENURIAL SYSTEM ON THE PRODUCTIVITY OF SMALLHOLDER RICE FARMERS IN NASARAWA STATE, NIGERIA	265
ANI	MAL SCIENCE/PRODUCTION	272
36	EFFECT OF DIFFERENT PROCESSING METHODS ON NUTRIENTS AND ANTI-NUTRIENT COMPOSITION OF LEUCAENA (<i>LEUCAENA LEUCOCEPH</i> ALA) SEEDS	273
37	PERFORMANCE OF ARBOR ACRE BROILER CHICKENS FED DIET ENRICHED WITH VARY LEVELS OF NANO ZINC SUPPLEMENTATION AT FINISHER PHASE	'ING 279
38	DIVERSITY AND COMPOSITION OF UNDERSTORY SPECIES IN CEDRELA ODORATA AN PINUS CARIBAEA PLANTATIONS IN OMO BIOSPHERE RESERVE, AREA J4, OGUN STATI	
39	ASSESSMENT OF MATING PROFILE OF RED SOKOTO BUCKS ADMINISTERED VARYIN DOSAGE OF ETHANOLIC EXTRACT OF TIGER NUT (CYPERUS ESCULENTUS)	G 296
40	APPARENT NUTRIENT DIGESTIBILITY AND PERFORMANCE OF GROWER PIGS FED ENEAGRO BY-PRODUCTS	RGY 301
41	FREE RADICAL SCAVENGING ACTIVITY OF SELIM POD (XYLOPIA AETHIOPICA)	306

42	ASSESSMENT OF DIFFERENT DOSES OF AQUEOUS RED HOT PEPPER EXTRACT ON TH GROWTH PERFORMANCE OF THREE BREEDS OF BROILER CHICKENS	311
43	ANTIOXIDANT ACTIVITY OF SUN-DRIED TROPICAL LEMON (CITRUS LIMON) PEEL	316
44	EVALUATION OF SYZYGIUM <i>AROMATICUM</i> (CLOVE BUD) AS A POTENTIAL SOURCE ON NATURAL ANTIOXIDANT	F 322
45	GROWTH PERFORMANCE AND NUTRIENT DIGESTIBILITY OF RABBITS FED DIETS CONTAINING SHEA BUTTER CAKE FERMENTED WITH <i>ASPERGILLUS NIGER</i>	326
46	MEAT QUALITY CHARACTERISTICS OF WEANER RABBITS FED DIETS CONTAINING SHE BUTTER CAKE FERMENTED WITH ASPERGILLUS NIGER	A 331
47	GROWTH PERFORMANCE AND CORRELATION MATRIX OF SAVANNA BROWN GOATS ENZYME TREATED SAWDUST DIETS AS REPLACEMENT FOR MAIZE OFFAL	FED 336
48	EFFECT OF BODY WEIGHT AND METHOD OF CASTRATION ON THE COLD AND HOT CARCASS CHARACTERISTICS OF SAVANNA BROWN GOATS	341
49	INFLUENCE OF CLIMATE EXTREMES ON LIVESTOCKS' DISEASES OCCURRENCE IN BURKINA FASO	347
50	MEAT YIELD AND CARCASS CHARACTERISTICS OF BROILER CHICKENS FED MORINGA OLEIFERA LEAVE POWDER AS ALTERNATIVE TO SYNTHETIC LYSINE	348
51	SEMEN CHARACTERISTICS OF RABBIT BUCKS FED GRADED LEVELS OF NEWBOULDIA LAEVIS LEAF MEAL	356
52	PROXIMATE COMPOSITION OF RAW AND PROCESSED FULL-FAT LEBBECK (ALBIZIA LEBBECK) SEEDS	364
53	GINGER (<i>ZINGIBER OFFICINALE</i>) AS FEED SUPPLEMENT: INFLUENCE ON GROWTH PERFORMANCE AND HEALTH OF GROWING RABBITS – A REVIEW	368
54	ASSESSMENT OF MIXTURES MEAL OF BREWERS DRIED GRAINS AND SORGHUM BREWERS DRIED GRAINS ON GROWTH AND NUTRIENT DIGESTIBILITY OF WEANER RABBITS	374
55	MODELLING OF PHENOTYPIC TRAITS AS DETERMINANTS OF BREEDING POTENTIALS CATTLE UNDER LOW EXTERNAL INPUT	OF 380
56	ASSESSMENT OF MATING PROFILE OF RED SOKOTO BUCKS ADMINISTERED VARYING DOSAGE OF ETHANOLIC EXTRACT OF TIGER NUT (CYPERUS ESCULENTUS)	386

57	FINISHER CHICKENS FED GROUNDNUT-COWPEA SHELL BASED DIETS	391
58	GROWTH PERFORMANCE AND EGG PRODUCTION OF JAPANESE QUAILS FED DIFFERENTLY PROCESSED (<i>LEUCAENA LEUCOCEPHALA</i>) BASED DIETS	396
59	EFFECTS OF THE METHANOL LEAF EXTRACT OF NEWBOULDIA LAEVIS ON OESTROGE LEVELS DURING PREGNANCY IN RABBIT DOES IN KADUNA STATE, NIGERIA	N 402
60	GROWTH AND BODY MORPHOMETRIC PARAMETERS OF BROILER CHICKENS ORALLY ADMINISTERED VARYING LEVELS OF LEMONGRASS EXTRACT, AT FINISHER PHASE	409
61	THERMOREGULATORY RESPONSES IN PERIPARTURIENT SAHELIAN AND WEST AFRICADWARF GOATS DURING THE HOT-DRY SEASON IN THE NORTHERN GUINEA SAVANNA ZONE OF NIGERIA	
62	GROWTH PARAMETERS OF RATS FED VARYING RATIONS OF EDIBLE CHITOSAN-STARGETILMS PACKAGING	CH 423
63	EFFECT OF BREED AND SEX ON PELT GROWTH OF NEW ZEALAND WHITE AND CHINCHILLA GIGANTAS RABBITS	433
64	WATER QUALITY ASSESSMENT OF THE PROPOSED KWADNA RESERVOIR WITHIN GID, KWANU MAIN CAMPUS, FEDERAL UNIVERSITY OF TECHNOLOGY, MINNA, NIGER STANIGERIA	
65	PHYLOGENY BETWEEN CLARIAS GARIEPINUS AND HETEROBRANCHUS BIDOSALIS INFERRED FROM SINGLE NUCLEOTIDE POLYMORPHISMS (SNPS) DNA MARKERS	447
66	EFFECTS OF PHOTOPERIOD AND FEEDING RATE ON THE GROWTH PERFORMANCE AI FEED UTILIZATION OF CLARIAS GARIEPINUS	ND 452
67	IMPACT OF COVID-19 INDUCED LOCKDOWN ON AQUACULTURE PRODUCTION IN MINNA, NIGER STATE, NIGERIA	457
68	EVALUATION OF ADAPTIVE PLASTICITY IN WILD SAROTHERODON GALILAEUS AND COPTODON ZILLII IN CONCRETE POND	464
69	EFFECT OF PROCESSED SELECTED MEDICINAL PLANTS DIETS ON HAEMATOLOGICAL PARAMETERS OF <i>CLARIAS GARIEPINUS</i> (BURCHELL, 1822)	472
70	ASSESSMENT OF INORGANIC FERTILIZER ON FRESHWATER FISH CULTURE IN NIGERIA	480
71	UTILIZATION OF MAIZE BRAN CHEMICAL HYDROLYSATE USING MINERAL ACID IN THE	• •

	PRATICAL DIETS OF CLARIAS GARIEPINUS	486
72	SENSORY AND PROXIMATE COMPOSITION OF 'BISCUIT' PRODUCED FROM SOME LES VALUED DRIED FISH POWDER	SS 492
73	PROXIMATE AND SENSORY ASSESSMENT OF AADUN PRODUCED FROM MAIZE AND GROUNDNUT PASTE	499
74	NUTRI-CEREALS AS POTENTIAL FUNCTIONAL INGREDIENTS: CHARACTERIZATION AN VALORIZATION	D 506
75	MICROBIOLOGICAL QUALITY OF DEVELOPED WHEAT BISCUIT FORTIFIED WITH EGGSHELL CALCIUM	507
CRC	P SCIENCE/PRODUCTION	512
76	EVALUATION OF SINGLE AND MIXED VIRUS-INOCULATED BAMBARA GROUNDNUT LANDRACES FOR NODULATION AND NITROGEN FIXATION	513
77	EFFECTS OF JATROPHA CURCAS LEAF EXTRACT ON THE GROWTH CHARACTERISTICS NEMATODE ASSOCIATED WITH TOMATO (SOLANUM LYCOPERSICUM)	AND 520
78	EFFECT OF FOLIAR FERTILIZER APPLICATION ON PHYSIOLOGICAL CHARACTERISTICS A	AND 529
79	YIELD AND POD SHATTERING BEHAVIOUR OF SOME SOYBEAN GENOTYPES ACROSS LOCATIONS IN NIGERIA	534
80	EFFECTS OF ROOT KNOT NEMATODE (<i>MELOIDOGYNE INCOGNITA</i>) ON THE GROWTH PERFORMANCE OF OKRA (<i>ABELMOSCHUS ESCULENTUS</i>) CULTIVARS IN MINNA, NIGI	
81	EFFICACY OF SOME BOTANICAL EXTRACTS ON THE MANAGEMENT OF FALL ARMYWORM (SPODOPTERAFRUGIPERDA).E. SMITH) ON MAIZE (ZEA MAYS L.)	555
82	RESPONSE OF CORCHORUS OLITORIUS (JUTE MALLOW) CULTIVARS INFECTED WITH ROOT-KNOT NEMATODE (MELOIDOGYNE INCOGNITA) TO PRE-SOWING TREATMENT	Г564
83	EFFECT OF FRUIT AGES ON MOTHER- PLANT SEED QUALITY OF "EGUSI" MELON (CUCUMEROPSIS MANNI NAUDIN) CULTIVARS	570
84	GRAIN YIELD OF EARLY MATURING PRO-VITAMIN A (PVA) MAIZE INBRED LINES UND STRIGA INFESTATION AND OPTIMAL CONDITIONS	ER 576

85	ESCULENTUS (L.) MOENCH)	584 584
86	MORPHOLOGICAL AND SEEDS DIVERSITY IN DRUMSTICK PLANT (<i>MORINGA OLEIFERI</i> LAM.) GERMPLASM FROM NORTHERN NIGERIA	4 589
87	EFFECT OF VARIETY AND INTRA-ROW SPACING ON GROWTH AND YIELD CHARACTER OF HYBRID MAIZE (<i>ZEA MAYS</i>) IN FCT	S 596
88	RESPONSE OF HYBRID LINES OF SWEET POTATO TO THE APPLIED PLANT EXTRACTS, WOOD ASH AND NPK FERTILIZER IN ABUJA, NIGERIA	603
89	RESPONSE OF GROWTH AND YIELD OF MAIZE (ZEA MAYS L.) TO LIME, INORGANIC A ORGANIC FERTILIZER IN MOKWA, NIGER STATE OF NIGERIA	ND 611
90	DETERMINATION OF IRON AND ZINC CONTENTS OF RICE VARIETIES AND LAND RACE NIGERIA USING ATOMIC ABSORPTION SPECTROMETRY ANALYSIS (AAS)	S IN 620
	EFFECT OF FRUI <mark>T</mark> AND MOTHER-PLANT AGES ON THE VIABILITY AND <mark>LO</mark> NGEVITY OF CELMOSCHUS ESCULENTUS L. MOENCH) SEEDS	KRA 628
92	INFLUENCE OF TEMPERATURE AND AIR VELOCITY ON THE MOISTURE DIFFUSIVITY A ACTIVATION ENERGY IN DRYING OF AFRICAN YAM BEAN (SPHENOSTYLIS STENOCAR TUBER	
93	PROFILING OF SOYBEAN (GLYCINE MAX (L.) GENOTYPES WITH HIGH FUNCTIONAL OIL CONTENT FOR INDUSTRIAL AND DOMESTIC APPLICATIONS	L 640
94	EFFECT OF SEED SIZE AND POSITION OF SEEDS IN POD ON THE SEED VIGOUR OF FLU PUMPKIN (TELFARIA OCCIDENTALIS HOOK)	TED 648
95	EFFECTS OF MOISTURE STRESS AT DIFFERENT GROWTH STAGES ON THE PERFORMA OF ONION (<i>ALLIUM CEPA</i> L.) VARIETIES	NCE 654
96	WEEDING FREQUENCY EFFECT ON GROWTH AND YIELD OF MAIZE IN SOUTHERN-GUINEA SAVANNAH	663
97	EFFECT OF FOLIAR FERTILIZER APPLICATION ON PHYSIOLOGICAL CHARACTERISTICS A HERBAGE YIELD OF AMARANTHUS AND CORCHORUS	AND 670
98	EFFECT OF FOLIAR FERTILIZER APPLICATION ON PHYSIOLOGICAL CHARACTERISTICS A HERBAGE YIELD OF AMARANTHUS AND CORCHORUS	AND 675
99	RESPONSE OF SESAME (SESAMUM INDICUM L.) TO DIFFERENT NUTRIENT SOURCES	IN

	ANYIGBA, KOGI STATE NIGERIA	680
100	EFFECTS OF INTEGRATED NUTRIENT MANAGEMENT ON WEED INFESTATION AND YIELD OF SOYBEAN IN SOUTHERN GUINEA SAVANNA ZONE OF NIGERIA	692
101	EFFECT OF MILLED GROUNDNUT SHELL AS A NUTRIENT SOURCE ON THE NUTRITIONAL QUALITY OF PEPPER IN COMPARISON WITH OTHER NUTRIENT SOURCES	699
102	REACTION OF RICE VARIETIES TO RICE BLAST - AN INSIGHT INTO UNDERSTANDING RICE RESILIENCE TO CLIMATE INDUCED RICE DISEASES	G OF 706
103	ALLELOPATHIC EFFECT OF MINTWEED (HYPTIS SUAVEOLENS (L.) POIT) GREEN AN BROWN LEAVES AQUEOUS EXTRACT ON SEED GERMINATION AND SEEDLING GROWTH OF COWPEA	D 717
104	PROXIMATE COMPOSITION OF SOME NEW SWEET POTATO VARIETES GROWN AT GIDAN KWANO, MINNA, NIGER STATE, NIGERIA	723
105	EVALUATION OF SACCHARUM <i>OFFICINARUM</i> GENOTYPES FOR JUICE QUALITY, CA	NE 731
106	IMPACTS OF SUGARCANE GENOTYPES AND WEED MANAGEMENT PRACTICES ON WEED DRY MATTER AND SUGARCANE PRODUCTIVITY	737
107	ISIGHTS INTO PREVALENCE AND DISTRIBUTION OF VIRUSES INFECTING MELON IN GEORGIA, UNITED STATE	744
108	GROWTH AND YIELD RESPONSES OF MAIZE TO PARTIAL SUBSTITUTION OF INORGANIC NITROGEN WITH FARM YARD MANURE AT GIDAN-KWANO	745
109	RESPONSE OF GROWTH AND YIELD OF MAIZE (<i>ZEA MAYS</i> L.) TO LIME, INORGANIC AND ORGANIC FERTILIZER IN MOKWA, NIGER STATE OF NIGERIA	753
110	UNDERSTANDING THE INTENTION TO USE GOOD AGRICULTURAL PRACTICES ON VEGETABLE FARMS – A COMPARATIVE STUDY OF FARMERS IN PUNJAB, INDIA AN NAKURU, KENYA	D 761
111	DETERMINATION OF DRY SEASON IRRIGATION WATER QUALITIES FOR VEGETABL PRODUCTION IN MINNA, NIGERIA	<u>-</u> 768
112	EFFECT OF DAYS AFTER ANTHESIS AND STORAGE PERIOD ON SEED QUALITY OF TO OKRA (ABELMOSCHUS ESCULENTUS L. MOENCH) VARIETIES	NO 778

113	ASSESSMENT OF SELECTED SOIL FERTILITY PARAMETERS ALONG A TOPOSEQUENC UNDER INTENSIVE CROP PRODUCTION AND IMPLICATIONS FOR SUSTAINABLE MANAGEMENT	Æ 782
114	EFFECT OF DAYS AFTER ANTHESIS AND STORAGE PERIOD ON SEED QUALITY OF TV OKRA (<i>ABELMOSCHUS ESCULENTUS</i> L. MOENCH) VARIETIES	VO 790
SOIL SO	CIENCE / ENVIRONMENTAL MANAGEMENT	794
115	EFFECT OF BIOCHAR AND NANOPARTICLES ON THE MICROBIAL DEGRADATION OF CRUDE OIL CONTAMINATED SOIL	795
116	INDIGENOUS KNOWLEDGE AND COPING CAPACITY OF FARMERS FOR CLIMATE CHANGE ADAPTATION: A CASE STUDY OF AGAIE LOCAL GOVERNMENT AREA, NIGHTSTATE. NIGERIA	ER 803
117	ASSESSMENT OF FERTILITY STATUS OF SOME SELECTED FADAMA SOILS IN BIDA LOCAL GOVERNMENT AREA OF NIGER STATE, NIGERIA	807
118	EFFECTS OF DIFFERENT NITROGEN SOURCES ON THE GROWTH OF MAIZE AND SOYBEAN IN AN INTERCROPPED SYSTEM	813
119	EVALUATION OF IRRIGATION SUITABILITY FOR SURFACE IRRIGATION IN BAKAJEBA, NIGER STATE, NIGERIA	, 820
120	EFFECTS OF BIOCHAR AND NITROGEN FERTILIZER ON SOIL ORGANIC CARBON, NUTRIENT RETENTION AND MAIZE PERFORMANCE	826
121	IMPLICATION OF LAND SURFACE TEMPERATURE AS A DRIVER OF LAND COVER CHANGES ON THE URBAN DYNAMICS OF ABUJA NIGERIA	833
122	PHOSPHATE MOBILIZATION BY ADDITION OF THREE ANIMAL MANURE SOURCES II THE SOILS OF KARU, NASARAWA STATE	N 840
123	COMBINATION OF ORGANIC MANURE, INORGANIC FERTILIZER AND BIO-FERTILIZE ON SOYBEAN [(GLYCINE MAX L. (MERRIL)] GROWTH	R 845
124	SPATIOTEMPORAL ASSESSMENT OF THE DYNAMICS OF DROUGHT IN NORTHEAST NIGERIA USING REMOTE SENSING TECHNIQUES	853

64 WATER QUALITY ASSESSMENT OF THE PROPOSED KWADNA RESERVOIR WITHIN GIDAN KWANU MAIN CAMPUS, FEDERAL UNIVERSITY OF TECHNOLOGY, MINNA, NIGER STATE, NIGERIA

¹ Ibrahim, S. U., ² Ibrahim, A., ³ Yusuf, J., ⁴ Adama, S. B., and ⁵ Abdulmumini, A.N.

^{1,3 & 5} Department of Water Resources, Aquaculture and Fisheries Technology, School of Agriculture and Agricultural Technology, Federal University of Technology, Minna, Niger State, Nigeria

² Marine Biology, School of Life Sciences
College of Agriculture, Engineering and Science
Westville Campus
University of Kwazulu-Natal
South Africa

⁴ Department of Animal Biology *Federal University of Technology, Minna Niger State, Nigeria

Corresponding author: ¹Ibrahim Saratu Usman suibrahim@futminna.edu.ng Phone: +234(0)7039025846, +234(0)8055957138

Abstract

The study assessed the Water Quality parameters of the Kwadna Reservoir located at Federal University of Technology, Minna guided by the following objectives: determine the variation of the physico-chemic<mark>a</mark>l parameter across the months, the stations and to pro<mark>vi</mark>de baseline data of the various physical and chemical properties of water. The Physico-chem<mark>ic</mark>al parameters of the river were studied for twelve weeks. Water samples was taken from five (5) stations in the reservoir, biweekly for the period of study. Samples collected was taken to the department of Water Resources, Aquaculture and Fisheries Technology (WAFT) laboratory for analysis. The results of the research depict that there was no significant variation in the mean values of physico-chemical parameters observed across the Station (p>0.05), pH range from 7.05 ± 0.03 -7.36 ±0.03 , Conductivity 281.5±45.5-311.7±44.9 µS/cm, Total alkalinity 115.7±22.0 -174.0±18.9, Total hardness 48.1±9.5-57.4±11.3mg/L, Calcium 17.3±5.4-24.6±4.6mg/L, Magnesium 4.0±2.4-8.6±2.9 mg/L, TDS 177.2±28.3-199.3±28.9mg/L,BOD 2.7±0.1-3.1±0.4mg/L, Dissolved oxygen 4.9±0.6-5.7±0.1mg/L, Carbon dioxide 1.7±0.9-3.4±2.3mg/L, BOD 2.7±0.1-3.1±0.4mg/L. Except Temperature and COD that has significant difference (p<0.05) range between 27.0±0.0- $28.5\pm0.5^{\circ}C$ and $18.1\pm2.8-104.9\pm45.9$ mg/L respectively. For monthly variations most parameters had significant difference (p<0.05) except Temperature and BOD that has no significant difference (p>0.05). Most of the physico-chemical parameters studied were within WHO range set standard for optimal fish production and survival. The mean temperature of the Reservoir (29.46°C) is in line with FAO (2006), which states that temperature requirements of $25^{\circ}\text{C}-30^{\circ}\text{C}$ is optimum for fish growth. This shows that temperature in the proposed Kwadna Reservoir is suitable to support the growth of fish Therefore, constant monitoring of the reservoir should be encouraged.

Keywords: Water Quality, Kwadna, Reservoir, Physico-chemical

INTRODUCTION

Water is one of the vital needs of all living beings. The quality of water usually described its physical, chemical and biological characteristics. Hence it becomes necessary to find out its the suitability for drinking, irrigation, fishing and Industrial purpose. The availability of good quality water is a necessary feature for preventing diseases and improving quality of life (Oluduro and Aderiye, 2007).

Dam reservoirs, which are an integral part of civilization development, have many features that distinguish them from natural lakes or rivers, hence they constitute a different category of surface water reservoirs. Their most important functions include water collection for municipal and agricultural purposes or flood protection. They are also used for recreational purposes, hydropower and fishing.

Water quality deterioration in reservoirs usually comes from excessive nutrient inputs, eutrophication, acidification, heavy metal contamination, organic pollution and obnoxious fishing practices. The effects of these "imports" into the reservoir do not only affect the socio-economic functions of the reservoir negatively, but also bring loss of structural biodiversity of the reservoir.

Water quality can generally be defined as chemical, physical and biological waters that are characteristic with respect to their suitability for a given use (Pawar, S.S, 2017). He further stated that the physico-chemical properties of water refer to the joint of physical characteristics and chemical composition of water body which include turbidity, colour, odour, temperature, pH, conductivity, dissolve oxygen (DO), Biological Oxygen demand (BOD), hardness, alkalinity, nitrate, chloride, CO₂, etc. Each of the designated uses has different definitions of the chemical, physical and biological standards needed to support the use.

These physiochemical compositions of water bodies need to be evaluated and compared with standards as a basis for the identification of the causes of the change in the water quality, high mortality or low population of aquatic animals in the dam reservoir.

Water quality parameters examined were pH, alkalinity, calcium, conductivity, magnesium, total hardness, Dissolved carbon dioxide, Chemical Oxygen Demand (COD) Total dissolved Solids (TDS), dissolved oxygen (DO), biochemical oxygen demand (BOD).

MATERIALS AND METHODS

Study area: The study was carried out at proposed Kwadna reservoir located within Federal University of Technology Minna, Bosso Local Government Area of Niger State, Nigeria which lies in between the latitude of 9^o 30' 40" N and longitude of 6^o 24' 50" E in the northern guinea savannah vegetation zone of Nigeria.




Figure 1: Nigeria indicating Niger State and the study area




Figure 2: The study area indicating Sampling Stations

Sample collection and preparation: The samples were collected bi-weekly for three (3) months March, April, May from five different stations (Figure 2). Inlet of the Reservoir, open water of the reservoir bank of the reservoir were anthropogenic activities and opposite sides of the bank. The samples collected were analyzed for pH, Temperature, Alkalinity, Calcium, Conductivity, Magnesium, Total Hardness, Dissolved Carbon Dioxide, Chemical Oxygen Demand (COD) Total dissolved Solids (TDS), dissolved oxygen (DO), Biochemical Oxygen Demand (BOD).

Determination of pH: The pH of the water samples was determined using the pH meter. It was standardized with a buffer solution of pH range between 4, 7 and 9.

Measurement of temperature: This was carried out in-situ at the site of sample collection using a mobile thermometer. This was done by dipping the thermometer into the sample and recording

the stable reading

Determination of conductivity: This was done using conductivity meter. The probe was dipped into the samples until a stable reading was obtained and recorded.

Determination of Alkalinity: 50mL of the sample was pipetted into a clean 250mL conical flask. Two drops of methyl red indicator were then added and the solution titrated against a standard 0.01M NaOH solution to a pink end-point. (American society for testing and Materials, 1982). Total alkalinity (mg/L) = V x M x 50,000 / mL of sample used Where V = volume of acid used M = Molarity of acid used

Determination of total dissolved solids (TDS): by Gravimetric Method: A portion of water was filtered out and 10mL of the filtrate measured into a pre-weighed evaporating dish. Following the procedure for the determination of total solids above, the total dissolved solids content of the water was calculated. Total dissolved solids (mg/L) = (W2-W1) mg x 1000 mL of filtrate used. Where W1 = initial weight of evaporating dish W2 = Final weight of the dish (evaporating dish + residue). Determination of Dissolved Oxygen: This was done using Winkler's method. Do bottle were used to collect the water sample at the sample site, 0.5ml of reagent 1(Magnose sulphate),and 0.5ml of reagent 2 (Potassium Hydroxide +Potassium Iodide) were added immediately into the collected sample,before it was transported to the laboratory 10ml of the sample was measured from the Do bottle into a conical flask,5drops of concentrated H₂SO₄ was added,5 drops of starch indicator was also added and was titrated with Sodiumthiosulphate until color changes from blue black colorations to colorless.

DO $(mg/l) = TV \times 0.025 \times 8 \times 1000 \ 20 \text{mls}$

Calculation DO (mg/L) = $16000 \times M \times V \times V2/V1$ (V1-2) Where = Molarity of thiosulpahte used. V = volume of thiosulphate used for titration V1 = Volume of bottle with stopper V2 = Volume of aliquot taken for titration.

Determination of Biochemical Oxygen Demand (BOD): The method involves filling the samples to overflowing, in an airtight bottle of the specified size.

Determination of total Hardness: 25mL of the samples was placed in different clean 250mL conical flask. To this were added 3mL of ammonium chloride in concentrated ammonia buffer (NH4CL/conc.NH3) and 2 drops of Eriochrome Black T indicator. This was titrated against 0.01M

EDTA solution until there was a color change from violet to blue.

Calculation: Hardness in mg/L CaCO3= V x M x 1000 mL of sample used Were M = Molarity of

EDTA Used V = Volume of EDTA used

RESULTS AND DISCUSSIONS

Physico-chemical parameters Variation across the stations and months

The findings of the station variations deduce that some parameters had significant variation from each other while some had no significant variation. The mean temperature of the Reservoir recorded (29.46°C) is in line with FAO (2006), which states temperature requirements at the range of $25^{\circ}\text{C} - 30^{\circ}\text{C}$ is optimum for fish growth, Conductivity was higher during the month of March $385\mu\text{S/m}$ and lowest during the month of May this is as a result of increased water evaporation and emergence from wind while mean value of Dissolved oxygen of 5.19 mg/l obtained agree with Cline (2012) value of 3-10mg/L.

The mean BOD values of 2.91mg/l is in line with the work of Boyd (2003), which states that the optimal BOD values ranges between 3–20 mg/l. Mean value of hardness was 53.91mg/l. indicate that the Reservoir water is a soft water which is within the desirable limit (WHO, 1984) that states hardness values below 300mg/l is potable water. For total Alkalinity the range value was 147.43mg/l this fell within the permissible range reported by Cline (2012), to be between 50-250mg/L suitable for fish growth.

The mean value for Calcium during the course of the research was 20.11mg/l and it fell within the permissible range as posited by WHO (1984), the maximum permissible limits for calcium in drinking water is 75mg/l. Thus the calcium level in this research is within the permissible limits. Though some parameters of the stations fall within the required range that support fish in the said period of March to May, some others are out of range which calls for greater attention to correct the causes to safeguard the future of fish survival in the Reservoir.

Table 1: Monthly variations of physico-chemical parameters of Proposed Kwadna reservoir

Parameters	March	April	May
pН	7.34±0.02 ^a	7.05±0.03 ^b	7.36±0.03a
Temperature (°C)	28.5±0.5 a	27.0±0.0 a	28.0±0.0 a
EC (μS/cm)	$346.5 \pm 10.84^{\ a}$	$336.80{\pm}10.63^a$	206.60 ± 6.40^{b}
TDS (mg/l)	216.00 ± 9.67^{a}	215.30±6.83 ^a	132.00 ± 3.98^{b}
Alkalinity(mg/l)	162.50±12.49 ^b	156.60±8.72 a	118.20 ± 18.35
Hardness (mg/l)	67.69±4.43 a	61.24±1.11 a	32.77 ± 1.62^{b}
Calcium (mg/l)	$20.75{\pm}2.58^a$	$26.32{\pm}2.01^a$	13.66 ± 0.72^{b}
Magnesium(mg/l)	12.85 ± 1.17^a	2.87 ± 0.95^{b}	4.03 ± 0.57^{b}
DO (mg/l)	4.70 ± 0.29^{b}	5.12 ± 0.17^{b}	5.19 ± 0.09^{a}
DCO2 (mg/l)	5.28±0.83 ^a	1.59±0.14 ^b	1.08 ± 0.20^{b}
BOD (mg/l)	2.71 ± 0.13^{a}	2.96±0.13 ^a	3.08 ± 0.21^{a}
COD (mg/l)	58.24±27.27 ^a	23.73±6.84 b	59.20±11.87 ^a

Mean in the same row followed by the same superscript are not significantly different (P>0.05) Mean not followed by the same superscript are not significantly different (P<0.05)

Table 2: Stations variation of physico-chemical parameters of Proposed Kwadna reservoir

Parameters	Station 1	Station 2	Station 3	Station 4	Station 5
Ph	7.26 ± 0.0^{a}	7.31±0.09 a	7.25±0.12 a	7.23±0.12 a	7.20±0.11 a
Temperature (°C)	28.5±0.5 b	27.0±0.0 a	28.0±0.0 a	28.2±0.0 b	27.5±0.5 a
EC (μS/cm)	297.3±51.1	311.7±44.9ª	291.8±49.7ª	299.8±41.0	281.5±45.5
TDS (mg/l)	189.8±32.8	199.3±28.9ª	177.2±28.3ª	191.8±26.3	180.7±29.1
Alkalinity(mg/l)	174.0±18.9	123.5±22.5 ^a	162.3±17.9 ^a	161.7±6.4 a	115.7±22.0
Hardness (mg/l)	57.4±11.3 a	53.8±13.5 a	54.9±20.6 a	55.4±10.8 ^a	48.1±9.5 a
Calcium (mg/l)	18.9±3.0 a	19.3±4.2 a	20.5±4.1 a	24.6±4.6 a	17.3±5.4 a
Magnesium(mg/l)	8.6±2.9 a	6.5±4.3 a	6.9±2.5 a	4.0±2.4 a	6.9±3.8 ^a
DO (mg/l)	5.1±0.4 a	4.9±0.6 a	5.1±0.2 a	5.1±0.3 a	5.7±0.1 ^a
DCO2 (mg/l)	1.7±0.9 ^a	3.4±2.3 ^a	2.3±0.8 a	2.7±1.2 a	3.0±1.5 ^a
BOD (mg/l)	3.1±0.4 a	2.9±0.1 a	2.7±0.1 a	3.0±0.3 ^a	2.9±0.2 a
COD (mg/l)	50.9±11.0	18.1±2.8 ^b	35.2±17.6 ab	26.1±2.3 ^b	104.9±45.9

Mean in the same row followed by the same superscript are not significantly different (P>0.05) Mean not followed by the same superscript are not significantly different (P<0.05)

Conclusion

The study on the assessment of physic-chemical parameters of Proposed Kwadna dam reservoir, Minna, Gidan Kwano Niger State was carried out bi-weekly for the period of three months in order to provide baseline information on the ecological status of the Reservoir. The physico-chemical parameters of the Reservoir varied with months and stations. The variations may be due to change in weather cycle during the study period.

REFERENCES

- Boyd C.E (2003) Guildelines for aquaculture effluent management at the farm-level. Aquaulture 226:101-112.https://doi.org/10.1007/978-3-319-17446-4
- Cline, D. (2012). Water quality in aquaculture. Freshwater Aquaculture, Alabama Cooperative Extension System, Auburn University.
- Food and Agricultural Organization FAO (2006): The state of the worlds land and water resources for food and agriculture. Stakeholder oriented valuation to support water resource management pp 71 https://www.fao.org
- Oluduro AO., Aderiye BI. (2007). Effeciency of moringa Oleifera seed extract on the microflora of surface and ground water. *Journal of Plant Sciences*. 6:453-438
- Pawar, S. S. (2017). Analysis of some physicochemical parameters and their effect on the productivity of fishes in two different types of aquatic bodies of Unamganon and Seipargaon of Patharkandi block of Karimaganj district, Assam, India. *International Journal of Life Sciences*, 5(4): 587-592
- World Health Organisation, (2011). Guidelines for drinking-water quality 4th ed. ISBN 978 92 4 154815 1 World Health Organization, Geneva 27, Switzerland. Accessed July 2019. http://www.who.int