



NIGERIAN JOURNAL OF CROP SCIENCE

VOLUME 4 NUMBER 1, JANUARY 2017

ISSN: 2350 - 2487

SPONSORED AND PUBLISHED BY:

CROP SCIENCE SOCIETY OF NIGERIA

MEMBERS OF THE EDITORIAL BOARD

Editor In-Chief

Professor P. E. Ogbonna
Dept. of Crop Science
University of Nigeria, Nsukka

Dr. Francis Chuks Ogbonnaya
Grains Research and Development
Corporation Level 1, 40 Blackall Street
Barton ACT 2600, Australia
Email: Francis.Ogbonnaya@grdc.com.au

Bello, Omolaran Bashir (Ph.D.) MGSN, MSAN,
Department of Biological Sciences,
Fountain University, Osogbo,
Osun State, Nigeria

Professor Uche Cyprian Amalu, FSSN
Department of Soil Science
Faculty of Agriculture, Forestry and Wild
Life Resources Management,
University of Calabar, Calabar, Nigeria.
E-mail: ucamalu@yahoo.co.uk

Dr. Lekan Sakariyawo
Department of Plant Physiology & Crop
Production. College of Plant Science &
Crop Production. Federal University of
Agriculture, Abeokuta.
Email: adetanwa@yahoo.co.uk

Prof. Gbadebo Olaoye
Department of Agronomy, University of
Ilorin, P.M.B. 1515, Ilorin, Nigeria.
Email: debolaoye@gmail.com

Prof. K. P. Baiyeri
Dept. Of Crop Science
University of Nigeria, Nsukka.
Email: Paul.baiyeri@unn.edu.ng

Dr. Nyaudoh Ukpabio Ndaeyo
Department of Crop Science,
University of Uyo, Uyo, Nigeria
Email: dr_nundaeyo@yahoo.com

Dr. Ibrahim Dugje Yakamba
Department of Crop Production,
University of Maiduguri, P.M.B 1069,
Maiduguri, Borno State, Nigeria
Email: dugje2011@gmail.com,

Prof. N. I. Odiaka
Dept of Crop Production
University of Agriculture
Makurdi
Email: niodiaka@yahoo.com

Prof. M. I. Uguru
Dept. of Crop Science
University of Nigeria, Nsukka
Michael.uguru@unn.edu.ng

Professor Peter Odo
Dept of Plant Breeding and Seed Science
University of Agriculture, Makurdi
Email: petereodo@yahoo.com

Dr. Aransiola Aaron Opeyemi
Dept. of Crop Production, Obafemi
Awolowo University, Ile-Ife
Email: bamujo@oauife.edu.ng

Professor Cosmas Muoneke
College of Crop and Soil Sciences
Michael Okpara University of Agriculture,
Umudike.
Email: muoneke.cosmas@mouau.edu.ng.

Prof. D.A. Okpara
College of Crop and Soil Sciences
Michael Okpara University of Agriculture,
Umudike.
Email: daokpara@yahoo.com

Professor P. I. Okocha
College of Crop and Soil Sciences
Michael Okpara University of Agriculture,
Umudike.
Email: okochaiwunor@gmail.com.

Prof. M. O. Adeyemo
Dept. of Crop Production
University of Agriculture
Makurdi

Professor B. C. Echezona
Dept. Of Crop Science
University of Nigeria, Nsukka
Email: bonaventure.echezona@unn.edu.ng

INSTRUCTION TO AUTHORS

The journal publishes original papers in Agronomy, Crop Productions, Genetics and Breeding, Crop Protection, Crop Germplasm, Soil Science, Agroforestry, Organic Farming, Post harvest technology, Crop Biotechnology. Manuscript in English should be prepared in double-spaced typing on one side of A4 paper. Papers should be subdivided into introduction, materials and method, results, and discussion and should contain an abstract that is not more than 250 words. Key words not exceeding (5) words should be included below the abstract. Tables and figures should be supplied on separate sheet. Photographs must be black and white. Every table should have a short descriptive caption at the top and carry no vertical lines or embellishments. The captions for the figures should be at the bottom. Numbering for tables and figures should be in Arabic numerals, that is, table 1, 2, ... and figure 1, 2, ..., etc. The appropriate positions of all the illustrations should be indicated in the text. The title which must be short and concise, should head the article. Author's name, consisting of the surname, followed by the initials should be written below the title. This should be followed by the author's affiliations with full address. References in the text must be given by the author's name with the year of publication. The literature cited must be given alphabetically according to the author's names at the end of the paper as follows:

Journals: Jana, S. Khanguata, B. S. (1986) Conservation of diversity in bulk production of barley, *Euphytica* 35: 761-776

Books: Leopold, A. C. (1989), Plant growth and development, 2nd ed. Tata McGraw Hill, New Delhi. pp. 55-60

SUBMISSION OF MANUSCRIPT

Manuscript, typed in character size 12, doubled spaced, with margin all round on A4 paper should be submitted in triplicate with the processing charge of N1,500, \$10 and \$30 to be paid on submission by contributors from within Nigeria, other parts of Africa, and outside Africa, respectively. Similarly, there will be publication charge of N15,000 and \$100 for accepted articles from within Nigeria and foreign subscribers, respectively. All payments are to be made into CSSN Account No. 2022702825, First Bank of Nigeria Plc. Nsukka Branch.

TABLE OF CONTENTS

Growth Analysis of Seven Sweetpotato Genotypes Grown in Lowland Humid Tropics of Southeast Nigeria <i>Eke-Okoro, Ogechi and Mbah, Emmanuel Ukaobasi</i>	1
Heritability and Correlation Studies of Intergeneric Hybrids of 'Egusi' Melon (<i>Colocynthis Citrillus</i> L.) and Watermelon (<i>Citrillus lanatus</i> Thumb). <i>Orji, V.O., Ogbonna, P. E. Onyia, V. N. Okechukwu, E. C.</i>	10
Influence of Poultry Manure and Phosphorus Fertilizer on Growth and Yield of Okro (<i>Abelmoschus Esculentus</i> , (L.) Moench) <i>Olla, N. O., Adekunle, I. O., Bello, W. B., Aribisala, L. A., Ogunjinmi, S. O and Olanlyan, M. I</i>	21
Genotype by Environment Interaction of Early and Intermediate Maturing Maize Varieties in the Southern Guinea Savanna of Nigeria <i>Gbadamosi A., Olaoye G., Akinwale R. A., Bankole F.A. and Takim F.O.</i>	27
Response of Growth and Yield on different Management Practices of Cucumber (<i>Cucumis Sativum</i> L.) in Rivers State, Nigeria <i>Orluchukwu, J. A. and Nwuke, Chidinma</i>	36
Effect of Phosphorus and Starter Nitrogen on Cowpea (<i>Vigna Unguiculata</i> L. Walp) Performance in Southern Guinea Savanna of Nigeria. <i>Afolabi, S.G, Adeboye, M.K.A., Lawal, B.A., Asin, E.C. and Uzoma, A.O</i>	40
Estimates of Genetic Variability in Native and Elite Cucumber (<i>cucumis sativus</i> L.) Varieties <i>Manggoel, W., Uguru M. I. and Ogbonna, P. E</i>	48
Food Quality Characteristics and Effects of Plant Spacing on Performance of New Cassava (<i>Manihotesculentum</i> Crantz) Genotype in a derived Savannah Agro-Ecology of Southeast Nigeria <i>Eze, S. C., Aba, S. C., Amuji, C. F., Ezeaku, P. I. and Ogbonna R.</i>	56
Effect of Pinching and Watering Frequencies on Growth and Flowering of Marigold (<i>Tagetes erecta</i> L.) <i>Ekwu, L. G., Nwanchor, K. Utobo, E.B. and Nwokwu, G. N.</i>	66
The Efficacy of Eco-Friendly Botanicals in the Management of Damping-Off and Downy Mildew Diseases of Cucumber, in Abakaliki, South Eastern Nigeria <i>Utobo, E.B., Ekwu, L.G., Nwokwu, G.N., Nwogbaga, A.C. and Nwanchor, K.</i>	75
Evaluation of Growth and Yield Performances of Three Varieties of Cassava (<i>Manihot Escalenta</i> Crantz) and Four Pigeonpea (<i>Cajanus Cajan</i> (L) Millsp) Genotypes in Intercropping Systems in Nsukka Agroecology <i>Ogbu, C.E.S., Astegbu, J.E. and Onyeonagi, C.C</i>	83

LIST OF CONTRIBUTORS

- Aba, S. C.**
Department of Crop Science,
University of Nigeria, Nsukka
- Adekunle, I. O**
Department of Soil Science Technology,
Oyo State College of Agriculture and
Technology, Igboora, Nigeria
- Adeboye, M.K.A.**
Department of Soil Science and Land Mgt, School
of Agric and Agricultural Technology, Federal
University of Technology, Minna, Niger State,
Nigeria.
- Afolabi, S.G**
Department of Soil Science and Land Mgt
School of Agric and Agricultural Technology
Federal University of Technology, Minna, Nigeria.
- Akinwale R. A.**
Department of Crop Production and Protection,
Obafemi Awolowo University, Ile-Ife, Nigeria
- Aribisala, L. A**
Department of Soil Science Technology,
Oyo State College of Agriculture and
Technology,
Igboora, Nigeria
- Asin, E.C.**
Department of Soil Science and Land Mgt
School of Agric and Agricultural Technology
Federal University of Technology, Minna, Nigeria.
- Asiegbu, J.E.**
Department of Crop Science, University of Nigeria
Nsukka
- Amuji, C. F.**
Department of Crop Science,
University of Nigeria, Nsukka
- Bankole F.A.**
Department of Agronomy,
University of Ilorin, Ilorin, Nigeria
- Bello, W. B**
Department of Soil Science Technology,
Oyo State College of Agriculture and
Technology,
P. M. B. 10, Igboora, Nigeria
- Eke-Okoro, Ogechi**
Department of Agronomy,
Michael Okpara University of Agriculture,
Umudike, Nigeria.
- Eze, S. C.**
Department of Crop Science,
University of Nigeria, Nsukka
- Ezeaku, P. I.**
Department of Soil Science
University of Nigeria, Nsukka
- Ekwu, L.G.**
Department of Crop Prod. and Landscape Mgt.
Faculty of Agric. and Natural Res. Mgt.
Ebonyi State University, Abakaliki, Nigeria
- Gbadamosi A.**
Teaching and Research Farm,
Landmark University, Omu-Aran, Nigeria
- Lawal, B.A.**
Department of Soil Science and Land Mgt
School of Agric and Agricultural Technology
Federal University of Technology, Minna, Niger
State,
- Mbah, Emmanuel Ukaobasi**
Department of Agronomy,
Michael Okpara University of Agriculture,
Umudike, Nigeria.
- Manggoel, W.**
College of Agriculture,
PMB, 001, Garkawa, Plateau State, Nigeria
- Nwogbaga, A.C.**
Department of Crop Prod. and Landscape Mgt.
Faculty of Agric. and Natural Res. Mgt.
Ebonyi State University, Abakaliki, Nigeria
- Nwanchor, K.**
Department of Crop Prod. and Landscape Mgt.
Ebonyi State University, Abakaliki, Nigeria
- Nwokwu G.N.**
Department of Crop Prod. and Landscape Mgt.
Ebonyi State University, Abakaliki, Nigeria
- Nwuke, Chidinma**
Department of Crop and Soil Science,
Faculty of Agriculture, University of Port
Harcourt, Choba, Port Harcourt, Rivers State,
Nigeria
- Orji, V.O.**
Department of Crop Science, University of Nigeria,
Nsukka, Enugu State Nigeria
- Ogbonna, P. E**
Department of Crop Science, University of Nigeria,
Nsukka, Enugu State Nigeria

Onyia, V. N.
Department of Crop Science, University of Nigeria,
Nsukka, Enugu State Nigeria

Okechukwu, E. C.
Department of Crop Science, University of Nigeria,
Nsukka, Enugu State Nigeria.

Olla, N. O
Department of Soil Science Technology,
Oyo State College of Agriculture and
Technology,
Igboora, Nigeria

Olaniyan, M. I
Department of Soil Science Technology,
Oyo State College of Agriculture and
Technology,
Igboora, Nigeria

Ogunjinmi, S. O
Department of Soil Science Technology,
Oyo State College of Agriculture and
Technology,
Igboora, Nigeria

Olaoye G.
Department of Agronomy, University of Ilorin,
P. M. B. 1515, Ilorin, Nigeria

Orluchukwu, J. A.
Department of Crop and Soil Science,
Faculty of Agriculture, University of Port
Harcourt,
Choba, Port Harcourt Rivers State, Nigeria

Ogbonna, P. E.
Department of Crop Science, Faculty of
Agriculture,
University of Nigeria, Nsukka

Ogbonna R.
Department of Crop Science,
University of Nigeria, Nsukka

Ogbu, C.E.S.
Department of Crop Science
University of Nigeria, Nsukka

Onyeonagu, C.C.
Department of Agronomy
Michael Okpara University of Agriculture ,
Umudike

Takim F.O.
Department of Agronomy, University of Ilorin, P.
M. B. 1515, Ilorin, Nigeria

Uguru M. I.
Department of Crop Science, Faculty of
Agriculture,
University of Nigeria, Nsukka

Utobo, E.B.
Department of Crop Prod. and Landscape Mgt.
Ebonyi State University, Abakaliki, Nigeria

Uzoma, A.O.
Department of Soil Science and Land Mgt
School of Agric and Agricultural Technology
Federal University of Technology
P.M.B. 65, Minna, Niger State,
Nigeria.

EFFECT OF PHOSPHORUS AND STARTER NITROGEN ON COWPEA (*VIGNA UNGUICULATA* L. WALP) PERFORMANCE IN SOUTHERN GUINEA SAVANNA OF NIGERIA

Afolabi, S.G*, Adeboye, M.K.A., Lawal, B.A., Asin, E.C. and Uzoma, A.O.

Department of Soil Science and Land Management School of Agriculture and Agricultural Technology Federal University of Technology, P.M.B. 65, Minna, Niger State, Nigeria.

*Corresponding Author e-mail: ooreoluwa14@gmail.com; a.gbolahan@futminna.edu.ng

Phone: +234(0)8035014309

ABSTRACT

Cowpea (Vigna unguiculata L. Walp) is an important grain legume that plays critical roles in diets of millions of people in Africa and other parts of the world. A pot experiment was conducted in 2016 at the Screen House of School of Agriculture and Agricultural Technology, Federal University of Technology, Minna to determine the effect of phosphorus (P) and starter nitrogen (N) on cowpea, SAMPEA 15 (IT99K-573-2-1) performance in Minna, Nigeria. Experimental soils were collected from the Teaching and Research Farm of the same institution. The treatments consisted of five rates of P application (0 kg P ha⁻¹, 10 kg P ha⁻¹, 20 kg P ha⁻¹, 30 kg P ha⁻¹ and 40 kg P ha⁻¹) and four rates of N application (0 kg N ha⁻¹, 10 kg N ha⁻¹, 20 kg N ha⁻¹ and 30 kg N ha⁻¹). The treatments were laid down in a completely randomized design and replicated four times. The results showed that the textural class of the soil was sandy loam, soil reaction was slightly acidic (pH = 6.5), low in organic carbon (4.22 g kg⁻¹), available P (6 mg kg⁻¹) and high total N (0.56 g kg⁻¹). The effect of starter N on plant height, number of leaves, days to flowering and podding, grain and haulms yields was not significant (p>0.05). Application of 10 kg P ha⁻¹ significantly (p<0.05) improved plant height, pods per plant, grain per pod, grains and haulms yield. The interaction between P and starter N was not significant (p>0.05) on all parameters measured except on number of days to flowering and fresh haulms yield. Application of 10 kg P ha⁻¹ is therefore recommended as optimum for the cowpea variety assessed.

Keywords: Cowpea, phosphorous, starter nitrogen, southern Guinea savanna

INTRODUCTION

Cowpea (*Vigna unguiculata* L. Walp) is regarded as one of the most ancient source of food for humans. It is of major importance to the livelihoods of millions of relatively poor people in less developed countries of the tropics (FAO, 2002). Islam *et al.* (2006) emphasized that all parts of the plant used as food are nutritious providing protein and vitamins. Immature pods and peas are used as vegetables while several snacks and main dishes are prepared from the grains. Nigeria is the 2nd highest consumer of cowpea in the world (Egho, 2009). Among the legumes, cowpea is the most extensively grown, distributed and traded food crop consumed in Nigeria. (Ogbo, 2009; Agbogidi, 2010). This is because the crop is of considerable nutritional and health value to man and livestock (Agbogidi, 2010). Cowpea forms a major staple in the diet in Africa and Asian continents (Awe, 2008). Their amino acid complements those of cereals (Asumugha, 2002). The mineral contents of cowpea (calcium and iron) are higher than that of meat, fish and egg and the iron content equates that of milk while the vitamins- thiamin, riboflavin, niacin (water soluble)

and their levels compare with that found in lean meat and fish (Achuba, 2006) which make them very useful in blood cholesterol reduction. Adaji *et al.* (2007) reported that daily consumption of 100-135gm of dry beans reduces serum cholesterol level by 20% thereby, reducing the risk for coronary heart diseases by 40%. It is a good food security item as it mixes well with other recipe (Muoneke *et al.*, 2012). It fixed atmospheric nitrogen through symbiosis with nodule bacteria (Shiringani and Shimeles, 2011). It does well and most popular in the semi - arid of the tropics where other food legumes do not perform well (Sankie *et al.*, 2012). Cowpea is an extremely resilient crop and is cultivated under some of the most extreme agricultural conditions in the world (Muoneke *et al.*, 2012).

Cowpea is the most economically important indigenous African legume crop (Langyintuo *et al.*, 2003). It has vital importance to the livelihood of several millions of people in West and Central Africa. Rural families that make up the larger part of the population of these regions derive

from its production, food, animal feed, alongside income from its production.

Cowpea responds to added nitrogen despite of its capacity to fix nitrogen with rhizobium (Sultana, 2003). Although there are divergent views of nitrogen application to legumes, especially cowpea, results of investigations in the tropics have indicated either no response or significant response to nitrogen fertilizer application (Akinola, 1978). Low level of soil P and N are the major constraints to crop growth and production in nutrients depleted sandy soil of Sub-Saharan Africa (Nyoki and Ndakidemi, 2014). Cowpea (Sampea 15) is high yielding variety, heat tolerance, drought tolerance, Striga and Alectra resistance. Despite all these attributes of the crop, few studies have been carried out to evaluate its effect on P and starter N in Minna. The objective of the study was to determine the effect of P and starter N on cowpea performance in Minna.

MATERIALS AND METHODS

Study site

The experiment was conducted in the screen house of School of Agriculture and Agricultural Technology, Federal University of Technology, Minna (latitude 09° 31.86' N and longitude 06° 27.28'E). Minna lies within the southern Guinea savanna zone of Nigeria. The climate of Minna is sub-humid with mean annual rainfall of 1300 mm and a distinct dry season of about 5 months duration occurring from November to March (Ojanuga, 2006). The rainy season commences most of the time in April and lasts till October, with fluctuations in amount of rainfall received per year.

Treatments and experimental design

The treatment consisted of five levels of phosphorus 0, 10, 20, 30 and 40 kg P ha⁻¹ and four levels of Nitrogen 0, 10, 20 and 30 kg N ha⁻¹. The experimental design was a 5 x 4 factorial arrangement, given a total of twenty treatment combinations laid out in completely randomized design (CRD) with four replicates. The cowpea variety used was SAMPEA 15 (IT99K - 573 - 2 - 1), a medium maturing variety. The cowpea used was sourced from Institute for Agricultural Research (IAR), Zaria.

Three seeds were sown per pot which was later thinned to one seedling per pot at two weeks after sowing (WAS). Watering of the plants was done daily. The treatments 10, 20, 30 and 40 kg P ha⁻¹ as single superphosphate and 10, 20, 30 kg N ha⁻¹ as urea was applied at 2 WAS. 10 ml of Lara force with active ingredients (Lambda-cyhalothrin 2.5% EC) dissolved in 2.5L of water was used to control insect.

Soil sampling and analysis

Surface soil (0-15 cm) sample was collected from the Teaching and Research Farm of Federal University of Technology, Minna using a shovel. The sample was air dried, gently crushed, passed through a 2 mm sieve and thoroughly mixed together. 10 kg of soil was used to fill each pot. 50 g of the soil samples was used for physical and chemical analysis some were further passed through 0.5 mm sieve to determine soil organic carbon and total nitrogen. The soil samples were analyzed using standard methods as described by Agbenin (1995). Particle size distribution was determined by Bouyocous hydrometer method. Soil pH was determined in a 1:2.5 soil to water and 0.1 M CaCl₂ using a glass electrode pH meter. The organic carbon was determined using Walkley-Black wet oxidation method. Total Nitrogen was determined by micro Kjeldahl method. Exchangeable bases (Ca²⁺, Mg²⁺, K⁺ and Na⁺) was extracted with 1N neutral ammonium acetate (NH₄OAc) solution and amounts of K⁺ and Na⁺ in solution was determined using a flame photometer while Ca²⁺ and Mg²⁺ by Na-EDTA titration. Exchangeable acidity (H⁺ and Al³⁺) was determined by titrimetric method

Data collection

Plant height was measured at 2, 4, 6, 8, and 10 WAS by measuring from the ground level of the soil to the top of the tallest leaf using a meter rule. Number of leaves was taken at 2, 4, 6, 8, and 10 WAS by counting the well-established leaves. Number of days to flowering (DF) and number of days to podding (DP) were recorded. Number of pods per plant (PP) was counted and recorded, the number of grain per pod (GP) was also counted and recorded. The pods were harvested, threshed manually, and the grain yield (GY) and haulms yield (HY) were weighed and recorded.

Statistical analysis

All the data obtained from the experiment were subjected to analysis of variance (ANOVA) using the SAS programme (SAS, 2002) with treatment means compared using Duncan Multiple Range Test (DMRT) at 5% probability level.

RESULTS AND DISCUSSION

Some physical and chemical properties of the soil prior to sowing

The results of the physical and chemical properties of the soil prior to sowing are shown in Table 1. The soil texture was sandy loam, the soil has a high proportion of sand fraction, 798 g kg⁻¹. The pH was slightly acidic which was suitable for plant growth as most plant nutrients are available for plant uptake at pH 5.5-6.5 (Brady and Weil, 2002). The organic carbon and available phosphorus were low

while total nitrogen was high (Esu, 1991). The low organic carbon content of the soil might be partly attributed to the rapid organic matter mineralization

Effect of phosphorus and starter nitrogen fertilization on plant height and number of leaves

The main effect of phosphorus and starter nitrogen on the plant heights are shown in Table 2. The application of nitrogen was not significantly ($P \geq 0.05$) different on plant height at 2, 4, 6, 8 and 10 WAS while the main effect of phosphorus on plant height was significant ($P \leq 0.05$) at 4 WAS, 6 WAS,

8 WAS, and 10 WAS. Application of phosphorus produced the tallest plant than control.

The main effects of phosphorus and starter nitrogen on the number of leaves are shown in Table 3. The effect of N on the number of leaves at 2, 4, 6, 8 and 10 WAS were not significantly ($P \geq 0.05$) different. The main effects of P on number of leaves of cowpea were significant ($P \leq 0.05$) at 4 WAS, 6 WAS, 8 WAS, and 10 WAS. Application of 20 kg P ha⁻¹ and 30 kg P ha⁻¹ produced which was significant ($P \leq 0.05$) different than 10 kg P ha⁻¹ and control at 6, 8 and 10 WAS.

Table 1: Some physical and chemical properties of the soil prior to sowing

Parameter	Value
Sand (g kg ⁻¹)	798
Silt (g kg ⁻¹)	90
Clay (g kg ⁻¹)	112
Textural class	Sandy loam
pH (H ₂ O)	6.5
Total Nitrogen (g kg ⁻¹)	0.56
Organic carbon (g kg ⁻¹)	4.22
Available phosphorus (mg kg ⁻¹)	6
Exchangeable bases (cmol kg ⁻¹)	
Calcium	4.00
Magnesium	1.00
Potassium	0.09
Sodium	0.16
Exchangeable acidity (cmol kg ⁻¹)	0.02
ECEC (cmol kg ⁻¹)	5.27

Table 2: Effect of phosphorus and starter nitrogen fertilization on plant height of cowpea

Treatment	Plant height (cm)				
	2 WAS	4 WAS	6 WAS	8 WAS	10 WAS
Phosphorus (P) (kg P ha ⁻¹)					
0	29.2 ^a	32.8 ^b	54.5 ^b	73.9 ^b	77.8 ^b
10	29.6 ^a	37.7 ^a	92.4 ^a	111.9 ^a	113.8 ^a
20	27.4 ^a	39.5 ^a	98.1 ^a	118.2 ^a	120.0 ^a
30	28.3 ^a	39.5 ^a	94.5 ^a	118.7 ^a	120.9 ^a
40	28.4 ^a	39.2 ^a	95.3 ^a	123.2 ^a	124.5 ^a
SE±	8.50	10.29	48.70	52.94	52.07
Nitrogen (N) (kg N ha ⁻¹)					
0	28.8 ^a	37.8 ^a	81.0 ^a	106.7 ^a	109.2 ^a
10	28.2 ^a	37.2 ^a	89.9 ^a	115.1 ^a	117.3 ^a
20	29.1 ^a	38.4 ^a	89.2 ^a	110.8 ^a	112.9 ^a
30	28.2 ^a	37.7 ^a	87.8 ^a	104.1 ^a	106.2 ^a
SE±	7.60	9.20	43.56	47.35	46.57
Interaction					
P*N	NS	NS	NS	NS	NS

Means with the same letter(s) in a column are not significantly different at 5% level of probability.

NS: Not significant

WAS: Weeks after sowing

Table 3: Effect of phosphorus and starter nitrogen fertilization on number of leaves of cowpea

Treatment	Number of leaves				
	2 WAS	4 WAS	6 WAS	8 WAS	10 WAS
Phosphorus (P) (kg P ha⁻¹)					
0	8 ^a	15 ^b	25 ^c	28 ^c	43 ^b
10	8 ^a	19 ^a	32 ^b	42 ^b	47 ^b
20	8 ^a	21 ^a	38 ^a	51 ^a	56 ^a
30	8 ^a	21 ^a	39 ^a	51 ^a	60 ^a
40	8 ^a	20 ^a	36 ^{ab}	48 ^a	54 ^a
SE±	1.00	11.34	14.68	23.44	25.71
Nitrogen (N) (kg N ha⁻¹)					
0	8 ^a	18 ^a	33 ^a	47 ^a	55 ^a
10	8 ^a	19 ^a	34 ^a	43 ^a	50 ^a
20	8 ^a	19 ^a	33 ^a	41 ^a	51 ^a
30	8 ^a	20 ^a	35 ^a	44 ^a	53 ^a
SE±	0.89	10.15	13.13	20.97	22.99
Interaction					
P*N	NS	NS	NS	NS	NS

Means with the same letter(s) in a column are not significantly different at 5% level of probability.

NS: Not significant

WAS: Weeks after sowing

The increased in plant height and number of leaves through the phosphorous application This result is in conformity to the results observed by Krasilnikoff *et al.*, (2003) and Nyoki *et al.* (2013). This could be attributed to the fact that phosphorus is required in large quantities in shoot and root tips where metabolism is high and cell division is rapid (Ndakidemi and Dakora, 2007). Afolabi *et al.* (2013) also observed increase in plant height of cowpea with application of phosphorus fertilizer. The interactive effects of P, and N on number of leaves were not significantly ($P \geq 0.05$) different from each other at 2, 4, 6, 8 and 10 WAS.

Effect of phosphorus and starter nitrogen fertilization on yield components

The main and interactive effect of P and N on growth parameters i.e days to flowering, days to podding, pods pot⁻¹, grains pot⁻¹, grain yield, fresh haulms yield and dry haulms yield are shown in Table 4. The main effects of P on DF, DP, PP, GP, OY, HY were significant ($P \leq 0.05$).

The main effect of P on days to flowering was significantly different among treatments with the treatment with no phosphorus having the longest number of days to flowering 46 days which was significantly different from all other treatment. The treatment with 20 kg P ha⁻¹ with number of days to flowering of 42 days was significantly different ($P \leq 0.05$) from the treatment with 0 kg P ha⁻¹ and 40 kg P ha⁻¹. The treatment with 40 kg P ha⁻¹ had the shortest number of days to flowering 39 days and is significantly different ($P \leq 0.05$) from treatment with no P and that with 10 kg P ha⁻¹ and 20 kg P ha⁻¹. Application of phosphorous fertilization reduced the number of days to podding. Application of 40 kg P ha⁻¹ podded earlier than other treatment and was significant ($P \leq 0.05$) to 0, 10, 20, and 30 kg P ha⁻¹.

The highest level of P of 40 kg P ha⁻¹ produced the highest number of pods which was significantly different ($P \leq 0.05$) from control. Application of 40 kg P ha⁻¹ produced the highest number of grains which was significantly higher than treatments with 10 kg P ha⁻¹ and control but similar to 20 kg P ha⁻¹ and 30 kg P ha⁻¹.

Table 4: Effect of phosphorus and starter nitrogen fertilization on yield components of cowpea

Treatments Phosphorus (P) (kg P ha ⁻¹)	DF	DP	PP	GP	GY(g plant ⁻¹)	HY(g plant ⁻¹)
0	46 ^a	51 ^a	3 ^b	12 ^c	1.6 ^b	2.8 ^d
10	42 ^b	47 ^b	7 ^a	33 ^a	6.3 ^a	11.6 ^c
20	41 ^b	47 ^b	7 ^a	33 ^a	6.4 ^a	19.4 ^b
30	40 ^{bc}	47 ^b	7 ^a	24 ^b	6.5 ^a	26.3 ^a
40	39 ^c	44 ^c	8 ^a	35 ^a	6.9 ^a	24.4 ^a
SE±	6.82	7.09	3.03	20.56	5.49	15.46
Nitrogen(N) (kg N ha ⁻¹)						
0	41 ^a	47 ^a	7 ^a	24 ^b	4.7 ^a	17.7 ^a
10	42 ^a	48 ^a	6 ^b	29 ^{ab}	5.6 ^a	15.2 ^a
20	42 ^a	48 ^a	5 ^b	25 ^b	4.8 ^a	17.8 ^a
30	42 ^a	46 ^a	6 ^b	31 ^a	5.5 ^a	17.0 ^a
SE±	6.10	6.35	2.71	18.39	4.91	13.83
Interaction				NS		
P*N	*	NS	NS		NS	NS

Means with the same letter(s) in a column are not significantly different at 5% level of probability using DMRT.

NS: Not significant, DF: Days to flowering, DP: Days to podding, PP: pod(s) per plant, GP: Grain of pod(s), GY: Grain yield, HY: Haulm yield, *: Significant at 5% level of probability.

The shortest number of days to flowering and podding was observed at the cowpea supplied with the highest amount of phosphorus fertilizer. Similar results was observed by Haruna and Usman, (2013) that phosphorus is an important nutrient in flower setting and grain formation. In a related study Ndor *et al.* (2012), reported that flower setting in cowpea is a function of levels of soil and applied phosphorus and genetic make-up of the variety which resulted in number of days to flowering. Early optimum P supply is essential for laying down the primordial for reproductive parts. Also, the enhancement of growth by phosphorus application induces early flowering.

The effect of phosphorus levels on days to podding was significantly different.

This could be due to higher level of phosphorus in those pots. Mokwunye and Bationo (2002) have reported that, P is essential for photosynthesis, pod development and grain filling in leguminous crops. This may also be due to the fact that the increase in phosphorus level contributed by SSP fertilizer application increases the number of pods as reported by Rani (1999) who stated that increase in phosphorus level results to an increase in the number of pod plant⁻¹. Phosphorus is responsible for nodulation in cowpea. Thus higher nodulation resulted in higher nitrogen fixation and eventually the number of pods per plant. Similar results was observed by (Ndor *et al.*, 2012; Haruna and Usman, 2013), who also discovered significant increase in pod number of cowpea in response to phosphorus application.

Application of P irrespective of the level produced significantly ($P \leq 0.05$) higher grain yield than control, while 30 kg P ha⁻¹ and 40 kg P ha⁻¹ produced the highest haulms yield than other treatments. The effect of N on number of days to flowering, number of days to podding, grain yield and haulms yield were not significantly different ($P \leq 0.05$) from each another.

The treatment 30 kg N ha⁻¹ producing the highest number of pods per plant which was significantly different from 10, 20 and 30 kg N ha⁻¹. Application of 30 kg N ha⁻¹ produced the highest number of grains per pods which was significant ($P \leq 0.05$) than treatments 20 kg N ha⁻¹ and 0 kg N ha⁻¹ but similar 10 kg N ha⁻¹.

Table 5: The interaction effect of phosphorus and starter nitrogen fertilization on days to flowering of cowpea

Treatment	Nitrogen (kg ha ⁻¹)			
	0	10	20	30
Phosphorus (kg ha ⁻¹)				
0	46 ^{ab}	43 ^{abc}	46 ^{abc}	48 ^a
10	39 ^c	42 ^{abc}	44 ^{abc}	39 ^c
20	41 ^{bc}	43 ^{abc}	42 ^{abc}	42 ^{abc}
30	42 ^{abc}	41 ^{bc}	39 ^{bc}	40 ^{bc}
40	38 ^c	39 ^c	39 ^c	40 ^{bc}
SE±			1.47	

The interactive effect of P and N on number of days to flowering was shown in Table 5. The application of 40 P and 0 kg N ha⁻¹ reduced the number of days to flower which was significantly different ($P \leq 0.05$) from 0 P and 30 kg N ha⁻¹ and 0 P and 0 kg N ha⁻¹ but similar to other treatment combinations.

The interaction for P and N on number of days to podding, number of grains per pod, grain yield and haulm yield were not significant different.

The increased in the number of grains with phosphorous application, collaborate with the finding of Olaleye *et al.*, (2011) that application of adequate levels phosphorus in the soil resulted in the highest number of grains. The application of phosphorus increased grain yield. The increased grain yield due to phosphorus addition may be attributed to increased leaf area, plant height and increased branching. Yusuf (1987) also observed optimum yield of cowpea yield when 40 kg P ha⁻¹ was applied. Similar findings was reported by other workers (Okeleye and Okelana, 2000; Nyoki *et al.*, 2013) who also discovered significant increase in yield of cowpea in response to phosphorus application. The significant response of the grain yield of cowpea to phosphorus application could be attributed to the role of phosphorus in seed formation and grain filling (Haruna, 2011). The significant response of the yield and yield characters of cowpea measured to P application could be attributed to the very low content of the soil available P.

Means in a column or row followed by the same letters are not significantly different at 5% level of probability.

Grain yield is also governed by number of factors, which have a direct or indirect impact. Among them are yield components such as number of pods per plant, number of seeds per pod and 100-seed weight over a given land area (Ayodele and Oso, 2014).

CONCLUSION

From the results of this study, phosphorus application gave the highest yield and growth response and the application of 10 kg P ha⁻¹ improved most of the parameters measured and therefore it can be considered useful for the cultivation of cowpea in a pot experiment. There was no response to nitrogen fertilization on the parameters measured.

REFERENCES

- Achuba, F. I. (2006). The effect of sublethal concentration of crude oil on the growth and metabolism of cowpea (*Vigna unguiculata*) seedlings. *The Environmentalist*. 21(1): 17-20.
- Adaji, M. J., Olufala, O. O. and Aliyu, L. (2007). Effect of intra-row spacing and stand density on the growth and yield of cowpea (*Vigna unguiculata* (L.) Walp). In: Olulaja, O. O., Omokore, D.F., Akpa, G. N. and Sanni, S. A. (eds.). Proceedings of the 41st Annual Conference of the Agricultural Society of Nigeria (ASN) held at the Institute for Agricultural Research, Samaru, Ahmadu Bello University, Zaria between 22nd and 26th October, 2007. Pp 153 – 157.
- Afolabi, S.G., Ogunesan, A.O., Uzoma, A.O., Adeboye, M.K.A., Bala, A. and Osunde, A.O. (2013). Growth, nodulation and yield response of cowpea to fertilizer inputs on Alfisols in Minna, southern Guinea savanna of Nigeria. Ojeniyi, S.O. Agbede, O.O. and Jayeoba, O.J. (eds.) Soil Science, Environmental Management and food security. Proceedings of the 37th Annual Conference of the Soil Science Society of Nigeria. pp 159-165.
- Agbenin, J.O. (1995). Laboratory manual for Soil and Plant Analysis. (Selected Method and Data Analysis). Published by Agbenin. 140pp.
- Agbogidi, O.M. (2010). Response of six cultivars of cowpea (*Vigna unguiculata* (L.) Walp.) to spent engine oil. *African Journal of Food Science and Technology*. 1(6): 139-142.
- Akinola, A. A. (1978). Influence of soil organic matter on cowpea: Response to N fertilizer. *American Journal of Agronomy*. 70 (1): 21-25.
- Asumugha, V. U. (2002). Sensory and functional properties of dry vegetable cowpea product (Akara). In: Ubbanu, C. N., Eke, O. S. and Uzomah, A. (eds.). Proceedings of the 26th Annual Conference of the Nigerian Institute of Food Science and Technology (NIFST) held at the Federal University of Technology, Owerri, Imo State between 4th and 8th of November, 2002. Pp. 66 – 67.
- Awe, O. A. (2008). Preliminary evaluation of three Asian yards long bean cowpea lines in Ibadan, Southern western Nigeria. In: Proceedings of the 42nd Annual Conference of ASN held at Ebonyi State University, Abakaliki, Nigeria between 19th and 23rd of October, 2008. Pp. 246 – 249.
- Ayodele, O.J. and Oso, A.A. (2014). Cowpea responses to phosphorus fertilizer application at Ado-Ekiti, South-West Nigeria. *Journal of Applied Sciences and Agriculture*. 9: 485-489.
- Brady, N.C. and Weil, R. (2002). The Nature and Properties of Soils. 13th edition. Singapore, Pearson Education. 976pp.
- Egho, E. O. (2009). Control of major insect pests of cowpea (*Vigna unguiculata* (L.) Walp) using conventional and non-conventional chemicals. A PhD Thesis submitted to the Department of Agronomy, Delta State University, Asaba Campus. 224p.
- FAO (2002). World Agriculture: towards 2015/2030. Summary report, Rome.
- Haruna, I. M., (2011). Dry matter partitioning and grain yield potential in sesame (*Sesamum indicum* L.) as influenced by poultry manure, nitrogen and phosphorus at Samaru, Nigeria. *Journal of Agricultural Technology*. 7: 1571-1577.
- Haruna, I.M. and Usman, A. (2013). Agronomic efficiency of cowpea varieties (*Vigna unguiculata* L. Walp) under varying phosphorus rates in Lafia, Nasarawa State, Nigeria. *Asian Journal of Crop Science*. 5: 209-215.
- Islam, S., Cowmen, R.C. and Ganer, J. O. (2006). Screening for tolerance of stress temperature during germination of twenty-five cowpea (*Vigna unguiculata* L. Walp) cultivars. *Journal of Food, Agriculture and Environment*. 4(2): 189 - 191.
- Krasilnikoff, G., Gahoonia, T., and Erik-Nelson, N., (2003). Variation in phosphorus uptake by genotypes of cowpea (*Vigna unguiculata* (L.) Walp) due to differences in root and root hair length and induced rhizosphere processes. *Plant and Soil*. 251: 83 - 91.
- Langyintuo, A.S., Lowenberg-DeBoer, J., Faye, M., Lambert, D., Ibro, G., Moussa, B., Ntougam, G. (2003). Cowpea supply and demand in West and Central Africa. *Field Crop Research*. 8: 215-231.
- Mokwunye, A. U. and Bationo, A., (2002). Meeting the phosphorus needs of soils and crops of west Africa: The role of indigenous phosphate rocks. In: Vanlauwe, B., Diels, J., Sanginga, N., and Merckx, R. (Eds). Integrated plant nutrition management in sub Saharan Africa; from concept to practise, CABI/IITA, Cornwall press, Trowbridge. pp.209-224.
- Muoneke, C. O., Ndukwe, O.M., Umana, P.E., Okpara, D.A. and Asawalam, D. O. (2012). Productivity of vegetables cowpea (*Vigna unguiculata* L. Walp) and maize (*Zea mays* L.) intercropping system as influenced by component density in a tropical zone of southeastern Nigeria. *International Journal of*

- Agricultural Research and Development*. 15 (1): 835-847.
- Ndakidemi, P.A. and Dakora, F.D. (2007). Yield components of nodulated cowpea (*Vigna unguiculata*) and maize (*Zea mays*) plants grown with exogenous phosphorus in different cropping systems. *Australian Journal of Experimental Agriculture*. 47: 583-589.
- Ndor, E., Dauda, N. Abimuku, E. Azagatu D. and Anzaku, H. (2012). Effect of phosphorus fertilizer and spacing on growth, nodulation count and yield of cowpea (*Vigna unguiculata* (L) Walp) in Southern Guinea Savanna agroecological zone, Nigeria. *Asian Journal of Agricultural Science*. 4: 254-257.
- Nyoki, D., Patrick, A., and Ndakidemi, R., (2013). Economic benefits of *Bradyrhizobium japonicum* inoculation and phosphorus supplementation in cowpea (*Vigna unguiculata* (L.) Walp) grown in northern Tanzania. *American Journal of Research communication*. 1(11): 173-189.
- Nyoki, D. and Ndakidemi, P.A. (2014). Effects of phosphorus and *Bradyrhizobium japonicum* on growth and chlorophyll content of Cowpea (*Vigna unguiculata* (L.) Walp). *American Journal of Experimental Agriculture*. 4: 1120-1136.
- Ogbo, E. M. (2009). Effects of diesel fuel contamination on seed germination of four crop plants-*Arachis hypogaea*, *Vigna unguiculata*, *Sorghum bicolor* and *Zea mays*. *African Journal of Biotechnology*. 8(20): 250-253.
- Ojanuga, A.G. (2006). Agroecological Zones of Nigeria Manual. FAO/NSPFS, Federal Ministry of Agriculture and Rural Development, Abuja, Nigeria, 124 pp.
- Okeleye, K. A. and Okelana, M. A O., (2000). Effect of phosphorus fertilizer on nodulation, growth, and yield of cowpea (*Vigna unguiculata*) varieties. *Indian Journal of Agricultural Science*. 67(1): 10-12.
- Olaleye, O., Fagbola, O., Abaidoo, R. C., and Ikeorah, N., (2011). Phosphorus response efficiency in cowpea genotypes. IITA, Ibadan. Published by Canadian Centre Of Sciences And Education. Pp.1-10.
- Rani, B.P. (1999). Response of soybean to nitrogen and phosphorus application in the black soils of Krishna-Godavari zone of Andhra Pradesh. *Annals of Agricultural Research*. 20: 367- 368
- Sankie, L., Addo-Bediako, K. O. and Ayodele, V. (2012). Susceptibility of seven cowpea (*Vigna unguiculata* L. Walp) cultivars to cowpea beetle (*Callosbruchus maculatus*). *Agricultural Science Research Journals*. 2(2): 65-69.
- SAS (2002). SAS user's guide; statistical Cary N.C. Statistical Analysis System Institute Inc.
- Shiringani, R.P. and Shimeles, H. A. (2011). Yield response and stability among cowpea genotypes at three planting dates and test environments. *African Journal of Agricultural Resources*. 6(4): 3259- 3263.
- Sultana, M. N. (2003). Effects of Rhizobium inoculum and nitrogen fertilizer on yield and nutrient quality of cowpea (*Vigna unguiculata*) forage at different stages of maturity. M. Sc. (A.H). Thesis, Dept. of Animal Nutrition, Bangladesh Agricultural University, Mymensingh.
- Yusuf, Y. (1987). Effect of Phosphorus levels on yield and yield components of Cowpea at Samaru. Cropping Scheme Meeting, Institute for Agricultural Research, Ahmadu Bello University, Zaria, Nigeria, Pp. 161.