



475
462

BIOTECHNOLOGY SOCIETY OF NIGERIA (BSN)




30TH ANNUAL INTERNATIONAL CONFERENCE

FEDERAL UNIVERSITY OF TECHNOLOGY
Minna, Niger State, Nigeria.

Book of **PROCEEDINGS**

Theme:

**BIOTECHNOLOGY AS A CHANGE AGENT
FOR NATIONAL DEVELOPMENT**



DATE: 27th - 30th August, 2017
VENUE: CPES Hall, Bosso Campus, FUT Minna, Niger State.

30TH INTERNATIONAL ANNUAL CONFERENCE OF BIOTECHNOLOGY SOCIETY OF NIGERIA (BSN) (MINNA, NIGERIA 2017)

LOCAL ORGANISING COMMITTEE (LOC)

CHAIR	Prof. S.A. Garba
CO-CHAIR	Prof. A.O. Falusi
SECRETARY	Dr. E.C. Egwim
ASST. SECRETARY	Dr. I.C.J. Omalu

CONFERENCE SECRETARIAT

Dr. E.C. Egwim
Dr. I.C.J. Omalu
Mr. S.S. Eke

EXECUTIVE COUNCIL

Prof. A.O. Falusi (Chair)
Prof. S.A. Garba
Prof. U.J.J. Ujah
Prof. E.O. Ogbadoyi
Prof. A.N. Saidu
Prof. H.O. Akanya
Dr. E.C. Egwim
Dr. I. C. J. Omalu
Dr. D Damisa
Dr. A. Y. Kabiru
Dr. I. K. Olayemi
Dr. (Mrs) H. L. Muhammad
Dr. (Mrs) F. F. Adefolalu
Dr. (Mrs) V. I. Chukwuemeka
Dr. (Mrs) A. V. Ayanwale
Dr. (Mrs) S. Daniya

SCIENTIFIC

Prof. E.O. Ogbadoyi (Chair)
Dr. E.C. Egwim
Dr. I.C.J. Omalu
Dr. M.O. Adebola
Dr. F.O. Arimoro

PUBLICITY

Dr. I. C. J. Omalu (Chair)
Dr. E. C. Egwim
Dr. D Damisa
Mr. S. S. Eke

CONTENTS ISSUE II

- THE ROLE OF NANOTECHNOLOGY IN FOOD INDUSTRIES- A REVIEW. Aigbogun, E. I., Mohammed, S. S. D., Orukotan, A. A. and Tanko, J.273
- INFECTION RATE OF SNAIL INTERMEDIATE HOST OF *Schistosoma* IN THREE WATERBODIES IN BOSSO LGA, MINNA. Ibeh, E. O., Omalu, I. C. J., Damashi, M. T., Pam, D. D., Eke, S. S., Ogunniyi, M. T. and Ocha, I. M.286
- THE ROLE OF HEALTH EDUCATION AND COMMUNITY INVOLVEMENT ON THE MANAGEMENT AND CONTROL OF MALARIA. Ramon, G. O., Abdullahi, D. A., Okpaje, J. O., Adeniyi, K. A., Ocha, I. M. and Adedeji, A. A.295
- ANTIMICROBIAL ACTIVITY OF AQUEOUS EXTRACTS OF THREE MEDICINAL PLANTS ON MULTI-DRUG RESISTANT DIARRHEAL SALMONELLAE AND SHIGELLAE BACTERIA. Aminu, S., Lawal, M. S. and Yahaya, U.307
- SPATIO-TEMPORAL VARIATION OF THE PLANKTON COMMUNITY OF TAGWAI LAKE MINNA, NIGERIA. Chukwuemeka, V. I., Auta, Y. I., Arimoro, F. O., Jonah, A. S. and Osaide, S. E. O.314
- EFFECT OF NITROGEN FERTILIZER ON GROWTH AND FRUIT YIELD OF TWO OKRA (*Abelmoschus esculentus* L. Moench) VARIETIES. Ibrahim, H., Adeboye, M. K. A., Tsado, P. A. and Badamasi, M. M.321
- MUTAGENIC EFFECTS OF GAMMA RAYS ON VEGETATIVE GROWTH OF THREE SESAME (*Sesamum indicum* L.) VARIETIES. Haruna, H. B., Falusi, O. A., Muhammad, M. L., Daudu, O. A. Y., Abubakar, A. and Muhammad, A. M.331
- SEASONAL VARIATION IN PROXIMATE COMPOSITION OF SOME SELECTED FISHES. (*Auchenoglanis occidentalis*, *Tilapia galilaea*, *Tilapia zillii*, *Alestes dentex* AND *Chrysichthys auratus*) : A CASE STUDY OF TAGWAI LAKE MINNA, NIGERIA. Chukwuemeka, V. I., Auta, Y. I., Agada, T. O., Ojutiku, R. O. and Erhabor, O. F.341
- DETERMINATION OF FREE AMINO ACIDS IN SELECTED BAMBARA GROUNDNUT (*Vigna subterranea* (L.) Verdc .) ACCESSIONS OF AFRICAN ORIGIN. Atoyebi, J. O., Osilesi, O., Abberton, M. and Adebawo, O.348
- IDENTIFICATION OF WEED HOSTS OF MAJOR LEGUME VIRUSES IN NIGER STATE, SOUTHERN GUINEA SAVANNA OF NIGERIA. Abdullahi, A. A., Salaudeen, M. T., Kolo M. G. M. and Ibrahim H.358
- EVALUATION OF BIOETHANOL PRODUCTION FROM *Gmelina aborea* SAWDUST USING *Saccharomyces cerevisiae* AND *Saccharomyces carlbergensis*. Abdusalami, M. S., Denwe, S. D., Ibrahim, J. and Yahaya, U.366

ANTICANCER POTENTIAL OF NATURAL PRODUCTS: A REVIEW. Adesina, A. D., Musa, H., Abdullahi, I. Z., Abdulsalam, B. D., Haruna, I. O. and Abubakar, M. D.	372
NUTRITIONAL COMPOSITION OF GINGER AND GARLIC FLAVOURED TIGERNUT-MILK DRINK FERMENTED BY INDIGENOUS LACTIC ACID BACTERIA DURING STORAGE AT AMBIENT AND REFRIGERATION TEMPERATURE. Maduka, N. and Ire, F. S.	390
ANTIMICROBIAL ACTIVITY OF AQUEOUS EXTRACTS OF THREE MEDICINAL PLANTS ON MULTI-DRUG RESISTANT DIARRHEAL SALMONELLAE AND SHIGELLAE BACTERIA. Lawal, M. S., Aminu, S. and Yahaya, U.	404
FUNGAL OCCURANCE IN SOME BREAD PRODUCED IN KANO STATE, NIGERIA. Abubakar, A. I. and Hamza, Z. A.	410
POTENTIALS OF YEAST ISOLATES FOR BIOSURFACTANT PRODUCTION. Oyeleke, S. B., Goro, M. A., Oyewole, O. A., Okeke, K. S., Ayisa, T. T. and Mohammed, S. S. D.	419
CORRELATION BETWEEN PHENOTYPIC AND GENOTYPIC RESISTANCE IN <i>Staphylococcus aureus</i> ISOLATED FROM COW MASTITIS. Ishaq, S. A. and Abubakar, A. I.	424
SUSTAINABILITY AND DIVERSITY IN FOREST SPECIES AND THEIR IMPROVEMENT NEEDS: THE CASE OF <i>Moringa oleifera</i> (A REVIEW). Amao, A. O. Mohammed, M. S. and Alamu, O. T.	434
PROXIMATE, MINERAL AND FATTY ACID COMPOSITIONS OF MELON (<i>Legnaria sphaerica</i>) SEED. Gado, A. A., Falusi, O. A., Muhammad, L. M., Daudu, O. A. Y., Dangana, M. C. Abejide, D. R. and Yahaya, S. A.	445
EXTRACTION AND PARTIAL PURIFICATION OF PHYTOCYSTATIN FROM <i>Calotropis procera</i> . Abdullahi, A., Umar, I. A. S., Ibrahim., E., Onyike., Kabiru, A. Y., Pala, Y. Y., and Gabi, B.	455
YOUNG TISSUES MORE PRONE TO DNA DAMAGE BY RADIATIONS FROM GSM MAST. Oluwajobi, A. O., Falusi, A. O., Dangana, M. C., Albert, C. O., Akinleye, B. A. and Daudu, O. A. Y.	463
NUTRIENT AND ANTI-NUTRIENT COMPOSITIONS OF FERMENTED AND UNFERMENTED SEED OF <i>CISSUS POPULNAE</i> FROM NIGER STATE, NIGERIA. Mathew, J. T., Dauda, B. E. N., Mann, A., Ndamitso, M. M., Fadipe, A. L. and Shaba, E. Y.	467

PROXIMATE, MINERAL AND FATTY ACID COMPOSITIONS OF MELON (*Leganaria sphaerica*) Seed

¹Gado, A. A., ²Falusi, O. A., ²Muhammad, L. M., ²Daudu, O. A. Y., ²Dangana, M. C. ²Abejide, D. R and ²Yahaya, S. A.

¹Department of Biological Sciences, Federal College of Education, Kontagora, Niger State, Nigeria.

²Department of Biological Sciences, Federal University of Technology, Minna, Niger State, Nigeria.

ABSTRACT

The chemical compositions of melon seed; *Leganaria sphaerica* were examined. Proximate, mineral and Fatty acid analysis were carried out on the shelled Cucurbit specie, Protein, fat content, ash content and moisture content respectively. The proximate compositions were determined by soxhlet extraction, AOAC, Kjeldhal and other standard procedure. Mineral contents were determined using Atomic Emission Spectrometry, Atomic Absorption Spectrophotometry and Vanado-molybdate method, while anti-nutrient determinations were carried out using standard procedures. The results obtained from proximate analysis showed: moisture content (7.61%), ash content (2.90%), crude protein (30.30%), crude lipid (43.35%), crude fiber (4.10%) and carbohydrate (11.41%). The mineral analysis revealed that potassium have the highest value (7.60mg/100g), followed by phosphorus (3.36mg/100g), zinc (0.04mg/100g) and magnesium (2.00mg/100g), calcium (2.00mg/100g) while iron was the least (0.01mg/100g). The fatty acid composition showed that alpha tocopherol has the highest 84.41, palmitic 58.10, linoleic 38.11, oleic 16.23 and stearic was the least 10.00. The results of the study revealed that *Leganaria sphaerica* seed flour is a good source of important nutrients such as fat, protein, fiber and minerals. The high contents of protein and fat make these seeds valuable dietary supplements especially for the less previlage people that cannot afford meat and diary products.

Keywords: Melon, *Leganaria sphaerica*, Proximate, Mineral, Fatty acid.

*Corresponding author: ayishatmoh@yahoo.com.

INTRODUCTION

Melon seed *Leganaria siceraria* are cucurbit crop that belong to the Cucurbitaceae family which consist of 119 genera and 825 species with fibrous and shallow root system. It is a creeping annual plant and an intercropping plant,

mostly grown assubsidiary crops interplanted with early maize and yam in some savannah belt of Nigeria, West African. Melons are major food crops with several varieties which serve as a major food source (Mabalaha et al., 2007). Cucurbit species are among the economically most important vegetable

crops worldwide and are grown in both temperate and tropical regions.

The seed kernels of the Cucurbitaceae family found in markets through out West Africa are an important source of oil used for food. Those oil-rich seeds are found in a range of genera of which the most important are *Citrullus* (watermelon), *Cucurbita* (pumpkin), *Lagenaria* (bottle gourd), *Cucumis* (melon), *Telfairia* (fluted pumpkin) and *Luffa* (sponge gourd) respectively (Schipper 2002). The seeds are obtained either in shelled or unshelled forms in West African markets and are used greatly in West African cookery. The melon seeds can be milled and used to prepare a popular egusi soup where it acts as a food thickener. It can also be fermented to produce ogiri and used as condiments to season or flavor soup (Yusuf, Adewuyi & Lasisi, 2006; Achinewu, 1987). The fruits vary much in size and seeds can be removed and roasted as an edible commodity (Soliman, EL Sawy, Fadel, Osman, & Gad, 1985), or as fried cake prepared from milled seeds (Okigbo, 1984; Odunfa, 1981). Egusi is very high in nutritional value. It is rich in protein, fat and vitamins A, B1, B2 and C. It is made up of 30 - 40 % protein, and about the same proportion of oil. The oil is cholesterol free. In terms of vitamins, it contains alpha-tocopherol, a component of vitamin E. It also contains palmitic, stearic, linoleic and oleic acids and very small amount of carbohydrate and calcium. 78 % of the fat is unsaturated fatty acid, which is protective to the heart. The alpha-tocopherol found in egusi is a component of vitamin E that helps in maintaining smooth young skin and good fertility. It also contains palmitic, stearic, linoleic and oleic acids important in protecting the heart. The egusi can also be an important supplementary baby food,

helping prevent malnutrition. Blending the seeds with water and honey produces a milky liquid that can be used as formula if breast milk is unavailable. The egusi plant is also easy to grow. It is extremely resilient to pests and diseases and because it blankets the ground as it grows, it can help suppress weeds. Because of this, farmers often intercrop egusi with other crops, including sorghum, cassava, coffee, cotton, maize, or bananas. Mature egusi melons can also remain in the field for a long time without rotting, so crop loss and waste is rare. And once the seeds are harvested, they can be a reliable year-round food source because they store well.

MATERIALS AND METHODS

Collection of melon seeds

The sample was collected from farmer's field in ilodaada, Ondo state in september 2015. Ripe fruits were cut and the seeds separated. Seeds were cleaned using filter paper to remove the pulp and air-dried at room temperature. The dried seeds were ground to flour using a grinder, the flour was then packed in a clean dry plastic containers; sealed and stored at 10°C until the time for analysis.

Mineral analysis

Determination of mineral elements was done according to the method of AOAC (1990) 2.0g of the sample was ashed in a furnace at 550°C for 18hrs and the ash dissolved in 10ml of 0.1M HCl, filtered into a 100ml volumetric flask and made up to mark with distilled H₂O. This was used to determine the mineral content by the use of Atomic Absorption Spectrophotometer (AAS) using prepared standards of the different mineral elements to be analyzed.

Proximate composition

The moisture content and the fat content were determined according to the procedures described by AOAC (2000) while the ash content, crude fibre and crude protein were estimated using procedures described by Pearson, (1979). The nitrogen was estimated based on the Kjeldhal procedure and the percentage nitrogen was converted to crude protein by multiplying by a factor of 6.25 while carbohydrate was determined by simple difference as follows: Carbohydrate = $100 - (\%Ash + \%Crude\ protein + \%Crude\ fat + \%Crude\ fibre)$. Energy value was obtained by the summation after multiplying percentage carbohydrate, protein and fat by factors of 4, 4 and 9 respectively and expressed in Kcal/100g. All analyses were carried out in triplicates. All the proximate values were reported in percentage %.

Tocopherol Analysis

Determination of tocopherols (TOC) was done according to the method of American Oil Chemists Society (2003), a solution of 250 mg of oil sample in 25 ml hexane were directly used for analysis. The HPLC analysis was conducted using a Merck-Hitachi low-pressure system fitted pump, a Merck-Hitachi F-1000 Fluorescence Spectro- photometer (detector wavelengths 295 nm and 330 nm for emission) and a D-2500 integration system. A sample in volume of 4 μ L was injected onto a normal phase silica column (HPLC mode: Shimadzu, column packing size: 150 cm x 4.6 mm). The column was eluted with a mobile phase of isopropanol : hexane (98 : 2). The flow rate was 1.0 ml per min and the absorption wavelength of 295 nm. Identification of the peaks for α

tocopherol was done using pure α tocopherol standards. The tocopherol content was expressed as ppm.

Fatty Acid Composition

Fatty acid methyl esters (FAME) were prepared as described by Joseph and Ackman (1992). FAMES were transferred into a separating funnel and 4 mL of hexane added. The contents were shaken vigorously at room temperature and left to stand. The hexane layer was collected and the aqueous layer was extracted again. The hexane fractions obtained were mixed together and washed with 3-4 portions of distilled water to remove acid present. Anhydrous sodium sulphate was added for dehydration purposes. The filtrate obtained was bubbled in nitrogen gas to concentrate it then about 0.5 mL was injected into the GC. The standard solutions were also injected and the procedure was repeated for all the samples, AOAC (2000).

RESULTS AND DISCUSSION

Proximate Analysis

The proximate composition of *Leganaria sphaerica* seed flour is shown in Table 1. The moisture content was 7.61% which is a little higher than what were reported for varieties of melon seeds; 4.78- 5.21% (Abiodun and Adeleke ,2010) and pumpkin seeds; 5.00% (Elinge *et al.* ,2012) , but lower than those reported for mango seeds; 12.50% Etong, et al. ,2013) and gardenia aqualla seeds; 49% (Dagogo *et al.*, 2011). The low moisture content in the melon will help to improve its lifespan. The ash content was 2.90% which is close to that reported for melon seed varieties; 3.35- 4.89% (Elinge *et al.* ,2012), but lower to that reported for melon seeds; 6.84-6.99% by (Bankole et al., 2005) . The high ash content in the

sample indicates the percentage of inorganic mineral elements present in melon seeds. High mineral elements in foods enhances growth and development, and also catalyses metabolic processes in human body. The crude fibre content was 4.10%, which is higher than those reported for four varieties of melon seeds, 1.66-2.16% (Abiodun and Adeleke, 2010) and *Mangifera indica* kernels, 2.22-3.95% cultivars grown in Western parts of Nigeria (Kayode *et al.*, 2011), but a little lower than those of ripe and unripe *Carica papaya* seeds, 7.85% and 7.40%; ripe and unripe *Citrus sinensis* seeds, 8.05% and 7.40% respectively (Abulude, 2000). It is believed that fibre reduces the level of cholesterol in human blood and decreases the likelihood of different cancers. The fat content was found to be 43.35% which is in agreement with that reported for four varieties of melon seeds, 40.26-45.21% (Abiodun and Adeleke, 2010) but lower

than those reported for *Citrullus vulgaris* 55.00% and *Citrullus lanatus* seeds from Southern Nigeria, 57.26% (Edidiong and Eduok, 2013) and *Colocynthis citrullus* seeds, 53.85% (Bankole *et al.*, 2005). The high value of fat in the melon seeds is the reason for it being referred to as the oils seed. Fat is very vital since it provides the body with tremendous amount of energy. The protein content is 30.30% which is comparable to those reported for *Colocynthis citrullus* seeds 28.63% (Bankole *et al.*, 2005) and *Cucurbitapepo L* seeds, 27.48% (Elinge *et al.* 2012), thus *Leganaria siceraria* seeds could provide the necessary protein requirement for the rural populace. Carbohydrate value of the sample was found to be very low, 11.41%. From this result, melon seeds cannot be considered a good source of carbohydrate compared with other sources such as cereals which contain 65-75% carbohydrate.

Table 1: Proximate composition of *Lagneria sphaerica* seed flour.

Composition	% weight
Moisture	7.61
Ash	2.90
Fat	43.35
Crude fibre	4.10
Crude protein	30.30
Carbohydrate	11.41

Mineral Composition

The mineral composition of *Leganaria sphaerica* seed flour is shown in Table 2. The concentration of calcium was found to be 2.00mg/100g, calcium is a constituent of bones and helps blood to

clot and the nerves to convey messages. The concentration of phosphorus in melon seeds was estimated as 3.36mg/100g. This value is very low in comparison with the phosphorus value of 47.68mg/100g reported for pumpkin

seeds (Elinge *et al.* 2012) but much higher than 0.87mg/100g reported for *Juglans regia* seeds. The recommended daily value for phosphorus is 1000mg. The low value of phosphorus in the sample may be connected with the fact that whole grains and vegetables are usually low in phosphorus. The copper content of 0.02mg/100g for *Leganaria shaerica* seeds in this study falls below its recommended daily allowance of 1.5-3.0mg per day for adult male and female. *Leganaria shaerica* seeds can therefore not be recommended as good source of copper. Copper helps the body to use iron and sugar properly. It is also necessary for bone growth and nerve function. Deficiency of copper may result to anaemia.

Na plays a very importance role in diet as it controls high blood pressure in the body. Studies had showed that lower sodium intake helps to reduce high blood pressure in hypertensive patients. The Na

value of 0.40 was obtained in *Leganaria shaerica* seeds in this study. The report of this investigation revealed that regular consumption of *Leganaria sphaerica* seed flour would help to prevent hypertension and might lower blood pressure in hypertensive patients. This result agrees with the finding of (Aremu et al 2006), who reported that Nigerian underutilized legumes are good sources of diets for lowering blood pressure. Therefore, from the results obtained, *Leganaria sphaerica* seeds cannot be regarded as a good source of minerals needed for the formation of bones in the body. However, the sample was richer in mineral than those reported for melon varieties (Abiodun and Adeleke ,2010) , mango seed, gmelina fruit (Edidiong and Eduok, 2013) and selected Nigerian oil seeds Onyeike and Achera (2002). The high value of some of the minerals may satisfy the nutritional needs of the consumer , F.A.O. (1968).

Table 2: Mineral Composition of Lagneria sphaerica seed.

Mineral	Composition mg/100g
Magnesium	2.00
Calcium	2.00
Sodium	0.04
Pottasium	7.60
Zinc	0.04
Iron	0.01
Phosphorus	3.36
Copper	0.02

Fatty acid composition

The result showed that *Leganaria sphaerica* seeds contains 38.11 linoleic ,16.23 oleic, 58.10 palmitic and 10.00 stearic acid .The linoleic acid content is similar to those reported for corn, cottonseed, sunflower, soya bean and sesame oils (linoleic acid is the most abundant) Fokou et al. (2009). Murkovic, et al (2004) reported 35.6-60.8% linoleic acid, 21.0-46.9% oleic acid, 9.5-14.5% palmitic acid, 3.1-7.4% stearic acid for

styrian pumpkin seed oil. The nutritional value of linoleic acid is due to its metabolism at tissue levels which is precursor to hormones like prostaglandins, which activity includes constriction of smooth vessels and lowering of blood pressure . The alpha-tocopherol is a component of vitamin E that helps in maintaining smooth young skin and good fertility. It also contains palmitic, stearic, linoleic and oleic acids important in protecting the heart.

Table 3: Fatty acid composition of *Lagneria sphaerica* seed flour.

Fatty acid	Composition%
Linoleic	38.11
Oleic	16.23
Palmitic	58.10
Stearic	10.00
Alfatocopherol	84.41

CONCLUSION

Leganaria sphaerica seeds are rich in oil, fibre and protein. The fatty acid profile, is similar to that from sesame, sunflower and soybean oils that are rich in polyunsaturated fatty acids. The seed oil can be considered as a new and valuable source of edible oil. The results of the study revealed that *Leganaria sphaerica* seed flour is a good source of important nutrients such as fat, protein and fiber. The high contents of protein and fat make these seeds valuable dietary supplements especially for the less previlage people

that cannot afford meat and diary products.

REFERENCES

- Abiodun, O. A. and Adeleke, R. O. (2010). Comparative studies of nutritional composition of four melon seeds varieties. *Pakistan Journal of Nutrition*, 9(9), 905-908.
- Abulude, F. O. (2000). Chemical composition and Nutritive values of *Carica papaya* and Citrus

- sinensis seeds. *The Journal of Technological Science*, 4, 24-27.
- Adeyeye, E. I. (2002). Determination of Chemical Composition of the Nutritionally Valuable Parts of Male and Female Common West African Fresh Water Crab (*Sudananoutes africanus*). *International Journal of Food Sciences and Nutrition*, 53,189-196.
- Aletor, V. A. and Adeogun, O. A. (1995). Nutritional and anti-nutrient components of some tropical leafy vegetables. *Food Chemistry*, 53, 375-379.
- Aliyu, A. B., Musa, A. M. and Oshaniyi, J. A. (2008). Phytochemical Analysis and Mineral Composition Analysis of Some Medicinal Plants of Northern Nigeria. *Nigerian Journal of pharmaceutical Sciences*, 7(1), 119.
- American Oil Chemists' Society. (2003). Official Methods and Recommended Practices of the American Oil Chemists' Society, Method Ce 8-89, AOCS Press, Champaign, IL, USA.
- AOAC. (2000). Official Methods of Analysis of the Association of the Analytical Chemists. 17th ed. Inc. Virginia, Washington DC, USA.
- Aremu, M. O., Olaofe, O. and Akintayo, T. E. (2006). A comparative study on the chemical and amino acid composition of some Nigerian underutilized legumes flours. *Pakistan Journal of Nutrition*, 7,381-384.
- Arinanthan, V., Mohan, V. R. and Britto, A. J. (2003). Chemical composition of Certain Tribal Pulses in South India. *International Journal of Food Sciences and Nutrition*, 3,103-107.
- Bankole, S. A., Osio, A., Joda, A. O. and Kuomehin, O. A. En. (2005). Effect of drying method on the quality and storability of *Colocynthisis citrullus*. *African Journal of Biotechnology*. 4(8), 799-803.
- Copper Supplement: www.mayoclinic.org/:/drg.20070120. Retrieved on 14th July, 2014. 46. Eleck, H. (1976). Introduction to nutrition, 3rd edition. Macmillan Company, New York. USA.
- Dagogo, S. M., Muhammad, A., Alerio, A. I., Tsafe, A. I. and Itodo, A. U. (2011). Proximate, Mineral and Anti-nutritional Composition' Gardenia aqualla Seeds. *Achieves of Applied Sciences Research*, 3(4), 485-492.
- Edidiong, A. E. and Ubong, M. E. (2013). Analysis of *Citrullus lanatus* seed oil obtained from southern Nigeria. *Elixir Organic Chemistry*, 54,12700- 12703.
- Elinge, C. M., Muhammad, A., Atiku, F. A., Itodo A. U., Peni, I. J., Sanni, O. M. and Mbongo, A. N. (2012). Proximate, Mineral and Anti-nutritional Composition of Pumpkin (*Cucurbita pepo* L.) Seeds Extracts. *International Journal of Plant Research*, 2(5),46-150.

- Etong, D. I., Ayeni, K. E., Ajayi, O. O. and Oladimeji, M. O. (2013). Physicochemical properties and Nutritional values of Melina fruit (*Gmelina arborea*) and mango (*Mangifera indica*) seed. *International Journal of Conservation Science*, 6(1), 56-62.
- FAO. (1968). Food composition table for use in Africa. Food and Agricultural organization US. Department of Health Education and welfare. Rome. 22.
- Fokou, E., Achu, M. B., Kansci, G., Ponka, R., Fots, M., Tchiegang, C. and Tchouanguép, F. M. (2009). *Pakistan Journal of Nutrition*, 8(9), 1325-1334.
- Guthrie, H. A. (1989). Introductory Nutrition, 7th edition. Time Mirror Mosby College Publishers, Boston. pp 155-159.
- Hegarty, V. (1988). Decisions in Nutrition, 5th edition. Time Mirror, Mosby London. Pp 80-132.
- Institute of Medicine, food and Nutrition Board. Dietary Reference Intake: Calcium, Phosphorus, Magnesium, Vitamin D and Fluoride. Washington DC: National Academy Press, 1997.
- Joseph, J. D. and Ackman, R. G. (1992) *Journal of Official Methods of Analysis of AOAC International*, 75, 488-506.
- Kayode, R. M., Sani, D. F., Apata, J. K., Joseph, O. A., Olorunsanya, A. A., Amnongu C. and Obalowu, M. A. (2011). Physico-chemical and anti-nutritional characterization of the kernels of some mango (*Mangifera indica*) cultivars grown in Western parts of Nigeria. *Bioresearch bulletin*. 61,1-8.
- McDonald, A., Edwards, R. A., Greenhulgh, F. D. and Morgan, C. A. (1995). *Animal Nutrition*. Prentices Hall, London. pp 101-122.
- Nieman, D. C., Butterworth, D. E. and Nieman, C. N. (1992). *Nutrition watch*, WMC Brown publishers, Dubugye, USA.
- Ogungbenle, H. N. and Atere, A. A. (2014). The chemical, fatty acid and sensory evaluation of parinari *curatellifolia* seeds. *British Biotechnology Journal*, 4(4), 379-386.
- Onyeike, E. N. and Achera, G. N. (2002). Chemical composition of selected Nigerian oil seeds and physicochemical properties of the oil extracted. *Food Chemistry*, 77,431-437.
- Payne, W. J. A. (1990). *An Introduction to Animal Husbandry in the Tropics*. Longman Publishers, Singapore. pp 92- 110.
- Rude, R. K., Magnesium, In: Coates PM, Betz JM, Blackman MR, Cragg GM, Levine M, Moss J, White JD, *Encyclopedia of Dietary Supplements*. 2nd ed, New York, NY: Informal Health Care. 2010:525-37.

Schippers, R. R. (2000). African indigenous vegetables. An overview of the cultivated species, Chathan, U. K. Natural Resources Institute ACP-EU Technical Center for Agricultural and Rural Cooperation. Pp 57-58.

Soliman, M. A., EL Sawy, A. A., Fadel, H. M., Osman, F. and Gad, A. M. (1985). Volatile components of roasted *Citrullus colocynthis* var *colocynthoides*. *Agr. Biol Chem.*, Tokyo 49, 269 - 275.

The Recommended Daily Value (DV).
www.fda.gov./:/ucm64928.htm.
Retrieved on 26th June, 2014.

Yusuf, A. A., Adewuyi, S. and A.A. Lasisi, 2006. Physico-chemical composition of leaves, meals and oils of fluted pumpkin (*Telfairia occidentalis*) and melon (*Citrillus vulgaris*). *Agricultural Journal*, 1, 32-35.