

ISSN 1927-646X (Online)

Journal of Mosquito Research

Mosquitoes, Flies Making Troubles to Humans



2017

JMR Vol 7



Publisher

Sophia Publishing Group

Edited by

Editorial Team of Journal of Mosquito Research

Email: edit@jmr.biopublisher.ca

Website: <http://jmr.biopublisher.ca>

Address:

11388 Stevenston Hwy,

PO Box 96016,

Richmond, V7A 5J5, British Columbia

Canada

Journal of Mosquito Research (ISSN 1927-646X) is an open access, peer reviewed journal published online by BioPublisher.

The journal publishes all the latest and outstanding research articles, letters and reviews in all areas of mosquito research, the range of topics including (but are not limited to) the research at molecular or protein level of mosquito, impact on the ecosystem, containing positive and negative information, natural history of mosquito, also publishing innovative research findings in the basic and applied fields of mosquito and novel techniques for improvement, as well as the significant evaluation of its related application field.



BioPublisher, operated by Sophia Publishing Group (SPG), is an international Open Access publishing platform that publishes scientific journals in the field of life science. Sophia Publishing Group (SPG), founded in British Columbia of Canada, is a multilingual publisher.

Open Access

All the articles published in Journal of Mosquito Research are Open Access, and are distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



BioPublisher uses CrossCheck service to identify academic plagiarism through the world's leading plagiarism prevention tool, iParadigms, and to protect the original authors' copyrights.

Latest Content

[Residual Effectiveness of Crude Ethanol Extracts of Mature Green Fruits of Endod *Phytolacca dodecandra* \(L' Herit\) against *An. gambiae* \(Diptera: Culicidae\) Larvae](#)

Yugi J.O., Anyango D.C., Ogot H.A.

Journal of Mosquito Research, 2017, Vol.7, No.1

[Laboratory Study on the Repellent Bioefficacy of Diethyl Phenyl Acetamide \(DEPA\) against *Aedes \(Stegomyia\) aegypti* \(Linnaeus\)](#)

Chee Dhang Chen, Othman Wan-Norafikah, Koon Weng Lau, Han Lim Lee, Chong Chin Heo, Cherng Shii Leong, Mohd Sofian-Azirun

Journal of Mosquito Research, 2017, Vol.7, No.2

[Seasonal Distribution and Micro-Climatic Factors Influencing the Abundance of the Malaria Vectors in South-East Nigeria](#)

Ebuka Kingsley Ezihe, Friday Maduka Chikezie, Chukwudi Micheal Egbuche, Edith N. Nwankwo, Angus Ejidikeme Onyido, Dennis Aribodor, Samdi Musa Lazarus

Journal of Mosquito Research, 2017, Vol.7, No.3

[An Observation on Virulence of Mosquito-killing Fungus, *Pythium guiyangense* after Rejuvenated through Its Host, *Culex quinquefasciatus* Larvae](#)

Xiaoqing Su, Xiao Yang

Journal of Mosquito Research, 2017, Vol.7, No.4

[Insecticide Resistance in *Anopheles gambiae* s.l Mosquitoes in Awka, Anambra State, Southeast Nigeria](#)

Nwankwo E.N., Okorie P.N., Acha C.T., Okonkwo O.E., Nwangwu U.C., Ezihe Ebuka K.

Journal of Mosquito Research, 2017, Vol.7, No.5

[Mosquito Species Associated with Refuse Dumps within Enugu Municipal, Enugu State, Nigeria](#)

Ezihe Ebuka Kingsley, Egbuche Chukwudi Micheal, Uka Job, Chikezie Friday Maduka, Anumba Joseph Uche, Nwankwo Edith N., Chukwuzoba Obinna Anthony, Umenzekwe Chukwudi Christian, Ejehu Zika Uchenna, Inya Agha Stella Ifeoma

Journal of Mosquito Research, 2017, Vol.7, No.6

[Some Determinant Factors of Malaria Prevalence in Nigeria](#)

Sunday Etim Bassey, Sylvester Chibueze Izah

Journal of Mosquito Research, 2017, Vol.7, No.7

[The Larvicidal and Repellent Effect of Bush Mint Plant \(*Hyptis suaveolens*\) and Hairy Sicklepod \(*Senna hirsuta*\) Extracts against the Larvae and Adult of *Culex quinquefasciatus*](#)

Ebuka Kingsley Ezihe, Chukwudi Micheal Egbuche, Edith N. Nwankwo, Angus Ejidikeme Onyido, Inya-Agha Mishael Uzochukwu, Anumba Joseph U., Chukwuzoba Obinna Anthony, Onyali Ikechukwu O.

Journal of Mosquito Research, 2017, Vol.7, No.8

[Influence of Meteorological Variables on Diversity and Abundance of Mosquito Vectors in Two Livestock Farms in Ibadan, Nigeria: Public Health Implications](#)

Opayele Adewale Victor, Adeniji Johnson Adekunle, Ibrahim Kolade Tahiru, Olaleye Olufemi David

Journal of Mosquito Research, 2017, Vol.7, No.9

[Levels of Insecticide Resistance to Temephos, and Associated Mechanisms in *Culex pipiens* Mosquitoes from Central Tunisia](#)

Jaber Daaboub, Ahmed Tabbabi, Ali Lamari, Mohamed Feriani, Chokri Boubaker, Hassen Ben Cheikh

Journal of Mosquito Research, 2017, Vol.7, No.10

[Status and Genes Involved in Insecticide Resistance in *Anopheles gambiae* Sibling Species in Lomé \(Togo\), West Africa](#)

Accrobessy K., OléL.M., Dorkenoo A.M., Oss èR., Akinro B., Sidick A., Akogbeto M.C., Glitho A.

Journal of Mosquito Research, 2017, Vol.7, No.11

[Larvicidal Efficacy of *Cleistanthus patulus* Muell. Arg. \(Euphorbiaceae\) Leaf Extract against Filarial Vector *Culex quinquefasciatus* \(Say 1823\)](#)

Jibon Kumar Pal, Aniket Singh, Goutam Chandra

Journal of Mosquito Research, 2017, Vol.7, No.12

[Pupicidal Activity of Ethanol and Water Extracts of *Phytolacca dodecandra* \(L' Herit\) on *Anopheles gambiae* \(Diptera: Culicidae\) Pupae](#)

Yugi J.O., Kiplimo J.J., Misire C.

Journal of Mosquito Research, 2017, Vol.7, No.13

[Mosquito Larvicidal Activity of Fractionated Methanolic Extract of *Rhizophora mucronata* Stilt Root and *Sesuvium portulacastrum* against Dengue Larvae](#)

P. Desiyamani, M. Syed Ali, V. Anuradha, N. Yogananth, J. Chitra

Journal of Mosquito Research, 2017, Vol.7, No.14

[Larvicidal Activities of Leaf Extracts of *Adansonia digitata* L. \(Malvales: Malvaceae\) and *Ficus sur* Forssk \(Rosales: Moraceae\) against *Culex quinquefasciatus* Mosquito \(Diptera: Culicidae\)](#)

Olayemi Israel Kayode, Samuel Oyiza Mariah, Ukubuiwe Azubuike Christian, Ande Adeolu Taiwo, Adeniyi Kamoru Abdulazeez, Shittu Kudirat Oluwatosin

Journal of Mosquito Research, 2017, Vol.7, No.15

[High Level of DDT Resistance and Reduced Susceptibility to Deltamethrin in *Anopheles gambiae*, *Anopheles coluzzi*, and *Anopheles arabiensis* from Urban Communities in Oyo State, South-West Nigeria](#)

Adedapo Olufemi Adeogun, Kehinde Olajide Popoola, Adedayo Olatunbosun Oduola, Abiodun Kanmi Olakiigbe, Taiwo Samson Awolola

Journal of Mosquito Research, 2017, Vol.7, No.16

[Influence of Agro-chemical Inputs on Suitability of Physicochemical Conditions of Rice-Fields for Mosquito Breeding in Minna, Nigeria](#)

Salihu I.M., Olayemi I.K., Ukubuiwe A.C., Garba Y., Gusau A.Y., Nma-Etsu M., Usman M.D.

Journal of Mosquito Research, 2017, Vol.7, No.17

Research Article

Open Access

Influence of Agro-chemical Inputs on Suitability of Physicochemical Conditions of Rice-Fields for Mosquito Breeding in Minna, Nigeria

Salihu I.M.¹ ✉, Olayemi I.K.², Ukubuiwe A.C.², Garba Y.³, Gusau A.Y.¹, Nma-Etsu M.¹, Usman M.D.¹¹ Department of Biological Sciences, Federal Polytechnic Bida, Niger State, Nigeria² Department of Biological Sciences, Federal University of Technology, Minna, Nigeria³ Department of Biology, Federal College of Education, Kontagora, Niger State, Nigeria✉ Corresponding author email: ismaikudi@gmail.comJournal of Mosquito Research, 2017, Vol.7, No.17 doi: [10.5376/jmr.2017.07.0017](https://doi.org/10.5376/jmr.2017.07.0017)

Received: 15 Aug., 2017

Accepted: 08 Sep., 2017

Published: 15 Sep., 2017

Copyright © 2017 Salihu et al., This is an open access article published under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Preferred citation for this article:

Salihu I.M., Olayemi I.K., Ukubuiwe A.C., Garba Y., Gusau A.Y., Nma-Etsu M., and Usman M.D., 2017, Influence of Agro-chemical inputs on suitability of physicochemical conditions of Rice-Fields for mosquito breeding in Minna, Nigeria, Journal of Mosquito Research, 7(17): 134-141 (doi: [10.5376/jmr.2017.07.0017](https://doi.org/10.5376/jmr.2017.07.0017))

Abstract The influence of chemicals used in rice farming in Minna, on physicochemical properties of rice-field mosquito larval habitats was investigated during the rainy season of 2013. Standard water quality analyses techniques were followed in determining the concentrations of the different physicochemical parameters. The results showed that Temperature, Dissolved Oxygen (DO) and Biological Oxygen Demand (BOD) did not vary significantly among the types of rice fields (range = 30.00±0.00 to 30.50±0.11°C, 6.00±0.00 to 7.00±1.41 mg/l, and 4.00±0.00 to 4.50±2.12 mg/l, respectively). Turbidity and Nitrates were significantly ($p<0.05$) highest in rice fields without chemical inputs (site A), with values ranging from 0.27±0.21 to 0.24±0.18 NTU and 9.92±1.94 to 7.72±2.23 mg/l, respectively. However, these two parameters were not significantly different ($p>0.05$) between site B (rice field with chemical fertilizer only) and site C (rice field with chemical fertilizer and herbicide). Levels of Alkalinity, Hardness, Sodium and Conductivity were highest in site B and least in Site C, with values ranging from 31.00±15.56 to 132.00±19.79 mg/l, 32.00±4.14 to 81.00±41.01 mg/l, 38.15±2.75 to 51.70±1.41 mg/l and 194.00±43.84 to 508.50±44.5 µs/cm, respectively. While the concentration of Potassium was significantly ($p<0.05$) lowest in site A (10.40±0.14 mg/l), the concentration of Chloride was significantly lowest in site C (23.41±15.03 mg/l) and highest in site A (31.45±4.03 mg/l). Significant positive and negative correlations between some of the parameters were also recorded. These results suggest differential suitability of rice field physicochemical condition for mosquito breeding in Minna and, thus, should provide baseline guide for mosquito vector control in relation to sustainable rice farming in the area.

Keywords Rice fields; Agricultural chemical inputs; Mosquito larval habitat

Background

Nigeria is the most populous country in Africa, with a population of over 130 million people. Its domestic economy is dominated by agriculture, which accounts for about 40% of the Gross Domestic Products (GDP) and two-third of the labour force (Akande, 2000). Generally, cereal crops including sorghum, maize, millet, rice and wheat, etc. are considered staple and of prime importance in ensuring food security and major source of income for households in Nigeria and of these cereals crops, rice is considered the most acceptable, resulting in about 10% increase in annual production, dictated by consumer preference. The Nigerian Government spends as much as \$300 million on rice importation annually (Akande, 2000).

Rice production and consumption in Nigeria cut across cultural, religious, ethnic or geographical boundaries. It is available in Five Star Hotels in the big cities and towns as well as in the 'most local' of eating places the remotest villages throughout the country. It is highly priced and widely accepted for festivities (Samson and Kadiri, 2007). Niger state is one of the major rice producing states in Nigeria (Saidu, 2008). The state has the highest average yield (on average 3 tons (t) of paddy) from a relatively limited area (on average less than two half) (Saidu, 2008).

Though, the use of agro-chemicals such as herbicides, insecticides and fertilizers, especially in rice farming is known to increase yield per unit area and time (Renato et al., 2011), such activities have widely resulted in the pollution of aquatic ecosystem, with particular negative effects on water quality. Moreby and Southway (1999),

reported loss of diversity resulting from the de-structuring of physico-chemical conditions and alteration of natural dynamics of biological communities in aquatic ecosystems. Rice agro-ecosystems provide ideal breeding habitats for mosquito species. For example, rice irrigation farming is associated with the production of vectors that transmit pathogens to humans, including those responsible for malaria (Muturi et al., 2006).

Large scale environmental modification concomitant with processes such as forest clearing, irrigation development, human settlement and rice cultivation, inevitably result in changes in surface water quality and affect the survival of mosquito species breeding in surface water habitats (Yusuf and Olayemi, 2015). The physicochemical factors in Minna that influence oviposition, survival and spatio-temporal distribution of mosquito species in rice fields, include salts, dissolve organic and inorganic matter, degree of eutrophication, turbidity, presence of suspended mud, vegetation, temperature, light or shade and hydrogen ion concentration, which also may be affected by agrochemical inputs used in rice farming (Yusuf and Olayemi, 2015).

According to Lacey and Lacey (1990), Rice cultivation has traditionally been associated with vector-borne diseases, especially malaria and Japanese encephalitis. The mosquito vectors of these diseases lay eggs in standing water and the larvae need about 7-10 days in an aquatic environment to complete development to adults. The intermittent drying of rice fields was, therefore, tested for its mosquito control potential as far back as the early twentieth century (Takken et al., 1990). With the introduction of DDT after World War II, water management and other environmental measures to control mosquitoes were neglected. It is only since the 1980s, after the failure of the DDT-based eradication campaigns of malaria that environmental control measures are receiving renewed attention (Van der et al., 2001).

Olayemi et al. (2014) have documented the presence of high densities and diversity of vector mosquito species in Minna, the capital city of Niger state, and reported that drainages and rice fields were the most productive mosquito larval habitats in the city. Rice is widely grown in Minna, and is one of the most suitable habitats for mosquito larva productivity (Olayemi et al., 2014). The impacts of agrochemical inputs on the productivity of rice field mosquito larval habitats in Nigeria, generally, and Niger state, in particular, have not been well investigated, thus making sustainable cost effective rice production a problem in the country.

1 Materials and Methods

1.1 Study area

The study was carried out in Minna, the capital of Niger State, Nigeria, located within longitude $6^{\circ} 33'E$ and latitude $9^{\circ} 37'N$, on a land area of 88 km^2 and having a population of about 1.2 million inhabitants. Minna has a typical tropical climate with mean annual temperature of $30.2^{\circ}C$, relative humidity of 61% and rainfall of 1334 mm. The climate in the area has two distinct seasons: rainy season (April-October) and a dry season (November-March), completely devoid of rains. Its vegetation is typically grass-dominated savannah with scattered short trees (Olayemi et al., 2009). Basically, four types of wetland rice ecosystems are present in Minna, which include irrigated, rain-fed low upland and wetland rice ecosystems, with the irrigated rice environment been the dominant (Akintayo, 2011). Farming is a major pre-occupation of the inhabitants of Minna and environs, with the area particularly famous for rice production; being a staple food in the area.

1.2 Selection of rice field mosquito larval habitats

Questionnaire were administered to farmers to determine the types and combinations of agricultural chemical inputs, i.e., chemical fertilizer, herbicides or both, commonly used in rice farming in Minna. Thereafter, guided by the results of the questionnaire test, three types of rice farming chemical inputs sites in two rice field categories, rain-fed lowland and irrigated ecosystems were selected each with three replicates. The three types of sites were named: type A (rice fields with no chemical input; serving as Control), type B (rice fields with only fertilizer application), type C (rice fields with both fertilizer and herbicide application). The sites were widely located at Fadukpe, Tunga and Chanchaga area of the metropolis (Figure 1).




Figure 1 Minna street guide map showing the study area (Source: Urban and Regional Planning Department, FUT, Minna, 2012)

1.3 Collection and analyses of water samples for physicochemical properties of rice fields

Water samples were collected from each selected rice field between the hours of 08:00 and 09:00 am, at a depth of 5 cm, on each sampling day. 500 ml of water sample was collected per rice field using sample bottles of 500 ml capacity. Temperature ($^{\circ}\text{C}$) and Hydrogen ion concentration (pH) were measured *in situ* at the sites, using mercury thermometer and pH meter, respectively. The other parameters namely Turbidity, Conductivity, Potassium, Sodium, Chloride, Nitrate, Phosphate, Alkalinity, Hardness, and Dissolved Oxygen (DO) were determined in the laboratory within 2-3 hours of collection, while Biochemical Oxygen Demand (BOD) was determined on day five after collection, using the standard methods described by American Public Health Association (1992).

1.4 Statistical analysis

Data collected from all the sites and sampling periods were processed as mean \pm S.D, using EXCEL programme. Mean values of the variables per treatment were compared for significance of statistical differences using appropriate tools of ANOVA and Duncan Multiple Range Test at P = 0.05 level of significance.

2 Results

2.1 Physicochemical condition of the rice field mosquito larval habitats

The indices of physicochemical conditions of the rice field mosquito larval habitats in Minna are presented in Table 1. Temperature, Dissolved Oxygen (DO) and Biological Oxygen Demand (BOD) did not varies significantly

($P > 0.05$) among the rice fields subjected to different agricultural chemical inputs. The mean temperatures however, ranged from 30.00 ± 0.00 to $30.50 \pm 0.00^\circ\text{C}$, while DO and BOD were 6.00 ± 0.00 to 7.00 ± 1.41 mg/l and 4.00 ± 0.00 to 4.50 ± 0.71 mg/l, respectively. On the other hand, Turbidity, Nitrate and Chloride were significantly ($P < 0.05$) higher in site A (rice-fields without agrochemical inputs), where they ranged as follows: 0.27 ± 0.24 to 0.24 ± 0.18 NTU, 9.92 ± 1.94 to 7.72 ± 2.23 mg/l and 31.45 ± 15.00 to 23.41 ± 15.00 mg/l, respectively. The distribution of the concentrations of chloride among the sites followed the same trend, been significantly highest in site A (mean = 31.45 ± 03 mg/l). Alkalinity, Hardness, Sodium and Conductivity were highest in site B (rice field with only chemical fertilizer application) (means = 132.00 ± 19.79 mg/l, 81.00 ± 41.01 mg/l, 51.70 ± 1.41 mg/l and 508.58 ± 44.54 $\mu\text{S/cm}$, respectively) and least in site C (rice field with both fertilizer and herbicide) (means = 31.00 ± 15.56 mg/l, 32.00 ± 14.14 mg/l, 38.15 ± 2.75 mg/l and 194.00 ± 43.84 $\mu\text{S/cm}$, respectively). The concentration of Potassium was significantly lowest in site A (10.40 ± 0.14 mg/l) and did not significantly differ between sites B and C (14.40 ± 0.14 mg/l and 13.40 ± 3.11 mg/l, respectively).

Table 1 Variation in physicochemical properties of rice-field mosquito larval habitats subjected to various agricultural chemical inputs in Minna, Nigeria

PHYSICO-CHEMICAL PARAMETERS	Site A	Site B	Site C
Temperature ($^\circ\text{C}$)	30.50 ± 0.71^a	30.00 ± 0.00^a	30.00 ± 1.41^a
pH	6.57 ± 0.51^a	7.17 ± 1.48^b	6.16 ± 0.11^a
Turbidity (NTU)	0.27 ± 0.35^b	0.24 ± 0.28^a	0.24 ± 0.21^a
Conductivity ($\mu\text{S/cm}$)	344.50 ± 61.51^b	508.50 ± 44.54^c	194.00 ± 43.84^a
Alkalinity (mg/l)	75.00 ± 21.21^b	132.00 ± 19.79^c	31.00 ± 15.56^a
Hardness (mg/l)	62.00 ± 19.79^b	81.00 ± 41.01^c	32.00 ± 14.14^a
Potassium (mg/l)	10.40 ± 0.14^a	14.90 ± 2.12^b	13.40 ± 3.11^b
Sodium (mg/l)	47.75 ± 2.33^b	51.70 ± 1.41^c	38.15 ± 2.75^a
Phosphate (mg/l)	5.58 ± 2.69^b	2.71 ± 1.77^a	9.98 ± 1.69^c
Nitrate (mg/l)	9.92 ± 1.94^b	7.93 ± 3.18^a	7.72 ± 2.23^a
Chloride (mg/l)	31.45 ± 4.03^c	27.47 ± 0.49^b	23.41 ± 15.03^a
D.O (mg/l)	7.00 ± 1.41^a	6.00 ± 0.00^a	6.00 ± 2.83^a
B.O.D (mg/l)	4.50 ± 0.71^a	4.00 ± 0.00^a	4.50 ± 2.12^a

Note: * Mean values followed by the same superscript alphabets in a row are not significantly different at $P = 0.05$; Site A: rice-field without agrochemical application; Site B: rice field with only chemical fertilizer application and Site C: rice-field with both fertilizer and herbicide application; * Dissolved Oxygen (DO) and Biological Oxygen Demand (BOD)

2.2 Relationships within the physicochemical conditions

Table 2 shows the relationships that exist between the physicochemical parameters, in which some were significantly correlated while others were not. Significant positive correlation among the physicochemical parameters include those between Chloride and Temperature ($r=0.95$), Conductivity and pH ($r=0.96$), Alkalinity and pH ($r=0.93$), Sodium and pH ($r=0.89$), Alkalinity and Conductivity ($r=0.96$), Sodium and Alkalinity ($r=0.88$), DO and Nitrates ($r=0.92$), DO and Chloride ($r=0.76$), DO and BOD ($r=0.94$). On the other hand significant negative correlation also exist among the following parameters Phosphate and pH ($r= -0.92$), Phosphate and Conductivity ($r= -0.97$), Phosphate and Alkalinity ($r= -0.97$), phosphate and Sodium ($r= -0.91$). Potassium distinctly showed a negative correlation between sodium, Phosphate, Chloride, DO and BOD.

3 Discussions

The results of this study showed that water temperature, Dissolved Oxygen (DO) and Biochemical Oxygen Demand (BOD) were not significantly affected by agrochemical inputs in the rice-field larval habitats. The temperatures recorded in this study were considerably lower than the 40.00°C maximum limit for the thriving of aquatic organisms by Federal Environmental Protection Agency (FEPA, 1990). This may be due to the fact that water temperature is generally dictated by atmospheric temperature, as influenced by the amount of absorbed solar radiation (Kemker, 2014). The conductive water temperature in Minna rice-fields may mean that such habitats are active vector-mosquito production sites, with its attendant public health implications in the area. According to Kawo et al. (2004) low water temperatures eliminate thermal pollution in aquatic habitats and promote the thriving of invertebrates.

Table 2 Cross correlation between the physicochemical parameters of rice field's mosquito larval habitats subjected to various agricultural chemical inputs in Minna, Nigeria

Physico-chemical Parameters	Temperature	pH	Turbidity	Conductivity	Alkalinity	Hardness	K	Na	Phosphate	Nitrate	Chloride	D.O	B.O.D
Temperature	1.00												
pH	0.23	1.00											
Turbidity	0.73*	0.38	1.00										
Conductivity	0.18	0.96*	0.30	1.00									
Alkalinity	-0.06	0.93*	0.20	0.96*	1.00								
Hardness	0.32	0.69	0.07	0.69	0.59	1.00							
Potassium	-0.70	0.19	-0.74*	0.14	0.30	0.21	1.00						
Sodium	0.31	0.89*	0.37	0.96*	0.88*	0.77*	-0.04	1.00					
Phosphate	0.02	-0.92	-0.07	-0.97*	-0.97*	-0.70*	-0.34	-0.91*	1.00				
Nitrate	0.50	-0.20	0.21	0.01	-0.20	0.12	-0.71*	0.22	0.08	1.00			
Chloride	0.95*	0.35	0.69	0.38	0.12	0.42	-0.70*	0.51	-0.18	0.64	1.00		
D.O	0.65	-0.14	0.23	0.03	-0.23	0.08	-0.68	0.17	0.09	0.92*	0.76*	1.00	
B.O.D	0.69	-0.25	0.19	-0.15	-0.41	-0.04	-0.62	-0.05	0.27	0.76*	0.72*	0.94*	1.00

Note: * = significantly correlate

The DO recorded in the rice-fields in this study (range = 6.00 – 7.00 mg/l) were higher than those (i.e., 4.42 – 4.70 mg/l) reported in a reservoir elsewhere in the state (Ibrahim et al., 2009). The higher DO in this study may be due to high density of phytoplankton that characterize rice-field ecosystem generally (Alshami et al., 2014). Phytoplanktons are the principal producers of Oxygen, as by-product of photosynthesis, in aquatic habitats (Alshami et al., 2014).

The BOD recorded in this study is lower than reported by Sangpal et al. (2011) from India, who attributed the high BOD in the aquatic habitat to excessive discharge of sediment and sewage into the water body. This may mean that the rice-fields in Minna are not organically or chemically polluted.

The concentrations of turbidity, Nitrate and Chloride were significantly higher in rice-fields where there were no applications of agrochemicals (i.e., Site A). The higher turbidity in this site may be due to higher run-off discharge from the surrounding land area. The magnitude of run-off into aquatic habitats is often influenced by anthropogenic activities in the surrounding area (Leroy et al., 2002), and this may vary considerably in different rice-field mosquito larval habitats. The higher Nitrate content of site A may also be due to higher deposition of sewage, the principal source of Nitrate in aquatic ecosystems (Mateo-Sagasta and Jacob, 2012). High Nitrate content may result in algal bloom in aquatic habitats (Glendon et al., 1999) and, hence, increased mosquito productivity, as the larvae will have access to abundant food supply (Alan et al., 2003).

The concentrations of Alkalinity, Hardness, Sodium and Conductivity were significantly higher in rice-fields with fertilizer application. This may be an indication of physico-chemical parameters singly affected by chemical fertilizers in aquatic habitats. Alkalinity and Conductivity, for example, are known to influence the survival and productivity of aquatic invertebrates (William, 2000). Therefore, this may mean that rice-fields subjected to only chemical fertilizer application in Minna produce mosquitoes at differential rates from the other counterpart habitats subjected to other kind of agrochemical practices. The level of Alkalinity, water hardness and Chloride recorded in the rice-fields in this study fell within ranges optimum for the productivity of aquatic organisms including mosquitoes (Ramatake and Morghe, 1998; Idowu et al., 2004; Girgin et al., 2010). The concentration of Potassium was significantly higher in sites B and C, where there were applications of inorganic fertilizer. This should be expected, as Potassium is one of the three major components of inorganic fertilizers (i.e., NPK used) in the study area. However, this may mean that rice farmers in Minna wrongly use the NPK fertilizer for their crops instead of the Urea (Nitrogen-based) fertilizer, recommended for optimum rice productivity. This assertion is buttressed by the facts that Nitrate concentration was even higher in rice-fields where there was neither fertilizer nor herbicide application.

4 Conclusions

Agro-chemical input practices in rice farming in Minna though may enhance yield, significantly alter critical physico-chemical parameters that condition the development and thriving of mosquitoes in the larval habitats. The little or no significant effects of the agro-chemical on Temperature, Dissolved Oxygen and Biological Oxygen Demand may not promote mosquito production but this gain is effectively countered by the enhancement of Alkalinity, Sodium, Potassium and Conductivity which are critical in mosquito immature development. The strong correlations between vital physico-chemical parameters indicate that such factors act collectively or perhaps, in-synergy on mosquito development. This probable relationships between mosquito immature development and collegiate of physico-chemical parameters requires urgent elucidation. The findings of this study should provide baseline information for cost-effective management of mosquito productivity in rice wetlands in Minna and, by extension, in Nigeria as a whole.

Author's contributions

OIK and SIM designed the experiment; SIM and GY analysed the data; SIM, GAY and UMD wrote the draft manuscript; OIK, SIM and UAC contributed to writing of the main manuscript; All authors agree with manuscript results and conclusion; finally the articulated manuscript were read and reviewed together.

Acknowledgement

We wish to appreciate the technologists of the Departments of Biological Sciences and Water, Aquaculture and Fisheries Technology (WAFTE), Federal University of Technology, Minna for analyses of the water analyses.

References

- Akande T., 2000, An Overview of the Nigeria Rice Economy, Nigeria Institute of Social and Economic Research, Ibadan, pp.1-11
- Akintayo O.I., Rahji M.A.Y., Awoyemi T.T., and Adeoti A.I., 2011, Determinants of Yield Gap in Lowland Rice Production in North Central Nigeria, *Agrosearch*, 1(2): 1-10
- Alan R.T., Robert W.H., Fakhri A.B., Mary S.B., Cory C.C., Sharon K.C., and Amir H.W., 2003, Human health effects of a changing global nitrogen cycle, *Front Ecology and Environ*, 1(5): 240-246
[https://doi.org/10.1890/1540-9295\(2003\)001\[0240:HHEOAC\]2.0.CO;2](https://doi.org/10.1890/1540-9295(2003)001[0240:HHEOAC]2.0.CO;2)
- Al-shami S.A., Che-Salma M.R., Siti-Aziza M.N., and Abu-Hassan A., 2010, The influence of routine agricultural activities on the quality of water in tropical rice field ecosystem, *Applied Ecology and Environmental Research*, 8 (1): 11-18
https://doi.org/10.15666/aeer/0801_011018
- American Public Health Association, 1992, *Standard Methods of Examination of Water and Waste Water*, 17th Edition, American Public Health Association Inc., United State
- Federal Environmental Protection Agency, 1990, *Limits for waste effluent discharge in to surface waters*, pp. 2-6
- Girgin S., Kazanci N., and Dögel M., 2010, Relationship between aquatic insects and heavy metals in an urban stream using multivariate techniques, *International Journal of Environmental Science Technology*, 7 (4): 653-664
<https://doi.org/10.1007/BF03326175>
- Glendon R., David P.M., and Corinne G., 1999, Eutrophication and Algal Blooms, *Environmental and Ecological Chemistry*, 2: 1-10
- Ibrahim B.U., Auta J., and Balogun J.K., 2009, An assessment of the physicochemical parameters of Kontagora Reservoir, Niger State, Nigeria, *Bayero Journal of Pure and Applied Sciences*, 2(1): 64-69
- Idowu R.T., Inyang N.M., and Eyo J.E., 2004, The physicochemical parameters of an African Arid Zone Manmade Lake, *Animal Research International*, 1 (2): 113-119
- Kawo A.H., Mohammed H.I.D., and Ajayi F.M., 2004, Microbiological and physicochemical qualities of Bayero University stabilization ponds effluent, Kano Nigeria, *Best Journal of Biological Sciences and Environmental science for the Tropics*, 1(1): 16-19
- Kemker C., 2014, *Water Temperature Fundamentals of Environmental Measurements*, Fondriest Environmental
- Lacey L.A., and Lacey C.M., 1990, The medical importance of rice-land mosquitoes and their control using alternatives to chemical insecticides, *Journal of the American Mosquito Control Association*, 6 (2): 1-93
- Leroy N.P., Mark M.B., and John W.D., 2002, *Aquatic Ecosystems and Global Climate Change: Potential Impacts on Inland Freshwater and Coastal Wetland Ecosystems in the United States*, Center for climatic and energy solution, pp.1-2
- Mateo-sagasta X., and Jacob B., 2012, *Water pollution from agriculture: A review*, Hugh Turrall Independent Consultant, Federation of Agricultural organization, 7-11
- Moreby S.J., and Southway S.E., 1999, Influence of autumn applied herbicides on summer and autumn food available to birds in winter wheat fields in southern England, *Agriculture, Ecosystems and Environment*, 72: 285-297
[https://doi.org/10.1016/S0167-8809\(99\)00007-9](https://doi.org/10.1016/S0167-8809(99)00007-9)
- Muturi E., Shilulu J., Jacob B., Gu W., Githure J., and Novak R., 2006, **Mosquito species diversity and abundance in relation to land use in a rice-land agro-ecosystem in Mwea, Kenya**, *Journal of Vector Ecology*, 31: 129-137
[https://doi.org/10.3376/1081-1710\(2006\)31\[129:MSDAAI\]2.0.CO;2](https://doi.org/10.3376/1081-1710(2006)31[129:MSDAAI]2.0.CO;2)
- Olayemi I.K., Ande A.T., Isah B., and Idris A.R., 2009, Epidemiology of malaria in relation to climatic variables in Minna, Nigeria, *African International Journal of Medical Sciences*, 2: 5-10
- Olayemi I.K., Ukubuiwe A.C., and Oyibo-usman L.A., 2014, Mosquito species occurrence and diversity in conventional larval breeding site in Minna Metropolis Nigeria, *International Journal of Innovation and Scientific Research*, 9 (1): 86-93
- Ramateke D.S., and Moghe C.A., 1988, *Manual on water and waste-water analysis*, Nagpur, National Environmental Engineering Research Institute (NEERI), pp. 43
- Renato Z., Martha. B., Adaime S.C., Peixoto C.A., Friggi O.D., Prestes L.O., and Ednei G.P., 2011, *Mechanisms and Mode of Action of Herbicides Persistence in Rice Paddy Water*, InTech Publisher, Southern Brazil, pp. 20-26
- Saidu M., 2008, *Economic Analysis of Rice Processing among Women in Paikoro Local Government Area of Niger State*, B.Tech project submitted to the Department of Agricultural Economics and extension Technology, Federal University of Technology Minna, pp 2-4
- Samson I.O., and Kadiri M.A., 2007, Problems and Prospect of Rice Production in Central District of Edo State, Nigeria, *Journal of Human Ecology*, 22(2): 123-128
- Sangpal R.R., Kulkarni U.D., and Nandurkar Y.M., 2011, An assessment of the physicochemical properties of Ujjani Reservoir, Solapur District, India, *Journal of Agriculture and Biological Science*, 6 (3): 34-36
- Takken W., Snellen W.B., Verhave J.P., Knols B.G.J., and Atmosoedjono S., 1990, Environmental measures for malaria control in Indonesia, *An historical review on species sanitation*, Wageningen Agricultural University Papers, 90-97

- Van-Der H.W., Sakthivadivel R., Renshaw M., Silver J.B., Birley M.H., and Konradsen F., 2001, Alternate Wet/Dry Irrigation in Rice Cultivation: A Practical Way to Save Water and Control Malaria and Japanese Encephalitis, International Water Management Institute, Research Report 47, Colombo, Sri Lanka
- William A., 2000, Organic fertilization in culture ponds, *World Aquaculture*, 35(2): 64-65 www.ca.uky.edu/wkrec/wurtspage.htm
- Yusuf G., and Olayemi I.K., 2015, Spatial variation in physicochemical characteristics of wetland rice fields mosquito larval habitats in Minna, North Central Nigeria, *International Conference on Agricultural, Ecological and Medical Sciences*, 3: 53-56



Reasons to publish in BioPublisher

A BioScience Publishing Platform

- ★ Peer review quickly and professionally
- ☆ Publish online immediately upon acceptance
- ★ Deposit permanently and track easily
- ☆ Access free and open around the world
- ★ Disseminate multilingual available

Submit your manuscript at: <http://biopublisher.ca>

