

ENVIRONMENTAL TECHNOLOGY & SCIENCE JOURNAL (ETSJ)

VOLUME 8

NUMBER 1

JUNE, 2017

ISSN: 2006-0459

PUBLISHED BY: School of Environmental Technology Federal University of Technology,

P.M.B. 65, Minna, Niger State, Nigeria Copyright@2017

Aim and Scope

The Environmental Technology and Science Journal (ETSJ) is devoted to the publication of papers which advance knowledge of practical and theoretical issues of the environmental technology. The aim of the journal is to provide an avenue for dissemination of academic research findings from various disciplines of the environment, engineering, pure and applied sciences, arts and social science, which have materials that emphasize on environmental issues.

ETSJ Policy

eTSJ prohibits an author from submitting the same manuscript for concurrent consideration by two or more publications. The author must ensure that when a manuscript is submitted to ETSJ, the manuscript must be an original work. The Author should check the manuscript for any possible plagiarism using any program such as TURNITIN or any other software before submitting the manuscripts to ETSJ. Authors are not permitted to add or remove any names from the authorship provided at the time of initial submission without the consent of the Journal's Editor-in-Chief.

Guide for Authors

Please read the guidelines and follow the instructions carefully; doing so will ensure that the publication of your manuscript is as rapid and efficient as possible. The Editorial Board reserves the right to return manuscripts without review that are not prepared in accordance with these guidelines.

Manuscripts should not be more than 15 pages of A4 paper size (including abstracts, tables, figures, references and appendices) typed in double spacing,

times new roman and font size 12.

2. Each paper must be compiled in one column; all text should be left and right justified. The page settings to be 25cm (1 inch) each for the top, bottom, left and right margins of an A4 paper. Decimal numbering of all sections of the article is recommended (with the exception of the reference and acknowledgement sections).

3. To ensure anonymity in the peer reviewing process, articles must be structured in such a way that the title of the paper, authors' names, address, and affiliation are written on the first page, followed by abstract and the body

of the paper in that order.

4. Each article should contain an abstract of not more than 300 words.

5. Each article should contain between 5 and 7 key words (avoid use of phrases).

6. Use the SI (*Systeme Internationale*) units and symbols, while Unusual Characters and symbols should be explained in a list of nomenclatures.

7. The journal's referencing style is the APA format.

8. Figures, Tables and Graphs are to be placed at the appropriate places in the paper and not at the end of the paper.

9. Acknowledgement is optional, except for researches being funded through

research grant awards.

10. Authors should avoid using headers, footers and special page breaks with

the manuscripts.

- 11. The manuscript is to be sent via electronic means to: estj@futminna.edu____
 for a blind peer review, which under normal circumstance will not exceed
 weeks, after which the status of the article will be communicated to
 Author(s).
- 12. Author(s) with positive review will be asked to effect corrections or revisor after which the camera-ready manuscript is to be emailed estj@futminna.edu.ng and to be accompanied by evidence of payment publication fee.

13. The Editorial board is not responsible for the information or views expressibly the author(s).

Frequency of Publication

The journal is published twice a year in June and December

Subscription Details

The 2017 subscription rates for hardcopies of the journals including postage are Individual within Nigeria: N2,000 per copy and N1,000 for postage Individual outside Nigeria including postage \$ 100 Institution within Nigeria: N5,000 per copy and N1,000 for postage Institution outside Nigeria including postage: \$ 150

Correspondence

All correspondence should be addressed to The Managing Editor
Environmental Technology & Science Journal SET, FUT, Minna, Nigeria
Email: etsj@futminna.edu.ng
Phone: 08051703663, 08036534507

Published By

School of Environmental Technology, Federal University of Technology. Minna, Nigeria.

	-,	
fies beneficial t		
roposed at desig	п.	
siy inadequate an	I.nntpnto	3
The author	-011101111	
firmer proactive	(
olicy actions and	1-13	Effect of Pozzolana on the Compressive Strength of Concrete in
es are necessary to		Chemically Aggressive Environment
maintain curren		Aka, A., Musa, A. A., Kaase, E.T. & Tukur, A.
hence make the		11. 11. 11. 11. 11. A., Rause, E.1. & Tukur, A.
ts sustainable in	14-22	Accomment of D 11' D
	17-22	Assessment of Public Perception of Climate Change Issues in
		Minna, Niger State, Nigeria
ues interrogated in		Odegbenro, F.J. & Ojoye, S.
s towards making		
ter place. Happy	23-35	Geospatial Analysis of Shoreline Dynamics in the Coastal Areas
		of Cross River State Nigeria
		Adebola, A. O, Adegboyega, S. A. & Ibitoye, M. O.
		2. O., The gooyegu, S. A. & Tolloye, M. O.
	36-46	Evaluation of Key Design Flaments for DI.
		Evaluation of Key Design Elements for Play-Learning
		Environment in Elementary Schools in Minna, Nigeria
		Ayuba, P. & Akpama, D. S.
	47-60	Car Dark Hann 134
u	, 00	Car Park Usage and Management in Federal Institutions of
		Zaria Metropolis, Kaduna State-Nigeria
		Oluwole, M. S., Masugari, D. Y. and Elegonye, I. I.
	61 74	D 11 1 111
	61-74	Reliability of the Liquidation Valuation Models of Selected
		Real Estate Collaterals in Niger State
		Babatunde, I. O.
	77.04	
	75-86	Appraisal of Recycling of Deconstructed Building Materials
		from Selected Renewal and Renovation Projects in Minna
		Ayuba, P. & Albert, B. S.
		3
	87-102	A Geospatial Analysis of Coastal Land use/Land Cover Pattern
		and Shoreline Changes in Akwa-Ibom State, Nigeria
		Adebola, A. O., Ojoye, S. & Ibitoye, M. O.
	103-122	The Risk of Residing in Proximity to Illegal Waste Dump Site
		in Sabon Wuse, North-Central, Nigeria
		Abd'razack N. T.A. Madawara S.O. II.
	3	Abd'razack, N. T.A., Medayese, S.O., Umaru, E.T. & Shaibu, S.I.

123-135 Developing a Mass Appraisal Approach for Residential Properties in Minna Metropolis Liman, H. S., Olatunji, I. A., Morenikeji, G., Olubajo, O. O. & Usman' Z. D. 136-150 Assessment of Urban Heat Island in Zaria Urban Area Fidelis, C. D. & Musa, W. K. 151-160 Assessment of Safety and Health Performance of Contractors' Construction Projects in Nigeria Using SHASSIC Method Yakubu. D. M. 161-174 A Simulation Framework for Housing Choice Optimality: Decision-Support Guide for Housing Procurement Service in Abuja Olatunji, I.A. 175-185 The Effects of Road Quality on Commercial Land Use Pattern in Makurdi Urban, Benue State, Nigeria Umoren, V. & Mchi. A. 186-206 Assessment of Residential Attributes of Lagos State Development and Property Corporation's Residential Schemes on Resident's Well-Being Johnson, M. B., Adebamowo, M. & Adejumo, O.

The Risk of Residing in Proximity to Illegal Waste Dump Site in Sabon Wuse, North-Central, Nigeria

Abd'razack, Nelson T.A., Medayese, S.O., Umaru, E.T. & Shaibu, S.I.

Department of Urban and Regional Planning, Federal University of Technology, Minna. Corresponding E-mail: abdrazacknta@futminna.edu.ng

Abstract

for

oing

and

eline

dhra

tock

1. K

ation

sing

For

ndia.

nces.

260)

of

This study investigated the spatial distribution of solid waste dump sites in the study area, established the residents' perceived level of exposure to solid waste hazards, and the effect of living near refuse dump sites. Systematic random sampling technique was used to administer 500 questionnaires to the households in the study area. The result of the study indicated that there is only one legal dump site located in Buntu area; there are 19 illegal dump sites within the town. The proliferation of illegal dumpsite was as a result of convenience and open spaces within the town. The most frequently used method of waste disposal is on weekly basis. The use of plastic bucket is favoured for storage. The level of health risk associated with living close to dumpsite shows that a total of 878 houses are at the severe risk, while, 1,898 houses are at mild risk level. The study concludes that There should be a total clearance of the existing illegal dump site and proper monitoring of the waste management in the town to forestall illegal dumping, and adequate information to residents and awareness on the danger of consequences of indiscriminate dumping of refuse in an undesignated dump site.

Keywords: Illegal Dump Site, Solid Waste, Waste Management, Environment, Urbanisation.

Introduction

Waste generation is a result of consumption of resources and other activities of mankind. As this is an unavoidable event in day to day living, there is need for waste generated to be managed. Lack of proper management of waste always results environmental and health challenges. Globally, the process of management requires a chain activities from collection, sorting, transporting and disposal. If any of these lines of management is disrupted, it affects the management of the waste. How this may be efficiently done poses a problem in many societies today. Another natural process, population growth, makes waste management even more challenging; more people in a specific geographic location would imply a higher level of waste generated, hence more waste to contend with in that area. As poorly managed wastes are perceived as environmental hazards of high significance, the societies' inability to manage waste generation effectively play no small role in increasing extant environmental pressures (Karanjit et al., 2007).

The double function of Sabon Wuse as the Local Government Headquarters and the resettlement of displaced people from Abuja due to its proximity to the Federal Capital has increased the population of the town and thereby increased the solid waste being generated. The responsible agencies, such as Niger State Environmental Protection Agency (NISEPA) and Tafa Local Government Council Public Health Department did not have any improvement in the capacity to handle the massive garbage being generated in the town. The local Government and NISEPA could not control the urban development as well as waste disposal sites that are scattered all over the town, which resulted into poor and unclean environment.

General Concept of Waste

Waste generally is an unwanted byproduct of man's activities. It is defined by Lutui (2001) as "materials that currently have negative value to their owner, that is the generator incurs costs in managing them (importantly this does not prevent them from having positive value to another owner at another location in space or time)" (Lututi, 2001: p3) and are disposed of. Solid waste is a matter of time and places, as what is a waste can become raw material in another place. For example, the recycling plant uses solid waste product as raw material for production of other items (Sanusi, 2010). Research has shown that there are two different variance of waste, that is, solid waste (effluence) and liquid waste (foul water, semi- liquid and the gaseous liquid Biogas). "Waste" does not have a generally concise definition, however waste is generally known to be any unwanted material (Lutui, 2001).

Morrisson, Wray, Dever, and Dusbaun (2000) define waste as "any matter prescribed to be waste under national legislation, any material listed as a

waste in appropriate schedules, and ingeneral, any surplus or reject material that is no longer useful and which is to be disposed off". This definition uses two words that is important to this study "waste is neither wanted nor has value to the owner" and which in turned to be disposed off. Waste has an unwelcome and often unnoticed effection the environment and hamped development and civilization (Kim and Gobalan, 1997) and can be traced to the growth of industrialization in modern economies (Holmes, 2000).

Solid Waste

Solid waste could be defined as any non-liquid and non-gaseous substance produced out of human activities which are regarded as being useless. It could take the form of garbage, refuse, and sludge (Nwosu and Olofa, 2015). Solid waste can also be regarded as materials which are no longer in use. This includes household garbage, unwanted materials from commercial and mining activities. Solid wastes are categorized in to three (3) types, based on the composition. These are:

i. Biodegradable; mainly composed of green waste, garbage. trash.

ii. Non-biodegradable; this consist of scraps, synthetic (plastic, rubber. leather) materials, metals etc.

iii. Semi-biodegradable; this consist of rubbish (paper, cartons, wood.)

Solid waste can also be defined as those materials that are generated, resulting from man's activities and are not in form of liquid or gas but are compacted and substantial.

The Ri

Types of D

fr w re n u it

C

I I

11.

iii.

iv.

abdra

Types of Solid Waste

and in

naterial

ch is to

in uses

to this

in turn

has an

effect

amper

im and

to the

nodem

as any

stance

which

could

terials.

Wanted

nining

orized

on the

rbage,

DESCRIPTION

DESERTO

nacted

This

- Domestic Solid Waste: These consist of rubbish and garbage from household (i.e Municipal waste). Other examples are reminant of food materials, old newspapers, spoilt kitchen utensils, cartoons, baby toys etc. it is always the principal focus of solid waste management till date (Abdrazack, Yusuf and Utange, 2013). In Sabon Wuse, The Niger State Environmental Protection Agency are those that saddled with the responsibility of collection and disposal of Municipal solid waste management.
- ii. Agricultural Solid Waste: These wastes resulting from different agricultural activities, include cultivated waste (weed); harvested wastes from plants, field and tree crop wastes; the dung from Animal production and waste from operation of feed lots as described by United Nation Environmental Programme (UNEP, 2010).
- iii. Commercial Solid Waste: These include all solid waste which emanate from business and profit making activities such as financial institution (Banks, market, stores, super market), educational institution (theatres, lecture halls and class rooms), hospitals (Abdrazack et al., 2013)
- iv. Industrial solid waste: These are solid waste product from the production of goods. They include all solid waste which result from both light and heavy

- industries. In most developed countries, industrial solid waste account for the most divasting tonnes of general waste hazards. In the USA for examples, industries generate one third of the general waste produced in the country. They are mostly as a result of industrial processes of manufacturing operation inherent in the country. Examples of such operations are: processing plant, repair and clearing establishment. refineries, manning and mineral operations (UNEP, 2014).
- v. Special Solid Waste: These types of waste comprise of waste from street sweeping, road side litter, debris, dead animals, abandoned vehicles and litters of store drains. They are called 'special waste' because it is impossible to predict where they will be found (Abdrazack et al., 2013).

Literature Review

Rapid increase in the types and volume of domestic waste (either liquid or solid waste) as a result of continuous economic growth, urbanization and industrialization, is becoming escalating problem for national, state and local governments to ensure effective and sustainable management of waste. In the year 2006, it was estimated that the total amount of municipal solid waste (MSW) generated worldwide reached 2.02 billion tons of waste, representing a 7% annual increase since 2003 (Global Waste Management Market Report, 2009). It was further estimated that

between year 2007 and 2011, global generation of municipal waste would rise by 37.3%, equivalent to roughly 8% increase per year. Based on incomplete reports from its participants, The Basel Convention estimated that about 318 and 338 million tonnes of solid waste were generated in 2001 (World Bank, 2010).

Owing to an increase in human population, industrial and technological revolutions, waste management has become increasingly complex (Akinbile and Yusuf, 2011). Marshall and Farabahksh (2013) observed that population growth and subsequent innovations have not only led to changing consumption patterns across borders but have also placed considerable pressure on waste management services. This is so because as the world races towards its urban and more complex future, the byproduct of the urban lifestyle, waste, is growing at even faster rates. At present, global solid waste generation is growing approximately at 1.3 billion tonnes per year and is expected to double by the year 2025 (Hoornweg and Bhada-Tata, 2012). The pressure placed on waste management services has consequently led to widespread inefficiencies that are mainly due to lack of funds, improper infrastructure, inadequate waste collection services, unlicensed waste management activities. insufficient minimisation and limited waste related legislation (Nahman and Godfrey, 2010). Inefficient waste management services have also led to extensive illegal dumping. According to Zurbrugg (2002), illegal dumping refers to an

instance where solid waste is disposed of in inappropriate manners in places such as in drains, roads, near rivers and on private and public land that is not legally approved for such a use Dumped materials typically include furniture, garden waste, discarded appliances, household rubbish, building rubble, broken glasses, old tires hazardous materials such as illegal pestilences, and metal contamination and abandoned automobiles (United States Environmental Protection Agency, 1998).

Illegal dumping of waste products is a recognised problem in much of the vorld and Nigeria. At a global level cumping has resulted in increased costs associated with clearing and clean-up efforts. For instance, in Britian, clearing waste that is dumped every 35 seconds has cost the government millions according to the British Broadcasting Corporation (BBC news, 2005). There have also been instances of developed countries dumping waste in the less developed countries. recen occurrence is the dumping of old and broken television sets in Ghana by one of the United Kingdom's leading was and recycling companies. This is said have been done in violation of the laws in relation to the flow of waste developing countries (Wasley, 2011). the city of Abidjan in Côte d'Ivoire vast amounts of toxic waste released a tanker registered under a Dutch O Trader resulted in the death of people and left many others suffer diarrhoea. vomiting nosebleeds (Johnson, 2006). It uncertain as to how the toxic pollutaentered the country. Such occurrence

thus make i monitor an ensure prop Bylaws for presented, b regulations i

Dumped s implications and the qua waste contar This occurs seeps throug with substar forms a sub According (2011), leac sites also Toundwater not properly make the su flooding as the waste can 1 culverts (Un Programme, combat illega effects, resid sites. This, ho on the enviro severe erosio which limit mimals such s they die fi from waste si Lebris (Projec

The presence cause serious reside treeding ground has rats tates Env Agency, 199

thus make it important for countries to monitor and to enforce policies that ensure proper waste disposal services. Bylaws for illegal dumping are clearly presented, but the enforcement of these regulations is unclear.

100

SÜS

ng

mg.

255

tio

W/S

Bo

In

Dil

Dumped solid waste has serious implications for the health, environment and the quality of life. Dumped solid waste contaminates both soil and water. This occurs when water from rainfall seeps through dump waste and mixes with substances within the waste and forms a substance known as leachate. According to Akinbile and Yusuf (2011), leachate released from waste sites also poses a high risk to groundwater and surface water if it is not properly managed. Dumpsites also make the surrounding areas prone to flooding as the different components of waste can block drains, creeks and culverts (United Nations Environment Programme, 2005). In an attempt to combat illegal dumping and mitigate its effects, residents in rural areas burn sites. This, however, has a direct impact on the environment because fires cause severe erosion due to burning of trees which limit vegetation growth. Small animals such as birds are also affected as they die from feeding on materials from waste sites and by being stuck in debris (Project Green Sweep, 2011).

The presence of an illegal dumpsite can cause serious health problems for nearby residents as it is an ideal breeding ground for disease-vectors such as rats and mosquitoes (United States Environmental Protection Agency, 1998). Certain respiratory

illnesses such as 'asthma tuberculosis are also linked to illegally dumped waste (Etengeneng, 2012). The occurrence of illegal dump sites not only affects health and the environment, but also the quality of life. According to Madava (2001), illegal dumpsites have adverse effects on the basic human rights of people with regards to the standard of living. This is so because hazardous wastes are prospective pollutants of the biophysical and human environment. This is so because the presence of dumpsites does not only deteriorate the quality of environment, but also breaches human rights as it has an impact on community pride. More often than not, the sight and smell that emerge from illegal dumpsites are unpleasant and diminish the land value. Illegal sites that mostly constitute of flammable substances and gases are vulnerable to fires. More also as stated by United State Environmental Protection Agency (USEPA), impact is not limited to the aesthetic of landscapes, but diminishes the value of surrounding properties significantly, forcing residents to vacate their homes to further places (USEPA, 2005).

The situation in Nigeria is different in that the process of urbanisation in the country is unplanned; there is population explosion as decentralisation of government has turned villages to local government headquarters thereby attracting more people to the urban centres (Okpala, 2004). This has led to high production of solid waste, the traditional attitude, poverty and high level of ineptitude on the agency responsible for waste management has

contributed significantly to turn our towns and cities to garbage cities rather than serene environment (Oyelola, and Babatunde, 2013). Table 1 shows the level of waste generation in some selected cities in Nigeria. There are many illegal waste dump sites round the cities. The population of the cities are grasping with the environmental and health risks associated with unclean environment. This has led to breeding of pathogenic condition (breeding ground for mosquitoes, rodents and airborne diseases (Ogwuelika, 2009).

Table 1: Volumes of solid waste generated in some Niperian cities (Tonnes/year)

Cities	Tonnage / Month	Densit y	Kg/Capital day
Lagos	255,556	294	0.63
Kano	156,676	290	0.56
Ibadan	135,391	330	0.51
Kaduna	114,433	320	0.58
Port/Cou	117,825	300	0.60
rt			
Makurdi	24,242	340	0.48
Onitsha	84,137	310	0.53
Nsukka	12,000	370	0.44
Abuja	14,785	280	0.66

Source: Ogwuelika, 2009

Table 1 has clearly shown that the city of Lagos generates more waste than every other city in Nigeria due to population of the state. It is worthy to know that Kano has a larger population when compared with Lagos state but Lagos state generates more waste than

Kano. Ogwueleka concluded that, on the average the Southern part of the country generates about 0.63 kg per capita per day while the Northern part produces about 0.56 kg per capita per day. On the average, the country produces about 0.60 kg per capita per day.

Study Area

Sabon Wuse is a Community in Tafa Local Government Area of Niger State. Nigeria, adjoining the Federal Capital Territory Abuja. It is located on latitude 9°33"N and longitude of 71°51"E with the projected population of 71,074 people as of 2015 (NPC, 2011). The geographical location of Sabon Wuse is shown in Figure 1 in the context of Nigeria and Niger State. The study area has a long range of hills and ridges on the western side of the town, which serve as restrictions to physical development in that part of the town. Presence of many slopes and valleys in the town encourage rapid erosion and formation of gully channels that are common in the town. Incidence of flash floods is a common occurrence especially in high density areas. While farming remains the chief occupation in Sabon Wuse, the town is noted for mat making and export of cotton weaving and dyeing.



Figure 1: 1 Source: U

Research

The meth was a m integrate Systems quantitat This app that the of the iss illegal d the toy neighbou dumpsite were ide of each taken ar GPS. Th were th Informat called A coordina displayi dumpsit dump dumpsit refuse samplin adminis

question the dun




Figure 1: Location of Sabon Wuse in the National and State context Source: URP Dept. FUT Minna.

Research Methodology

DO

a per

Tafa

tate

with

.074

The

se is

I of

area

S OE

sical

WIL

SI

and

are

moe

The methodology adopted for the study was a mixed-methods approach which integrated Geographical Information Systems (GIS) mapping, qualitative and quantitative research (Bryman, 2006). This approach was selected to ensure that the data presented a holistic view of the issue at hand. In order to map out illegal dumpsites throughout the town, the town was divided into four neighbourhoods and all illegal dumpsites within the neighbourhoods were identified. The GPS co-ordinates of each illegal dumpsite were then taken and recorded using a hand-held GPS. The co-ordinates of the dumpsites were then entered into Geographic Information Systems (GIS) software called ArcMap. This software used the coordinates to generate map displaying the location each dumpsite, the coordinate of each refuse dump point, height of the refuse dumpsite and area coverage of each refuse dump site. Systematic random sampling technique was adopted in administering a set of pre-tested questionnaire on households living near the dump sites. To do this, ten of the

twenty illegal dumpsites in the town were randomly selected from the four neighbourhoods in the town. A set of self-administered questionnaires designed to include dichotomous, Likert rating, checklist and open-ended questions, was used to obtain primary information on respondents' attitudes and views on solid waste dump sites. Forty residential buildings within the closest proximity (as the crow flies) from all the dumpsites purposively selected. A household was randomly sampled in each of the selected building and the household head provided the required data. It was only in the absence of the head that available oldest adult member of the household was interviewed. 500 copies of the questionnaire were administered out of which 370 copies were correctly filled, returned and used for analysis of the study. Both descriptive inferential statistics (Analysis Variance and Tukey Post-Hoc tests) were used in analyzing data collected at $p \ge 0.05$.

The level of exposure to environmental and risk hazard when living in

proximity to dump site was measured using Chung and Poon (2001) scale. The scale measured the distance to dump site to determine the level of rsk and exposure to hazard thus:

- i. 0.00 50.00 metres = Very Severe Risk
- ii. 50.01 100.00 metres = Severe Risk
- iii. 100.01 150.00 metres = Moderate Risk
- iv. 150.01 200,00 metres = Mild Risk
- v. >250.00 metres = No Risk

Research Findings and Discussion

Results of the study are presented under four main headings: Spatial distribution of illegal dumpsites; Assessment of solid waste disposal method; Residents' perception of level of exposure to solid waste hazards; and, Effects of living near solid waste dump sites.

Spatial Distribution of Waste Dump in Sabon Wuse

the bv Investigations conducted researchers revealed that although there is only one legally recognized solid waste dump site in Sabon Wuse town. nineteen illegal ones were identified by the authors. Spatial distribution of the dump sites are presented in Table 2 and Figure 2. From Figure 2, it is revealed that illegal dumpsites of solid waste were found throughout inhabited parts Distance of the legal of the town. dumpsite and inadequate enforcement of environmental sanitation laws and edits, especially in relation to illegal dumping of solid wastes, might be responsible for the proliferation of illegal solid waste dump sites in the town.

The Risk of I

Location o the Dump Site

Buntu

AngwanYas

Katampe

Katampe Bridge Aso

Behind SarkinAso House Central Mosque Aso AngwanTof

Behind Aso Market AngwanYa

Hausawa

Abuja Kadı Expressway Abuja Kad Expresswa Abuja Ka Expressw Abuja Ka Expressw Abuja Ka

Expressw

abdrazackni


Abd'razack, Medayese, Umaru, Shaibu The Risk of Residing in Proximity to Illegal Waste Dump Site in Sabon Wuse, North-Central, Nigeria

of living


te Dump

by the ugh there zed solid ise town, ntified by on of the ble 2 and revealed id waste ted parts he legal prcement aws and o illegal night be ation of s in the

SN	le 2: Spatial Loc Location of	Status	Location and Coordinate		Size of the Dump Site					
	the Dump	of		mp Site						
D.	Site	Dump Site	Northern	Eastern	Length (m)	Breath (m)	Area (m²)	Height (m)	Vol. of Waste (m ³)	
	Buntu	Legal			100	70	7,000	3	21,000	
-	A 37 1.	711	9°17.932'N	7°13.707'E						
-	AngwanYashi	Illegal	0010 00001	701442015	5	3	15	1	15	
3	Katampe	Illegal	9°18.265'N	7°14.430'E	10	-	50			
	raumpo	megai	9°18'290"N	7°14.876'E	10	5	50	1	50	
4	Katampe	Illegal	7 10 250 11	7 14.870L	12	6	72	1	72	
	Bridge		9°18.125'N	7°14.718'E	12	· ·	12	1	12	
5	Aso	Illegal			50	20	1,000	1	100	
-			9°17.962'N	7°14.575'E						
5	Behind	Illegal			40	30	1,200	5	6,000	
	SarkinAso House		9°17.884'N	7°14.542'E						
7	Central	Illegal			20				2020/0000 2000	
	Mosque Aso	megai	9°17.924'N	7°14.417'E	20	11	220	1	220	
3	AngwanTofa	Illegal	J 17.524 IN	/ 14.41/E	70	45	3,150	4	12 (00	
	3		9°17.997'N	7°14.096'E	70	43	3,130	4	12,600	
9	Behind Aso	Illegal		. 1.10,02	10	4	40	1	40	
	Market	90 -1 00	9°18.169'N	7°14.228'E			.0	* 1	40	
10	AngwanYashi	Illegal			10	5	50	1	50	
	**	***	9°18.327'N	7°14.365'E						
11	Hausawa	Illegal	0010 50507		4	2	8	1	8	
12	Abuja Kaduna	Illagal	9°18.535'N	7°14.366'E						
	Expressway	Illegal	9°18.591'N	7°14.388'E	6	3	18	4	72	
13	Abuja Kaduna	Illegal	9 10.391 N	/ 14.388 E	18	2	26	2	100	
	Expressway	mogui	9°18.574'N	7°14.389'E	10	2	36	3	108	
14	Abuja Kaduna	Illegal		7 11.507 E	16	4	64	Ĭ	64	
	Expressway		9°18.549'N	7°14.367'E	7.7		0.1	*	UT	
15	Abuja Kaduna	Illegal			5	3	15	1	15	
	Expressway	100001 KS	9°18.529'N	7°14.345'E						
16	Abuja Kaduna	Illegal	0010 75 70 7	V22 24 24 24 25 25 25 25 25 25 25 25 25 25 25 25 25	20	10	200	4	800	
17	Expressway Abuja Kaduna	111 1	9°18.525'N	7°14.343′E	12121	100				
	Expressway	Illegal	9°18.494'N	701// 21017	40	20	800	8	6,400	
18	Abuja Kaduna	Illegal	7 10.474 IN	7°14.318'E	15	7 •	105		400	
	Expressway	1110541	9°18.477'N	7°14.307'E	13	1 •	105	4	420	
19	Abuja Kaduna	Illegal		, 14.50/L	6	4	24	2	48	
	Expressway		9°18.484'N	7°14.293'E	<i>y</i>	out 1	۷.	2	70	
20	Abuja Kaduna	Illegal			12	8	96	5	480	
N	Expressway		9°18.472'N	7°14.282'E				0770		



he most used acilities in the acket (23.4% ylon Bag (11.7%), reasonsy to converse posal because the undesign often saddle mping it off boices. Table acilities used least the use.



Nisk Associate
Vaste Proxin
VabonWuse

mdrazacknta@fu




Figure 2: Spatial location of both legal and illegal waste dump site in SabonWuse

Assessment of Solid Waste Disposal Methods

Solid Waste Disposal Methods in the Study Area

Table 3 shows the level of assessment of waste collection and disposal in Sabon Wuse. The analysis indicated that 21.4% of respondents dump their solid waste to an informal collector on daily basis; about 61.7% dump their solid waste indiscriminately on weekly

basis which is a clear indication that they often dump their waste when the are less busy which happens that a Saturday and Sunday when they do not go to work, While 14.9% fortnightly, this deal with dumping refuse at any time they were able to see the informal refuse collectors which does really have a prescribed day and 6.0% on monthly basis.

Table 3: Frequency of Waste Disposal in

requency of Disposal	Number of Respondents	(%)	
Daily	79	21.4	
Weekly	228	61.7	
Fortnightly	55	14.9	
Monthly	7	1.9	
Total	370	100.0	

The most used municipal waste storage acilities in the study area are; Plastic Bucket (23.4%), Cotton Sack (18.2%), Nylon Bag (21.4) and Bagco Bag 11.7%), reasons being that they are easy to convey to various area of disposal because of their light weight and they are less expensive and very few people in the study area are aware of the undesignated dump site so they are often saddled with no option than dumping it off available place of their choices. Table 4 shows various storage facilities used by households in Sabon Wuse.

Table 4: Method ofWaste Collection in

Storage Facilities	Number of Respondents	(%)
Palm Tree Front Basket	24	6.5
Plastic Bucket	87	23.4
Cotton Sack	67	18.2
Drum	60	16.2
Nylon Bag	79	21.4
Bagco Bag	43	11.7
Others	10	2.6
Total	370	100.0

Risk Associated with Illegal Solid Waste Proximity to Houses in SabonWuse

Figures 3 and 4 show the extent of closeness of compounds to refuse dump in the study area. The risk of staying close to refuse dump was measured between 100 meters and 200 meters. The risk ranges from fatal, severe and mild (Chung and Poon, 2001). Satellite image of the study area was used to capture the number of compounds that were within the 100 meters (Severe) and 200 meters (mild) risk. This process is known as buffering. The buffering uses radius to determine the area of coverage. The total numbers of structures at the study area is 3,571. The total number of compounds at severe risk (100m) is 878 which is 24.6% and at mild risk (200m) is 1,898which is 53.2% of the total number of structure within the study area. Dumped and untreated solid waste has serious implications for the health, environment and the quality of life of people, therefore, Chung and Poon, 2001 indicated that residents within the buffering region of the refuse dump at 100m are at higher risk than those at 200m distance. This is due to the fact that level of exposure to hazard is high, hence the ability of rodents and other vectors to transmit infections is also high. The implication of this is that the closer the residents to dump site the higher the risk of both environmental and health hazard in Sabon Wuse.

The analysis of the proximity to risk of exposure to hazard is measure using the buffer analysis as shown in Figures 3 and 4. The buffer that was used followed the 100 metres and 200 metres risk zones. The 100 metres shows a severe risk associated with living within the catchment of such environmental and health risk, while, 200 metres

of

buffer on the other hand shows a mild risk of exposure to both environmental and health risk as posited by Chung and Poon (2001), it is better to avoid living

in an area where there is risk when mild or s evere.




Figure 3: 100m Risk radius showing houses closer to waste dump site in Sabon Wuse

Figure 4: 20

Residents of Hazard Area.

Table 5 residents to solid we The exten measured range of to (SA); Agr (D) and St




Figure 4: 200m Risk radius showing houses closer to waste dump site in Sabon Wuse

Residents' Perception of the Types of Hazard Exposed to in the Study Area.

Table 5 shows the perception of residents to various risks of exposures to solid waste hazard in the study area. The extent of perception of the risk is measured using Linkert Scale. The range of the scale is Strongly Agreed (SA); Agreed (A); Fair (F); Disagreed (D) and Strongly Disagreed (SD).

The specific risk and hazard exposed to by the residents based on their perception include the following: Foul water, Dust during hamattan, Dust at the beginning of rainy season, Dust at the end of rainy season, Rats and Rodents attack. Houseflies, Mosquitoes, Defacing the physical appearance of the environment, Children playing off the dump, Injuries from sharp objects of the dumps,

Pollution of water bodies and Smoke from burning of the waste.

Table 5: Residents' Perception of the Type of Hazard exposed to in the Study Area

S/	le 5: Residents' Perception of the Type of Huz Hazard	Fi	xed Ha	zard G	rade (1-	5)	Ground
N	Allen C	SA	A	F	D	SD	Total
1	Domestic Sewage	47	89	62	130	41	370
2	Sand Dust	29	26	202	60	53	370
3	Erosion	38	50	38	67	187	370
4	Poor Physical Apperance of the	199	24	103	17	24	370
5	Environment Children Playing on dump sites	7	19	48	125	171	370
6	Rats and Rodents	7	46	38	55	223	370
7	Mosquito Breeding	147	72	50	82	19	370
8	Houseflies	113	159	55	14	29	370
9	Injuries from Sharp Objects	5	14	19	183	149	370
10	Epidemics	7	34	46	84	199	370
11	Pollution of Water Source	17	7	10	130	206	370
12	Smoke from burning of solid waste (open	31	10	50	58	221	370

Table 6 shows the responses of the residents in respect to their exposure to hazard of solid waste. These data are known as Variable Numbers of Respondents (VNR) and the Fixed Hazard Grade (FHG) is ranged from 1-5 of which 5 is the most dangerous. The procedure of calculation of the households' level of exposure to solid waste Hazards is shown thus:

$$AG = \underline{\sum} FHG \ XVNR \tag{1}$$

$$(SS_0)$$

Actual Grade (AG) = <u>Fixed Hazards Grade</u> (FHG) X Variable Numbers of Respondents (VNR)

Sample Size (SSo)

Where:

Fixed Hazards Grade (FHG): ranging from 1 to 5, were '1' is Very Good; '2' Good; '3' Average; '4' Very Poor and 5 is Extremely Dangerous.

Variable Numbers of Respondents (VNR): the total number of responses for each grade.

Sample Size (SS_o) - it is referred to as the sample size of the study area, which is 370.

Table 4 also shows that erosion, children playing on the dump, injuries from sharp objects of the dumps fall within the fair condition, meaning that their effect as a result of the presence of these refuse dumps is minute/ fair on the people around that area. In the same vein, it shows that: foul odour, dust at the beginning of rainy season and house flies falls with the average condition, which indicate that most of the just itemized hazards exist on an average scale in the study area as a result of those refuse dumps being present in that area.

The Risk o

In addition, dust during rodent outbre the physic environment smoke from within the vanking scale

Table 6: Hous

S/ N	Hazard
1	Domestic
- 200	
2	Sand Du
3	Erosion
4	Poor Phy
	Environr
5	Children
6	Rats and
7	Mosquite
8	Housefli
9	Injuries:
10	Epidemi
11	Pollution
12	Smoke f
	dump)

Perception
Waste Haz
Table 7 sh
exposed to
site in the s
exposed to
following:
from foul
sight of t
leaving th
persecution
sanitation
being repu
friends an
visit.

abdrazackni

In addition, Table 4 also indicates that: dust during hamattan, rats and other rodent outbreak, mosquitoes, defacing the physical appearance of the environment, pollution of water and smoke from burning of waste fall within the very bad condition on the ranking scale, which means that there is

und

70 70 70

dents onses

to as

sion.

uries

s fall

that.

ice of

ir on

same

ast at

iouse

ition.

just

erage

alt of

n that

a very high existence of health hazards due to the presences refuse dump sites in the study area. It has made the area susceptible solid waste hazard and if urgent measures are not taken, it would be more terrifying in the nearest future.

Table 6: Households Level of Exposure to Solid Waste Hazards

S/	Hazard		Fixe	d Hazard	Grade (1-5)	
N		Poo r	Fai r	Averag e	Very Bad	Extremel y Dangerou s	Actual Grade
1	Domestic Sewage	.13	.48	.51	1.40	.55	3.07
2	Sand Dust	.01	.08	.16	1.97	2.01	4.23
3	Erosion	.08	.14	1.64	.65	.71	3.22
4	Poor Physical Apperance of the Environment	.02	.18	.37	.91	2.69	4.18
5	Children Playing on dump sites	.31	.86	.45	.12	.39	2.12
6	Rats and Rodents	.02	.25	.31	.73	2.53	3.95
7	Mosquito Breeding	.10	.10	.39	1.35	2.31	4.17
8	Houseflies	.02	.25	.31	.60	3.02	4.19
9	Injuries from Sharp Objects	.40	.39	.41	.88	.26	2.34
10	Epidemics	.54	.13	.84	.18	.36	2.05
11	Pollution of Water Source	.05	.04	.08	1.40	2.79	4.36
12	Smoke from burning of solid waste (open dump)	.08	.05	.41	.62	2.99	4.16

Perception of Residents to Solid Waste Hazard in Sabon Wuse

Table 7 shows level of health hazard exposed to when residing close to dump site in the study area. The health hazard exposed to varies, and it included the following: fear of disease, discomfort from foul odour, irritation from the sight of the dump, fear of people leaving the neighbourhood, fear of persecution by the environmental sanitation body, the neighbourhood being repulsive to needed businesses, friends and relatives not wanting to visit.

Table 8 shows the response of the residents to risk of health hazard when residing close to refuse dump in the study area. The variable was then grouped as Variable Numbers of Respondents (VNR) and the Fixed Hazard Grade (FHG) is ranged from 1-5 of which 5 is the most dangerous. The total number of respondent is 370. The procedure of its calculation and judgement follows the same procedure.

Table 7: Residents' Perception to Level of Health Hazard in the Study Area

S/	Perception of Risk Exposed to						
N	rer i de Rok de 12 de se de la mir p	Poo	Fai	Averag	Very	Extremel	Actual
		r	r	e	Bad	y Dangerou	Grade
						S	
1	Fear of disease	12	38	101	144	75	370
2	Discomfort from foul Odour	22	34	139	82	175	370
3	Irritation from the sight of the dump	26	12	79	197	113	370
4	Fear of people leaving in the neighborhood	34	77	108	86	65	370
5	Fear of persecution by the environmental sanitation body	26	82	77	82	103	370
6	The neighborhood being repulsive to needed businesses	31	10	43	127	159	370
7	Friends and relatives not wanting to visit my family	26	14	67	123	139	370
8	Blocking of roads	21	46	53	151	99	370

The result of Table 8 also shows that fear of disease, discomfort from foul odour, irritation from the sight of the dump, fear of people leaving the neighbourhood, fear of persecution by the environmental sanitation body, friendsand relatives not wanting to visit and blocking of roads falls within the average condition, it therefore indicate that due to the presence of the refuse dump the area suffers the just listed health hazard in an average manner, that is to say, the occurrence are minimal though exist in that area. Furthermore, The Table 8 shows that only one falls within the very bad condition, this indicates that the presence of the refuse dump has made the neighbourhood to be repulsive to needed businesses. According to the popular Nigerian saying "The eyes eat before the mouth". Due to the unpleasant odour, unpleasant view and other environmental deterioration caused by the present of the refuse dump, this has drastically harper business activities with the affect area of the study area.

Recommendations

Based on the findings of this study, the following recommendations are hereby proposed;

There should be a total clearance of the existing illegal dump sites and proper monitoring of the waste management in the town to forestall illegal dumping. The offenders of this provision should be punished to serve as a deterrent to others. There should also adequate information to residents and awareness on the danger of consequences of indiscriminate dumping of refuse in an undesignated dump site. System of refuse collection by the responsible that is Niger agency Environmental Protection Agency should be improved upon to include door to door system of refuse collection, putting into consideration durable storage facilities.

Regular evacuation of the solid waste from dump site by the government and its agency. Prompt and timely evacuation of municipal solid waste from residence to the nearest landfill and sorting of time of decom recyclable was if there is a delain evacuation of

Tab	le 8: Resident
S/	Hazard
N	

1	Fear of disea
1	real of disea
2	Discomfort

- 3 Irritation from
- 4 Fear of people neighborhood
- 5 Fear of persent sanitation bod
- 6 The neighborneeded busine
- 7 Friends and my family
 8 Blocking of my

Conclusion

The study has that there is on dump sites in Sal its close proxim Nigeria and inf town. The quart shows that it kg/capita/day kg/capita/day.The dumping of so surrounding is method. There nformal waste the waste the town. T esidents to the refuse dump var 100m) and mild mber of building both environ azard is about

and sorting of the waste to reduce the time of decomposing and recycling of recyclable waste. It should be noted that if there is a delay or irregular frequency in evacuation of municipal solid waste,

it might lead to the temptation of residents returning back to their usual habit of indiscriminate refuse dumping.

Table 8: Residents' Level of Risk of Health Hazard Exposed to in the Study Area

S/	Hazard	Fixed Hazard Grade (1-5)					
N		Po	Fai	Avera	Ver	Extreme	Actua
		or	r	ge	y Bad	ly Dangero us	Grad e
1	Fear of disease	.03	.21	.82	1.56	1.01	3.62
2	Discomfort from foul Odour	.06	.18	.47	.88	2.40	3.99
3	Irritation from the sight of the dump	.07	.05	.64	1.51	1.53	3.80
4	Fear of people leaving in the neighborhood	.09	.42	.88	.94	.88	3.19
5	Fear of persecution by the environmental sanitation body	.07	.44	.42	.88	1.40	3.21
6	The neighborhood being repulsive to needed businesses	.08	.05	.35	1.38	2.14	4.01
7	Friends and relatives not wanting to visit my family	.07	.08	.55	1.32	1.88	3.90
8	Blocking of roads	.06	.25	.43	1.64	1.33	3.70

Conclusion

the

the

per i in

ng. uld

ate

ess of

an

of

ble

ate

1CY

ıde

use

ion

iste

and

ely

iste

fill

The study has been able to establish that there is one legal and 19 illegal dump sites in Sabon Wuse Area despite its close proximity to Federal Capital of Nigeria and influx of people to the town. The quantity of waste generated shows that it ranges between 0.52 kg/capita/day and kg/capita/day. The result also shows that dumping of solid waste into the surrounding is the most favoured method. There is also the use of informal waste collectors who also dump the waste in the illegal dump sites in the town. The perception of the residents to the risk of staying close to refuse dump varies between severe (at 100m) and mild (at 200m). The total number of buildings exposed to the risk of both environmental and health hazard is about 25.6% at 100 m and about 53.2% at 200 m buffer. This shows that a greater number of residents are exposed to danger on health and environmental hazard. This has an effect on the health care delivery as many residents are prone to mosquito attack and epidemic as a result of houseflies.

Refrences

Abd'Razack, N.T., and Bin Muhamadludin A.N. (2013): Environmental Assessment tool & its rating. A Review. Australian Journal of Basic and Applied Sciences.7(7), 20-78. Retrieved from www.researchgate.net, 2015.

Abd'Razack, Nelson T.A., Yusuf, A.E. and Utange, J.Z. (2013). An Appraisal of Solid Waste Generation and Management in

Univers

Jalingo City', Nigeria. Journal of Environment and Earth Science, 3(9), 20-29.

Agunwamba, J. C. (2003). Analysis of Scavengers' Activities and Recycling in Some Cities of Nigeria. *Journal of Environmental Management*. 32(1),116–127.Retrieved from www.link.springer.com..

Agwu, M. O. (2012). Issues and Challenges of Solid Waste Management Practices in Port-Harcourt City, Nigeria: A Behavioural Perspective. American Journal of Social and Management Sciences. 3(2), 83-92. Retrieved from www.scihub.org,.

Akinbile, C.O. and Yusoff, M.S. (2011). Environmental Impact of Leachate Pollution on Groundwater Supplies in Akure Nigeria. International Journal of Environmental Science and Development, 2, 81-89.

BBC News. (2005). Illegal Dumping 'Costs Millions'. Available online at http://news.bbc.co.uk/2/hi/uk_n ews/4310267.stm. [retrieved 1st April 2014].

BBC News. (2005). *Illegal dumping* 'costs millions'. Available: http://news.bbc.co.uk/2/hi/uk_n ews/4310267.stm. Accessed on 01 April 2014.

Bryman, A. (2001). Social Research Methods. UK: Oxford University Press.

Chung, S.S and Poon, C.S. (2001). A
Comparison of Waste Reduction
Practices and New
Environmental Paradigm of

Rural and Urban Chinese Citizen. Journal of Environmental Management 62(1), 3-19.

Etengeneng, D. (2012). Municipal Solid Waste Management in Grahamstown, Republic of South Africa. Novia: Novia University of Applied Sciences. A MSc Thesis. [pdf].

Holmes, D. (2000). Waste Minimization in the Small Islad Developing State of the South Pacific; Implication for Strategy & Policy. Workshop on Waste Management of Developing Countries. University of Wollongong, Australia.

Hoornweg, D. and Bhada-Tata, P. (2012). What a Waste: A Global Review of Solid Waste Management. Urban Development, 15, 22-32.

Johnson, K. (2006). Toxic Dumping in Africa Elicits calls for Better Controls. Available online at http://www.nationalgepgraphic.com/news/2006/10/061030-toxic-waste.html [Accessed on 9th Dec., 2015

Karanjit, H. G., Chowdhury, B., Alam T. R., Shrestha, F. S., and Hassan, U. (2007). Household waste collection – factors and variations. Luleå University of Technology, Sweden University Press, 12-34.

Kariuki, P. (2008). Examining
Consumers' Perceptions
Water Supply and Sanitation
Services: A case study of
Ohlange Township, Durban
South Africa. Durban

(MA the Kim, J.T. and Public

Environ Municipo Manager Countre Small in Manager

Countrie
Wollong
Lutui, V. (200
practices

attitudes Waste www.ro

Madava, T. (20)
Toxic War
Rights.
Political

Marshall, R.E. 2013. Sy integrated managem countries. 33, 988-10

Morrisson, R.S.. V &Dusbaur Workshop Manageme Countries

Wollongor
Tahman, A. and
Economic
Waste Ma
Africa:
Constraints
Conservant
521-531.

Agency
Profile

University of KwaZulu-Natal (MA thesis) [Pdf].

Kim, J.T. and Gobalan, H.N.B (1997): Public Health and Environmental Aspect of Municipal Solid Waste Management in Developing Countries with Special Focus on Small Island Developing State Workshop on Waste Management of Developing University Countries. Wollongong, Australia.

Lutui, V. (2001). Waste management practices, perceptions and attitudes in Tonga: Australian Waste Database. Cited in www.ro.uow.edu.au. 2015.

Madava, T. (2001). Illicit Dumping of Toxic Wastes breach of Human Rights. Review of African Political Economy, 28, 288-290.

Marshall, R.E. and Farahbakhsh, K. 2013. Systems approaches to integrated solid waste management in developing countries. *Waste Management*, 33, 988-1003.

Morrisson, R.S., Wray, R., Dever, S &Dusbaun, L. (2000). Workshop on Waste Management of Developing Countries. University of Wollongong, Australia.

Nahman, A. and Godfrey, L. (2010).

Economic Instruments for Solid
Waste Management in South
Africa: Opportunities and
Constraints. Resources,
Conservation and Recycling, 8,
521-531.

Niger State Environmental Protection Agency (NISEPA) (2012). Profile of Solid Waste Management in Niger State (2000-2012). Government of Niger State Press, Minna.

NPC (2015). 2006 National Population and Housing Census Results. Publication Units, National Population Commission. Presidency, Abuja, Nigeria.

Nwosu, A.E., and Olofa, S.A. (2015).

Effect of waste dump sites on proximate residential property values in Ibadan, Oyo State, Nigeria. Ethopian journal of Environmental studies and management, 8, 45-76.

Retrieved from www.ajol.info.com. 2015.

Ogwueleka, T. (2009). Municipal Solid Waste Characteristics and Management in Nigeria. Journal of Environmental Health, Science and Engineering. 6(3), 17-123. Retrieved from www.ijehse.tums.ac.ir. 2015.

Oyelola, O. T. and Babatunde, A.I (2008).Characterization domestic and market solid Wastes at source in Lagos metropolis, Lagos, Nigeria. African Journal of Environmental Science and Technology, 3, 430-437.

Project Green Sweep. (2011).

Environmental, Health and
Economic Effects of Illegal
Dumping. Available online at
http://pulse.pharmacy.arizona.ed
u/resources/EffectsofDumping.p
df [Accessed on 3rd/04/2016].

Sanusi, Y.A (2010). Water, sanitation and human development in urban fringe settlements in Nigeria. Theoretical and empirical researches in urban

- management. 5(7). Retrieved from www.search.proquest.com. 2015.
- UNEP (2005). Environmental Disaster and Development. *Journal of Environment Science*. 3(1), 2005.8-15.
- UNEP (2010): Solid Waste Management in Sub-Saharan Cities. UNEP publication Unit. Nairobi, Kenya.
- UNEPA (October 2014).

 Environmental Preparedness
 and Response. Geneva: press, 832.
- United **Nations** Environment Programme. (2005).Closing Dumpsite and Open Shifting from Open Dumping to Controlled Dumping and to Sanitary Landfilling. Available online http://www.unep.or.jp/ietc/Publi cations/spc/SPC Training-Module.pdf [Accessed 17th/08/2015].
- United States Environmental Protection Agency.(1998). Illegal Dumping Prevention Handbook.

 Available online at http://www.epa.gov/region5/waste/illegal_dumping/downloads/il-dmpng.pdf [Accessed on 21st/02/2016].
- USEPA (2005). National policy on Environmental risk reduction: Fact Sheet No. 134. Revised May 2008.
- Vika, L.(2001): Waste Management Practice, Perception & Attitude in Tonga. An Unpublished M.Sc Thesis. School of

- Geoscience. University of Wollongong, Australia.
- Wasley, A. (2011). UK e-waste illegally dumped in Ghana. The Guardian. Available online at http://www.theguardian.com/en_vironment/2011/may/16/uk-ewaste-dumped-ghana [Accessed on 23/07/2015].
- Waste Management Strategy in England and Wales (2000)

 Waste Management Strategies. London: square publisher, 5-27.
- World Bank (2010). Global Strategy for Health and Environment. World Bank/EHE/51.8.
- Zurbrügg, C. (2002). Urban Sohn Waste Management In Lou-Income Countries of Asia: How to cope with the Garbage Criss Urban Solid Waste Management Review Session. Durban, South Africa.

Develop: Resident

Hassan Shur Gbenga Mor Usman³

Department of Es Nigeria
Department of Bu Department of Es Corresponding aud

Abstract

Mass appraisal has taxation purpose. It countries as well a mass appraisal mooto analyse a set of evealed the contribusults, a mass appraisal the model was evaluated a good mass and a good mass and a suddy area by reduce ax assessment procephanatory variable keywords: Mass A

Introduction

he significance ociety and overemphasised.

The basic hum onstitutes a subsequent wealth of many forensic Technic Tech