

IMPACT OF HUMAN ACTIVITIES ON WATER QUALITIES AND SUSTAINABLE WATER DEVELOPMENTAL PROGRAMMES IN NIGERIA: A REVIEW

M. M. Ndamitso¹, S. O. E. Sadiku², A. I. Ajai^{1*}, Y. A. Iyaka¹.

1. Department of Chemistry, Federal University of Technology, Minna, Nigeria
2. Department of Fisheries Federal University of Technology, Minna, Nigeria.

Correspondence email: talk2alex2005@yahoo.com

Abstract

Water occupies the largest volume of space in the world, and serves as the most prominent sustainable factor for the existence of life. However, human activities have altered the qualities and quantities of most water bodies in Nigeria, in particular and the whole world in general. For instance, construction of dams has led to changes in the physical and chemical composition of the aquifer, which has consequently given rise to the growth of weeds that have rendered most water bodies unfit for navigation and agricultural activities. Industrial effluents, detergents, agricultural chemicals such as fertilizers and pesticides, as well as other surface run offs from industrial and farmland areas serve as ready reservoirs containing high concentrations of toxic elements and contaminants that could be harmful to life. This is a review of some of these activities of man that render such water bodies unfit for human use. This review has also shown that there is necessary need for government regulatory bodies and agencies on management of aquatic resources to make laws that will ensure safe water for consumption in Nigeria.

Key Words: Aquifer, effluents, navigation, aquatic.

INTRODUCTION

The most prized resource of any country in the world is water. This may not, however generally be realized until its uses are threatened by reduced availability or quality. Water occupies a prominent position among the people of the world (both rural and urban dwellers) and without it, life, in any form, is not possible (Lohani, et al., 1978). It may be said to have been the single most dominant factor in the settlements, civilizations and growths of ancient empires and kingdoms. For example, the Nile River provides the two basic necessities of human existence for Egypt and much of Sudan (Robert, 1988). Even in the modern ages, human settlements still tend to have been limited to those areas where water supply is relatively in abundance as it is not possible for man to survive where there is no water and to multiply without fertile lands that are adequately watered.

In most cases, the major quantities of water supply of these settlements come from the surface (oceans, rivers, prepared catchments, storage reservoirs and lakes) and underground (wells, natural springs, shallow wells, deep and artesian wells and horizontal galleries) sources (Kalmbach, 1982). The availabilities of water from these two sources vary with both season and the nature of the soil of a given region. Hence, depending upon the technological advancement of a given country, these two sources are heavily used for community water supply with the ground water lending itself more readily to smaller communities while the larger municipalities use surface water as their major source of water supply (Offodile, 2002).

The most important challenges of today's water resources management throughout the world are considered to be water scarcity, environmental degradation, economic development and ineffective administration. Sustainable

development in general offers a scientific, economic and social alternative approach that ensures the healthy coexistence of mankind and the environment. The questions arising on the use of water these days concern the prospect of sustainable water management as well as the shifts that the whole process will require at the environmental, economic, social, administrative and political levels. It is therefore a great challenge to us that to have sustainable water resources management, recovering of costs, policy integration, and participatory planning, there is need for an envisioning programme that will ensure long-term water supply with the right balance between the expansions of water supply projects and water conservation (Yannis, 2003).

Importance of both Surface and Ground Waters

Both surface and subsurface waters are the direct sources of our developed water resources (Glenn *et al.*, 1981). Surface waters exist in natural basins and stream channels. In areas where flows in streams or rivers are large in relation to water demands of adjacent lands, towns and cities, development of surface water is accomplished by direct withdrawal from flow. Reservoirs are however, constructed to hold flow during seasons or years of high run off and later release for beneficial use. In most cases, these reservoirs do incorporate hydroelectric, flood control and recreational features in addition to their water supply function. They may include such functions as irrigation and fisheries. These reservoirs range in size from several million hectares for large multipurpose reservoirs to small ponds. Good examples of large multipurpose reservoirs are Grand Coulee and Hoover Dams in the United States of America, which are 168 and 221m high with storage capacities of 0.74 and 3.46 million hectares respectively (Glenn *et al.*, 1981). Usually the subsurface water available for development

is referred to as ground water, which predominantly results from precipitation that has reached the zone of saturation in the earth through infiltration and percolation. While, groundwater is developed for use through well, springs and dugout ponds. However, in areas where groundwater is an important source of water supply it is being withdrawn much faster than being replenished from infiltration and percolation of precipitation which consequently results in ecological effect on the environment.

The Hydrology and Water Resources of Nigeria

Nigeria is a West African country covering an area of about 200,000 km² (Ademiluyi, 1988) with an estimated population of about 140 million people. It lies between longitudes 2° 49'E and 14° 37'E, and latitudes 4° 16'N and 13° 52' N of the equator. The climate is tropical, characterized by high temperatures and humidity as well as marked wet and dry seasons. The durations of these seasons however, vary from the South to the North with decrease in the rainfall from the coast northwards. The South has an annual rainfall ranging from 1,500 to 4,000mm, while the extreme North has between 500 and 1000mm annual rainfall (Ita, 1993).

The country is endowed with both surface and underground water with the latter being about 30 times the former (Ademiluyi, 1988). These water resources form a vast expanse of inland freshwater and brackish ecosystem although their full extent cannot be accurately stated as they vary with season and from year to year depending upon rainfall. These water resources are however, spread all over the country from the coastal region to the arid zone of Lake Chad basin (Ita, 1993).

Two great river systems, namely, the Niger- dominate the hydrology of Nigeria and with the exception of a few rivers that empty directly into the Atlantic Ocean (Cross River, Ogun, Osun, Imo, Kwa-Ibo and a few others), all other flowing waters finally find their way into the Chad Basin or down the lower Niger to the sea. Figure 1 shows the hydrological map of Nigeria with the major inland waters.

Benue and the Chad

The Niger-Benue and Chad systems are separated by a primary watershed which extends from North-East and North-West from the Bauchi Plateau-the main source of their principal tributaries (Ita, 1993; Ita and Sado 1984; Ita, et al, 1985) and to the North-West of the Plateau lie the drift covered plains of central Hausa land, drained by many streams flowing outwards to form the major tributaries.

The rivers that flow into Lake Chad come from Central highland and the high plateau both of which converge to

form the Komadugu-Yobe River just before flowing into Lake Chad (Offodile, 2002). Some of these rivers originate from the Cameroon Mountains. However, only a small part of Lake Chad lies within Nigeria which is not more than a mere vast swamp along the border. Most of the rivers flowing into this lake rise from the watershed areas of the Jos plateau and the Adamawa highlands and these include the Hadejia-Yobe, Alo and Yedseram from the southwest, and the Ba Mbassa, Chari, Illi and Lagone from the southeast. From the southeast to the northwest, the Rivers Yedseram-Ngadda together with all other minor tributaries and rivulets empty into the Lake Chad (Offodile, 2002). Within Nigeria, River Niger is fed by rivers flowing into it from all directions with head waters originating from the central plateau in the North, from the Yoruba highlands in the south, from Benin Republic to the west and from the eastern highlands. From outside Nigeria, we have a significant flow from the watersheds stretching westwards right up to the Fouta Djallon mountains of Guinea. The Benue River on the other hand, is fed by rivers emanating both from the high central plateau and also from the Cameroon mountains and Ogoja hills (Ita, 1993). Of the rivers flowing to the Atlantic Ocean, both the Ogun and Osun rivers are fed by rivers originating from the Yoruba highlands. They flow slowly from the north to south into the lagoons before discharging through creeks and swamps into the Atlantic Ocean. The Cross River on the other hand, is fed by many tributaries originating in the Cameroon Mountains. It flows east and then turns southwards and empties into the Atlantic Ocean with limited Delta formation. Those rivers flowing into the Atlantic Ocean in the south include; Imo, Kwa Iboe, Ase, Orashi, Benin and numerous creeks in the Delta and River states.

In general, the water bodies of Nigeria are divided into marine, brackish and fresh-waters (Ita, 1993) while the brackish waters with their saline wetland cover the surface area of 858,000 hectares, the freshwaters cover about 3,22,500 hectares. Other water bodies, including small reservoirs, fishponds and miscellaneous wetland suitable for rice cultivation cover about 4,108,000 hectares. The total surface area of the water bodies in Nigeria excluding marine and miscellaneous wetland suitable for rice cultivation is estimated to be about 14,991,900 hectares or 149,919 km² constituting about 15.9% of the total area of the country.

Human Activities that affect Hydrological Systems
Various human activities create many environmental problems ranging from global warming, biodiversity, ocean/water pollution, and ozone layer depletion to even total extinction of some species of plants and animals (Thomas, 2002). Among the activities that bring about

these deleterious effects are:

(i) **Construction of Dams for the generation of Electricity:** Water is a unique substance that forms a vital and integral part of man's life when put into many uses. These uses are however, affected by the presence of other substances (which could be from natural sources or as a result of human activities) in it. In different parts of the world, man made lakes and dams have been constructed for various purposes. For instance, the Volta Lake in Ghana, the Kainji and Jebba Dams in Nigeria constructed mainly for generation of electricity; the Nasser Dam constructed on the Nile for the regulation of water for the irrigation of Egypt. Others include the construction of seasonal reservoirs at Jabel Auliya (White Nile) and Sennar or Abbay (Blue Nile), Bakolori, Anka, Yomtera and Zobe all on the Sokoto System (Offodile, 2002 and Abdullahi, 2007) mainly for irrigation, while Lake Nwahdingusha and Koni on the River Lufira, and Nzilo on Lunlaba in Zaire are mainly for the supply of electricity (Beadle, 1981).

All these dams and Lakes have been constructed to improve the lots of humanity but they also have their own demerits especially in the tropics where they have adverse biological consequences (White, 1968), Principally, because construction of lakes and dams leads to the change of environment from riverine to lacustrine and consequent changes in the physical and chemical compositions of the former. Human activities in these environments have led to the growth of sudd forming weeds (e.g. *Vossia cuspidate*, *Pistia stratiotes*, and *Salvinia auriculata*). These usually render the waters unnavigable and inaccessible to other activities especially fishing. They also cause deoxygenation of these waters thereby making life unbearable to fish and other aquatic fauna. They also prevent the penetration of light energy into the waters making photosynthesis very difficult for the phytoplankton. These weeds, having formed mats, can also be blown towards the dam, which can eventually block the turbines. The decayed vegetation in these waters can also cause the accumulation of hydrogen sulphide (Ewer, 1966). The economic importance of the growth of water hyacinth (*Eichhornia crassipes*) on these water bodies cannot however, be over-emphasized. These weeds, having been established, are not easily eradicated and their eradication always involve huge economic losses. For instance, in Sudan it is stated that it costs the government up to 500,000 pounds each year to clear the waterways of label Auliya in as far back as the sixties (Hassan and Heinen, 1964).

The newly created environments also serve as conducive habitats for the aquatic vectors of certain important

diseases plaguing man and domestic animals such as bilharzias and liver fluke (snails); malaria and filariasis (mosquitoes). Eventually, water pollution problems due to rural-urban migration are also observed in these areas. For instance, water pollution caused as a result of effluent discharge into these water bodies usually lead to the overloading of the marine environment as a result of certain chemicals and heat that have such deleterious effects act as harm to marine activities including fishing, impairment of quality for use and reduction of amenities (GESAMP, 1980; Deju, 1971; Richard, 1992).

Seasonal alternation between inundation and desiccation of considerable areas of land as a result of opening of the sluice gates in order to make room for the incoming floodwater in rainy season (annual draw down) also affects the ecology of these areas. In addition, a large number of people are displaced along the downstream of these dams like Kainji, Shiroro and Jebba Dam in which thousands of people are displaced every year. The socio-economic problems created by these displacements are enormous especially as some large agricultural lands are either washed away or drowned tectonic equilibrium is also offset by water bodies especially in regions of tectonic instability.

(ii) **Agriculture:** Agriculture is another human activity, which affects both surface and underground water quality. In agricultural practices, the use of chemical fertilizers, herbicides, pesticides and fungicides is employed to improve the output by farmers. The clearing of agricultural lands and eventual burning of the cleared vegetation, lead to the emission of high quantities of undesirable polyatomic gases like carbon dioxide, water vapour and oxides of nitrogen which reduces the earth's infra-red radiation with a consequent rise in the warming up of the earth's surface and the atmosphere. This ultimately leads to the destabilization of the earth's natural global climatic regime with a gradual but steady rise in the global temperature of between 1.5°C and 4.5°C and finally the rising trend in temperatures, evapotranspiration and the intermittent droughts as well as extension of the deserts most especially in the Sahel region (Offodile, 2002). Most of the herbicides and pesticides in common use in this country are harmful to the environment and some of them have been banned by some countries that are critical about their environmental degradation. For instance, fenthion, an insecticide, is effective also as an avicide and therefore widely used in many states in Northern Nigeria like Kano, Bauchi, Yobe and Borno which are prone to bird pests. DOT and gammalin, which have rodenticide properties although outlawed, are still being used illegally in some parts of the

country. These insecticides are classified as moderately to highly hazardous by World Health Organization and as harmful to extremely harmful to aquatic life (Ita, 1993).

(iii) **Industrial:** Paradoxically, while industrialization in most of our cities and towns have propelled the economic growths of these centers by creating wealth and employment opportunities for the youths, it has done a lot of damage to the environment. In these areas, fossil fuels are flared and toxic gases like sulphur dioxide and nitrogen dioxide are emitted to the environment and appreciable amounts of these substances (e.g. sulphur), build up in the soil and subsurface waters. Acidification of the soil and soil water also occurs (Michael and Frank, 1991). Unfortunately, despite the number of industries in this country that discharge these substances of varying environmental toxicity, less than 10% of them manage the waste in environmentally friendly manner (Louis, 2001). In most cases, these wastes are dumped in any available space on water or land. The lagoons, streams, and rivers which are the final recipients of these wastes and other surface runoffs are incidentally the sources of drinking water, fishing, livestock rearing, washing and recreation for a large proportion of the populace despite the fact that they serve as ready eutrophic reservoirs collecting high levels of toxic heavy metals like the case of the Jankara reservoir in Kano city (Butt, 1985; Adeniji and Mbagwu, 1990). When these lakes, rivers and aquifers become loaded with high concentrations of these chemicals and heat, they are no longer suitable for consumption nor can they support fish and wild life (Deju, 1971). Furthermore, the soils within the premises of these industries as well as the underground water are heavily contaminated with these hazardous wastes posing high risks to consumers of crops. Water from wells of such soils like the case of the Kurmin Gwari, a suburb of Kakuri, in Kaduna City has been rendered a danger zone for human habitation as well as agricultural activities (Thomas, 2002). The overloading of the Kaduna River with industrial wastes by the Kaduna textiles which have been reported to be discharging between 27 million and 35 million litres of untreated water daily and the large volume of waste discharged into the fragile coastal and marine environment from Rivers to Lagoons is estimated to be about 313,000 tons per annum. These are other examples of the gruesome impacts of human activities on the fragile environment (Louis, 2001; Thomas, 2002).

(iv) **Oil and Gas Explorations:** The same sad tales are told in the oil and gas sector with oil spills and release of noxious gases into the atmosphere accompanying crude oil production. A large amount of gases is produced following the production of a single barrel of crude oil (about 1,000 standard cubic feet per barrel or about two

billion standard cubic feet per day) but only less than 30% of this is being converted to useful fuel with the remaining being flared. In fact it has been estimated that Nigeria released 30 million tons of carbon dioxide into the environment in 1989, a value which rose to 35 million in 1994, thereby placing the country as contributing about 28% of the world's total gas flared and hence a significant contributor to the emission of greenhouse into the atmosphere and the acid rain phenomenon in the Niger Delta (Louis, 2001).

Deforestation: While many countries are in the bid of checking global warming which eventually affects the hydrology of the world by making efforts to conserve the forests, falling of trees is still unabated in this country due to the fact that many Nigerians are poorly served by electric energy with only about 34% of the country's households having access to this essential commodity. Nearly three-quarters of the country's households are still dependent on firewood for their energy.

On the domestic scene, whenever the raining season sets in Nigeria, residents of most of the major cities in the country, including our young and prestigious new Federal Capital-Abuja, are gripped by the reality of the annual charade of flood resulting mostly from lack of drainages or their blockage due to improper refuse disposal. As a result, surface runoffs and other sewage waters are directed into place of greater importance making the sites to stink, and a common example is the Abuja Monday Market (Maimuna, 2004). Our major roads can also be over flooded and submerged by this phenomenon as experienced in the case of Zone five, and some parts of Areas 2, 8 and 11 all in the Capital City of this great country (Yusuf, 2004). The use of automobiles which run on leaded gasoline resulting in increase of lead in both surface and ground waters of our big cities and towns due to vehicular emissions. The use of detergents in laundry can also lead to increase in the phosphate levels of our waters (Ita, 1993).

Mining: Mining activities also affect the potability of waters most especially the ground water by raising the dissolved organic matter, acidity and corrosive properties. A typical example of this is the pollution of Aboh and Onyeama coal field waters (in Enugu) where the samples have high temperatures of 30- 40°C and considerable quantities of hydrogen sulphide (Offodile, 2002).

Suggestions: In the past, lack of care by man for his environment, though devastating in some areas, could apparently be solved by abandonment and immigration. Today, with rapid population growth and accumulated degradation, the easy option of just finding another source for our needs is no longer easy to come by (John and Michael, 1988). Consequently, solutions have to be

sought since it is not morally sound for us to leave our problems for the coming generations to solve while we enjoy the benefits today (UNESCO, 1977; Zylicz, 2002). Furthermore, of all the natural resources that are in abundance, only water seems to be in the least abundance when potability is taken into consideration. Hence, despite the fact that water occupies the largest volume of space in the whole world only 0.62% of the total body is available for potability (Wilson, 1975).

In the industrially advanced countries, it is normal to provide at least some degree of treatment for water from any source and the public expect standard water supply services, efficient collection, treatment and disposal of waste-waters. This is because they have well-developed techniques and equipment needed to attain these goals. In rural schemes in developing countries, e.g. Nigeria treatment will not be feasible in many circumstances since in most cases, the facilities used to obtain potable water in these areas, if at all provided, are fashioned in accordance with the sophisticated developed-country style. These facilities can only be proficiently handled by the donor governments and agencies. Consequently, little value is derived from these facilities since no appropriate operation and maintenance back up are provided (Tebbut, 1983). In such areas, it is only necessary for us to consider water resources in relation to what is likely to be the most important parameter (bacteriological quality). This is because, local communities, especially those located down streams of urban sewers that do not have adequate provisions for their water treatment are highly prone to severe bacteriological water borne diseases like typhoid, cholera and dysentery (White, 1986; Terence, 1991).

Apart from adequate provisions by the Water Boards of every state of the federation for routine chemical, physiological and biological checks. Care should also be taken to ensure that the ecological upset caused by these pollutions is adequately measured to ascertain the severity of the pollutions. Also, since some sewage contain high organic manure useful for agricultural activities and for the enhancement of aquatic life which usually lure the rural and even some city dwellers into their direct use, despite the fact that they contain some toxic substances, which are detrimental to both aquatic and human lives. Hence, every effort should be made to dissuade these people from the use of such bodies for agricultural and other practices prior to proper treatment.

Traditionally, agricultural activities in river basins are known to have little effect on water quality and may even be regarded as a non-competing joint use of the resource.

With the increasing human population and the need for more crop production, agricultural methods which are more advanced, requiring the retention of water in large dams for use in sophisticated irrigation systems have been adopted in various parts of these country. These have resulted in desiccation of downstream farmlands to the detriment of traditional agriculture and loss of fishery resources as a result of habitat loss (Welcomme, 1993). In addition, the use of inorganic fertilizers that accompany these aggressive agricultural methods, coupled with build up of salts in the soil, can lead to a sudden excessive growth of phytoplankton in stagnant ponds and pools when they are washed into other water bodies. Consequently, excessive bloom of these algae can lead to high biological oxygen demands resulting in aquatic pollution. In order to avoid these situations, it is suggested that water organizations of this country in conjunction with other non governmental organizations (NGOs) saddled with the responsibilities of providing wholesome water for communities be adequately funded to be able to fill all these ponds and also provide adequate sewage treatment plants for our big cities and towns. In rural areas where the achievement of the above goal may not be easy to come by, the use of hardy species of fish that can tolerate such polluted waters like Clarias and Tilapia could be of help to some extent (Ita, 1993). In order to prevent high health hazards, it is necessary to have and also enforce laws by state and Federal Ministries of Health prohibiting the catching and hence, consumption of fish from such polluted reservoirs and ponds.

We also, are aware of the fact that the intensive agricultural activities embarked upon in the Northern parts of the country not only require the application of inorganic fertilizers but also, the application of pesticides for increased yield. These pesticides used in this country include certain chemicals that for environmental reasons have been partially or completely banned in developed countries (FEPA, 1991). These chemicals still continue to find their ways into Nigeria for pest control because the effective cheap substitutes have not yet been evolved and have been found to cause serious environmental problems, especially in the dry season when the dilution capacity of water systems is very low leading to an increase in the risk of high concentration of toxic chemicals with severe consequences (Ita, 1993). In light of these problems above, it is suggested that the government ban the use of these hazardous pesticides and herbicides and the more environmentally friendly substitutes provided for the farmers. It is also suggested that in light of criticisms being leveled

against the nation's environmental laws including those of the water resources decree number 101 of 1993 (Offodile, 2002) which are still viewed as being ineffective in ensuring sustainable expectation and management of the nation's environmental codes to make the best practice standard in the world (Nnimmo, 2004), water and the Federal environmental agencies be given a kind of leadership that has seen the like of Prof. (Mrs.)Dorothy Akunyeli of NAFDAC to eliminate the ineptitude that bedevils the implementation of these codes (Louis, 2001).

Furthermore, although surface water tends to be the major source of water supply to towns and cities, the prevailing advance and unpredictable climatic conditions have made the need for a thorough study and development of underground water a necessity in Nigeria. This however has to be done with great care in order not to destroy some of the country's most important ground water resources (Offodile, 2002).

In general, the management of solid waste is so complex that this task should not be vested solely on the local government authorities but all the three arms of government should fully participate (Louis, 2001).

In order to curb the menace of greenhouse effect as a result of the release of large amount of noxious gases flared into the atmosphere by the oil sector, the Federal Government has set up the Nigerian Liquefied Natural Gas (LNG) in

Bonny and also harness the vast potential to generate further foreign income for the economy. This plant was originally planned to be in full operation by the year 2004. However, we still experience the flaring of gases with hope that in the nearest future, this act will be banned. Finally, it is important that the government curtails the spate of indiscriminate land use abuses particularly land reclamation in the urban areas. This is because; if this should continue unabated more wet lands will be filled for residential purposes which will force the water down streams into rivers, seas or oceans. This will eventually lead to increased water volume and consequent ocean surge and coastal flooding. As a result, the residential and commercial buildings within such areas will be at the danger of being washed away causing a great water pollution problem.




Fig.1. Hydrological map of Nigeria showing the major inland waters.

REFERENCES

- Abdullahi, M. G. 2007. Ethiopia: Echoes of the Queen of Sheba. Daily Trust, Friday May 16th, No. 49, pp 28 and 29.
- Ademiluyi, J. O. 1988. Regional Profiles of Selected Groundwater Quality Measures in Nigeria. *Water International, Official Journal of the International Water Resources Association*, 13: 130 - 135.
- Adeniji, H.A.; Mbagwu, I.G. 1990. Study of physico-chemical characteristics of some heavy metals in Jankara Reservoir, Kano State, Nigeria, NIFFR Annual Report, pp 136 - 140.
- Beadle, L.C. 1981. *The Inland Waters of Tropical Africa* (2nd Edn), Longman Inc., New York.
- Butt, I.A. 1985. Report on the industrial pollution of Jankara Dam Ministry of Health, Kano, Nigeria.
- Deju, R. A. 1971. *Regional Hydrology Fundamentals*. Published by Gordon and Breach, Science Publishers, Inc. 440 Park Avenue South New York, N.Y. 10016.
- Ewer, D.W. 1966. Biological Investigations on the Volta Lake, May 1964 to May 1965. In *Man-made Lakes*, Ed. R.M. Lowe McConnell, Academic Press, London.
- FEPA, 1991. Guide lines and standards for environmental pollution control in Nigeria. Federal Environmental Protection Agency (FEPA), Nigeria.
- Gbenm, O. S.; Richard, K.F.; Takatt, W.E; Kenneth, K.B. 1981. *Soil and Water Conservation Engineering*, Third Edition.
- GESAMP. 1980 Report of the eleventh Session. Reports and Studies No. 10. UNEP. [GESAMP IMO/FAO/UNESCO/IMO/WHO/IAEA/JUNI/UNEP. [Joint Group of Experts on the Scientific Aspects of Marine Pollution].
- Hassan, A.; Heinen, E T. 1964. *Water hyacinth Control on the Nile River*. Sudan Department of Agriculture, Khartoum.
- Ita, E.O. 1993. *The Inland Fishery Resources of Nigeria*. CIFA Occasional Paper No. 20.
- Ita, E. O.; Sado, E K. 1984. Inventory survey of Nigerian inland waters and their fishery resources with special reference to ponds, lakes, reservoirs and major rivers: surface area survey and potential yield estimate. Kainji Lake Research Institute 1983 Annual Report.
- Ita, E. O.; Sado, E.K.; Pandogari, A.; Balogun J. K.,; Ibitoye, B. 1985. Inventory Survey of Nigerian inland waters and their fishery resources 1. A preliminary checklist of inland water bodies in Nigeria with special reference to ponds, reservoirs, and major rivers. Kainji Lake Research Institute Technical Report Series.
- Kalmbach, O. 1982. *Encyclopedia Britannica (Macropaedia) Volume 19, (15th Edn)*. Hartmolls, Bodmin, Cornwall.
- Lewis, I. 2001. Pollution, Waste Management, Desertification, Erosion Flood Control. The Nigeria Environmental Score Sheet. The PUNCH, Thursday, January 11, pp 29
- Maimuna, Y. 2004. Abuja Monday Market Traders decry stinking refuse. A DAILY TRUST article on Property and Environment. Wednesday, April 14, pp 26.
- Offodile, M. E. 2002. *Groundwater study and development in Nigeria*. Published by Mecon Geology and Engineering Services. Jos, Nigeria.
- Robert, O. C. 1988. The Jonglei Canal: Illusion or Reality? *Water International. The Official Journal of the Int. Water Res. Ass.* 13(3):144-153.
- Tebbutt, T.H.Y. 1983. *Principles of Water Quality Control*. (3rd Edn). Pergamon Press
- Terence, J.M. 1991. *McGraw-Hili Series in Water Resources and Environmental Engineering* (6th Edn). Published by McGraw-Hili Publishing Company.
- Thomas, A. 2002. Industrial Pollution: Agonies of Kaduna South residents. *New Nigerian Weekly*, July 13, pp 4.
- Welcomme, R.L. 1983. *River basins F AO Fisheries Technical paper* (202).
- White, E 1968. The Place of Biological Research in the Development of the Reservoirs of Man-made Lakes, in *Man-made Lakes, the Accra Symposium* Ed L.E. Obeng, Ghana UP., Accra.
- White, J.B. 1986. *Waste Water Engineering, (International Student Edition)* Printed and bound by Arnold Allion Printing Co. Ltd. Hong Kong.
- Wilson, D.C. 1975 *Encyclopedia Britannica*.
- Yannis, M. 2003. Urban Water Management in Greece, Present Conditions and Perspectives of Sustainability, *Water International*. 28: 43-51.
- Yusuf, O. U. 2004. Environmental Rights Action (ERA) and friends of the Earth Nigeria (FOEN). A DAILY TRUST article on Property and Environment. Wednesday, April 14, pp 26.
- Zylicz, T.J. 2002. The role of Science in sustainable development policy. Retrieved September 2003, from URL: <http://www.sustainability.at/easv/easv/ecollpdr/zylicz/paper.pdf>

International Journal of Chemical Sciences is a six monthly & peer-reviewed scientific journal published by Department of Chemistry, Nasarawa State University, Keffi, Nigeria. The journal is devoted to the publication of contributions in all fields of chemistry, including chemistry of natural products, new materials, macromolecules and polymers, industrial chemistry, physical chemistry, inorganic chemistry, agrochemistry, petrochemistry, geochemistry, clinical chemistry, biological chemistry, analytical and environmental chemistry, nanotechnology and surface chemistry.

We would be honoured if you could join this new and exciting project by submitting your sound and current manuscripts through cdnsukijcs@gmail.com for consideration. Your article will undoubtedly be an extremely valuable contribution to the Science Community. You may wish to request for "Guide for Authors" by contacting us;

The Managing Editor
Dr. Leke Aremu
Department of Chemistry,
Nasarawa State University,
PMB 1022, Keffi, Nigeria.

The Editor-in-Chief
Dr. E. O. Ojeka
Department of Chemistry,
Nasarawa State University,
PMB 1022, Keffi, Nigeria.

Director of Publication
Dr. A. Olonisakin
Department of Chemistry,
Nasarawa State University,
PMB 1022, Keffi, Nigeria.

CENTRE FOR MULTIDISCIPLINARY RESEARCH AND ACTION (CMRA)

P.B. No. 55, Thodupuzha, Kerala 685 584, India.

Trivandrum Office: TC 12/304. PLAMOOD, PATTOM, TRIVANDRUM -4.

Director: **Dr. Stephen Joseph**

Registrar: **Mr. Salvy Thomas**

Dr. Ravindran Ankathil
Dr. G. S. Gireesh Kumar

Dr. Jomy Augustine
Prof. Joseph Augustine

Dr. T. M. Joseph

Centre for Multidisciplinary Research and Action (CMRA), is a registered Autonomous Research Institute with Offices at Thodupuzha and Thiruvananthapuram. CMRA envisages launching of programmes helpful for fostering and promoting Research activities in a multidisciplinary approach focusing emphasis on problems that confront contemporary issues relevant to the society at large. The Quarterly International Journals of CMRA are **INDIAN JOURNAL OF MULTIDISCIPLINARY RESEARCH** and **INDIAN JOURNAL OF BOTANICAL RESEARCH**. All articles published in these Journals have been reviewed by the referees and are deemed to reflect the opinions of authors and not of the CMRA. Please send your research papers (3 copies and CD) for publication to the Managing Editor in the Thodupuzha address.

Dr. Stephen, Managing Editor, CMRA, P. B. No. 55, Thodupuzha, Kerala-584, INDIA.
E-mail: drstephen@sanchamnet.in; drstephenjoseph@gmail.com