

## EFFECT OF LAND CONFIGURATION AND CROP RESIDUE – MULCH ON MILLET PERFORMANCE IN MINNA, SOUTHERN GUINEA SAVANNA ZONE OF NIGERIA

Eze, P. C<sup>1</sup>., Odofin, A. J<sup>1</sup>., Attahiru, A<sup>1</sup>., Onyekwere, I. N<sup>2</sup>. and Lawal, B. A<sup>1</sup>.

<sup>1</sup>Department of Soil Science and Land Management, Federal University of Technology, Minna, Niger State.

<sup>2</sup>National Roots and Tuber Crops Research Institute, Umudike, Umuahia, Abia State.

### ABSTRACT

A 2 x 3 factorial experiment was conducted during the 2012 cropping season at the old Teaching and Research Farm, Federal University of Technology, Minna, to determine the effect of land configuration and crop residue – mulch on the growth and yield of millet in minna, Southern guinea savanna zone of Nigeria. It was a Randomized Complete Block Design with two types of land configuration (ridge and flat surface) and three mulch application rates (0, 10 and 15 t/ha), replicated three times. Composite soil samples were collected from the experimental site at 0 – 15 and 15 – 30 cm depths prior to the commencement of this study, for the determination of initial soil physicochemical properties. Crop growth parameters measured were plant height, number of leaves per plant, number of tillers per plant and leaf area index at 3, 6, 9 and 12 weeks after planting. Yield indices determined were panicle length and stover yield. Data collected were subjected to analysis of variance at 0.05 level of significance, while means separation was done using Duncan's multiple range test. Findings in this study showed that planting on ridge resulted in taller ( $P \leq 0.05$ ) millet plants and higher number of leaves per plant, higher number of tillers per plant and higher leaf area index than planting on the flat. Plant height, number of leaves per plant, number of tillers per plant, leaf area index, panicle length and stover yield increased with increasing mulch application rates.

**Key words:** land configuration, crop residue – mulch, soil moisture content, millet, growth and yield parameters.

### INTRODUCTION

Major constraints to high productivity of most soils of the savanna zone of Nigeria are low organic matter content, low nutrient status, poor structural stability, low water holding capacity and extremely high permeability (Yandev and Sachan, 1985). Modifying soil surface condition of these soils in combination with application of organic residues as amendments have been found to improve their physical and chemical properties, with a consequent improvement in crop growth and higher crop yields (Chiroma *et*

*al.*, 2006a &b). Khan (2002), reported that alteration of soil surface condition plays a significant role in preventing loss of water through evaporation and in making available water beneficial for crop growth. Modification of soil surface could be in the form of ridge tillage with different configurations such as open-ridge, tied-ridge and a combination of the two. It could also be in the form of furrows, flat beds, raised beds, mounds, mulching and crop residue incorporation.

Ridges, due to their height, are exposed to

more sunlight and exhibit higher temperature than furrows and flat soil surface. Radke (1982) reported that ridges had higher temperature than other forms of land configuration. Growth of crop is positively influenced by temperature, especially within certain limit. It has been documented that conservation tillage practices such as ridge tillage, lower soil bulk density and reduce penetration resistance (Materechera and Mioza-Banda, 1997; Sow *et al.*, 1997). This results in promoting root elongation and improving crop growth and yield. Organic residues could be chopped into smaller bits or as whole plant parts and then applied on the soil surface as mulch or incorporated into the soil. These technologies are gradually gaining popularity due to rapid degradation of agricultural lands in recent times, and the need to increase and sustain the productivity of arable lands. Researchers that made use of grain-cereal straw as crop residue-mulch noted that tillage practices that retain crop residues on the soil surface enhance the yield potential of crops by conserving moisture (Alhassan *et al.*, 1998; Chiroma, 1993) and improving soil fertility status (Batiano *et al.*, 2000; Biolders *et al.*, 2002). Rice husk is available in huge quantities around rice mills where it is set on fire because it is not a desirable livestock feed. Therefore, it faces less competition unlike so many other crop residues such as maize, rice and wheat straws. Little or no attention has been given to this huge waste generated and its impact on the environment. A more serious issue is that, there is no deliberate attempt to effectively utilize the rice husk, or at least, to safely dispose it.

Millet (*Pennisetum glaucum* (L.) R.Br.) is commonly grown on 5.2 million hectares for its grains, with an average yield of 9 t/ha (FAO and ICRISAT, 1996; Gupta, *et al.*, 2000). It is cultivated under rain-fed condition in Northern

Nigeria mainly because of its drought-resistant characteristics (Eze, 2015). It is a major staple food usually grown as sole crop or intercropped with legumes such as cowpea and groundnuts. However, the yield of millet has continued to decline to as low as between 5000 – 8000 kg/ha due mainly to low soil fertility status, declining annual rainfall, insect pests and diseases, striga, birds and low-yielding varieties (Kwari and Babinu, 2002). The objective of this research was to determine the effect of land configuration and crop residue-mulch on the growth and yield of millet in Minna.

## MATERIALS AND METHODS

### Site Description

The research was conducted in 2012 at the old Teaching and Research Farm, Federal University of Technology, Minna under rain-fed condition. The experimental site is located at longitude 9° 3'E and latitude 6° 27'N, and 240.50 m elevation above mean sea level. Minna is found within the Southern guinea savanna zone of Nigeria, which is characterized by alternate wet and dry seasons. Mean annual rainfall ranges between 1,000 and 1,200 mm. Rainfall commences between late April and early May, and ends between late September and early October. Minimum temperature ranges from 18 – 22 °C, while mean maximum temperature is 40 °C between February and March, and 36 °C between November and December. Soils around Minna are derived from basement complex rocks. They range from sandy clay to loamy sands. The vegetation is characterized by vast grassland interspersed with few tree species and shrubs. Most of the arable crops (sorghum, millet, yams, rice, cowpea, maize and groundnuts) are usually grown under rain-fed conditions. Also, vegeta-

bles such as tomatoes, okra, pepper, spinach and cocchorus are grown under irrigation during the dry season.

### Treatments and Experimental Design

The experiment was a 2 x 3 factorial in laid out in a Randomized Complete Block Design with three replications. The treatment combinations were six in number, consisting of two land configuration (ridge and flat) and three mulch application rates (0, 10 and 15-t/ha).

### Soil Sampling and Analysis

Prior to the commencement of the trial, soil samples were collected randomly at 5 m interval within the experimental site at two depth intervals (0 – 15 and 15 – 30 cm) using soil auger. The samples were bulked to obtain a composite soil sample, air-dried, gently crushed and passed through a 2 mm sieve for physicochemical analysis in accordance with standard procedures. The Bouyoucos hydrometer method (Gee and Bauder, 1986) was used for the determination of particle size distribution. Textural class of the soil was determined using a textural triangle. Soil pH was determined using glass-electrode pH meter. The pH was determined on a 1:2.5 soil : solution ratio in both water and 0.01M  $\text{CaCl}_2$  solution (Mc Lean, 1982). Total nitrogen was determined by employing the modified macro-Kjeldahl method (Bremner and Mulvaney, 1982). The organic carbon content of the soil was determined by Walkley - Black (1934) wet - oxidation method. Available phosphorus extraction was carried out by employing the Bray No.1 method (Bray and Kurtz, 1945). Exchangeable bases ( $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{K}^+$  and  $\text{Na}^+$ ) were extracted with 1N neutral ammonium acetate ( $\text{NH}_4\text{OAc}$ ) solution (Grant, 1982).  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  were determined by titration, while  $\text{K}^+$  and  $\text{Na}^+$  were measured by flame photometry (Grant,

1982). Exchangeable  $\text{H}^+$  and  $\text{Al}^{3+}$  were extracted with 1N KCl and the extract titrated with 0.05 N NaOH. The total exchangeable acidity was estimated by summing the amounts of  $\text{H}^+$  and  $\text{Al}^{3+}$  determined (Grant, 1982). Cation exchange capacity (CEC) was determined by summation of the total exchangeable bases and total exchangeable acidity. Percent base saturation was computed as follows:

$$\% \text{Base-Saturation} = \frac{\text{Exchangeable bases, cmolkg}^{-1}}{\text{Exchangeable cations, cmolkg}^{-1}} \times 100$$

### Cultural Practices

The land was cleared, tractor-ploughed and levelled manually. The experimental plots were laid out. Fifteen (15) cm high ridges were constructed manually, 0.75 m apart, using a hand-hoe. The crop residue-mulch (rice husk) was uniformly applied on the appropriate experimental plots shortly before planting. Five to six Apron-plus-treated seeds of millet (LCIC 9702 variety) were hand-sown per hole at a spacing of 75 x 50 cm. The gross and net plots dimensions were 6 x 3 m and 4.5 x 2 m, respectively. Seedlings were thinned to two plants per stand, two weeks after planting. NPK fertilizer (15:15:15) was applied in two split doses, at the recommended rate (60:30:32 kg/ha) at 3 and 7 weeks after planting. Weeding was carried out manually using a hand-hoe as at when due. Harvesting of the stover was done at physiological maturity, 12 weeks after planting. The grains were destroyed by weaver birds.

### Measurement of Crop Growth Indices

The crop growth indices determined at 3, 6, 9 and 12 weeks after planting were plant height, number of leaves, number of tillers and leaf area index. Ten plants were tagged within the net plot for these measurements. Plant height was

measured from the base of the plant to the highest flag leaf using a metre-rule and a calibrated wooden pole. The number of leaves and tillers were counted on each of the tagged plants. Leaf area index was determined following the modified length and breadth product method by Dugje and Odo (2006):

$$LAI = (P \times L \times A) / GA$$

where, P is plant population per hectare, L is number of fully expanded green leaves per plant, A is single leaf area (cm<sup>2</sup>) and GA represents ground area (hectare).

### Determination of Crop Yield Indices

Crop yield indices determined at physiological maturity were stover yield and panicle length. The above-ground plant part (stover) was harvested 12 weeks after planting and left in the field for about two weeks to allow for thorough sun-drying, and weighed on a Salter balance. The length of panicles was measured from the base to the tip of the panicles using a metre-rule.

### Statistical Analysis

Data collected were subjected to analysis of variance at 0.05 level of probability using Statistix 8.0 (2005) statistical software. Where significant differences exist between means, Duncan's Multiple Range Test was employed for the purpose of means separation.

## RESULTS AND DISCUSSION

### Soil Physico-chemical Properties before Cultivation

Results in Table 1 show the physicochemical properties of the soil of the experimental site. The results indicate that the soil is sandy loam in texture. According to the fertility ratings for soil data by FDALR (1985) and FAO (2004),

the soil was neutral in reaction, low in organic carbon, total nitrogen, exchangeable magnesium, potassium and cation exchange capacity. It is very low in available phosphorus and exchangeable calcium, and moderate in organic matter, C:N ratio, sodium and percent base saturation. The sandy nature of this soil and its low fertility status necessitates the application of soil amendments such organic residues to improve soil physical condition, soil nutrient status and enhance its productivity (Chiroma *et al.*, 2006a & b).

### Crop Growth Indices

#### *Plant Height, Number of Leaves per Plant, Number of Tillers per Plant and Leaf Area Index*

The crop growth indices studied were plant height, number of leaves per plant, number of tillers per plant and leaf area index at 3, 6, 9 and 12 weeks after planting (WAP). The results of the effect of land configuration and crop residue-mulch on plant height and number of leaves per plant are presented in Table 2. Planting on ridge produced significantly taller ( $P \leq 0.05$ ) plants than planting on the flat, at 9 and 12 WAP. Also, plant height increased with increase in mulch application rate throughout the growing season. A similar trend was observed in which planting on ridge and increase in mulch rate resulted in significantly higher ( $P \leq 0.05$ ) number of leaves per plant. Interaction between land configuration and mulch had significant influence on plant height at 6 WAP. Table 3 indicated that combination of ridge treatment and 15 t/ha mulch produced the tallest millet plants. This attests to the superiority of planting on ridge and 15 t/ha mulch application rate compared to planting on the flat and lower mulch application rates

Results of the effect of land configuration and crop residue-mulch on number of tillers per

plant and leaf area index are presented in Table 4. Ridge treatment produced higher number of tillers per plant than planting on the flat at 9 and 12 WAP. Fifteen t/ha mulch resulted in the highest number of tillers per plant at 3, 6, 9 and 12 WAP, followed by 10 t/ha rate. The lowest in number was produced when no mulch was applied. Similarly, planting on ridge resulted in higher leaf area index than planting on the flat at 12 WAP, while leaf area index increased with

increase in mulch application rate. Interaction between land configuration and crop residue-mulch had no significant influence on number of tillers per plant and leaf area index.

In summary, ridge tillage resulted in taller millet plants, higher number of Leaves per plant, more tillers per plant and higher leaf area index than raising plants on the flat. The better performance of plants grown on ridges than those raised on the flat is well supported by the find-

**Table 1: Soil physicochemical properties of the experimental site at Minna.**

Parameters	0 – 15	15 – 30
Depth, cm		
Sand, g kg <sup>-1</sup>	745	655
Silt, g kg <sup>-1</sup>	103	153
Clay, g kg <sup>-1</sup>	152	192
Textural Class	Sandy loam	Sandy loam
pH (1:2.5) in Water	6.70	6.90
pH (1:2.5) in 0.01M CaCl <sub>2</sub>	5.70	5.60
Organic Carbon, g kg <sup>-1</sup>	7.70	7.40
Total Nitrogen, g kg <sup>-1</sup>	7.40	7.00
C : N Ratio	1.04	1.06
Available Phosphorus, mg kg <sup>-1</sup>	0.85	1.26
Exchangeable Calcium, cmol kg <sup>-1</sup>	0.49	0.46
Exch. Magnesium, cmol kg <sup>-1</sup>	0.51	0.49
Exch. Potassium, cmol kg <sup>-1</sup>	0.28	0.25
Exch. Sodium, cmol kg <sup>-1</sup>	0.52	0.50
Exch. Acidity (H <sup>+</sup> + Al <sup>3+</sup> ), cmol kg <sup>-1</sup>	1.10	1.20
Cation Exch. Capacity, cmol kg <sup>-1</sup>	2.90	2.90
Base Saturation, %	62.10	58.60

Table 2: Effects of Land configuration and mulch on plant height and number of leaves per plant of millet at Minna.

Treatment	Plant height (cm)				Number of leaves/plant			
	3 WAP	6 WAP	9 WAP	12 WAP	3 WAP	6 WAP	9 WAP	12 WAP
A: Land configuration								
Ridge	24.50a	110.60a	157.40a	171.50a	6.6a	6.8a	6.0a	4.9a
Flat	21.90a	93.60b	114.90b	136.20b	5.7b	5.9b	5.9a	4.2b
SE ±	1.9	5.0	6.2	5.9	0.2	0.2	0.3	0.1
B: Mulch rate								
0 t/ha	18.90b	72.20c	116.70b	122.20c	5.4b	5.6b	4.9b	4.2b
10 t/ha	23.90ab	102.10b	131.60b	151.90b	6.2a	6.5a	6.0ab	4.7ab
15 t/ha	26.80a	132.00a	160.20a	187.50a	6.8a	7.0a	6.9a	4.8a
SE ±	2.3	6.1	7.6	7.2	0.2	0.2	0.4	0.1
Interaction								
A x B	NS	*	NS	NS	NS	NS	NS	NS

Means with the same letter (s) in the columns are not significantly different according to Duncan's Multiple Range Test (DMRT) at 5 % level of probability.

NS: Not Significant; \*: Significant at 5 % level of probability

WAP: Weeks After Planting

ings of other researchers. Materechera and Mioza-Banda (1997) and Sow *et al.* (1997) reported that improvement in crop growth resulted from decrease in soil bulk density and reduction in penetration resistance to root elongation. Ridge is a conservation tillage practice which raises the seed bed above the land surface, gathers the top fertile soil layer and conserve moisture on soils of Nigerian savanna characterized by low nutrient status (Lal, 1995; Hatfield *et al.*, 1998; Chiroma *et al.*, 2006a). Haque *et al.* (2002) also noted that tallest plants, highest number of leaves per plant and higher garlic bulb diameter were produced on ridges than on furrows and flat planting methods.

The increase in growth parameters with increase in mulch application rate is adduced to better soil surface cover under the highest mulch treatment (Akambi and Ojeniyi, 2007; Ojeniyi *et al.*, 2009). Another reason could be improved nutrient status following the application of 15 t/ha mulch. Eze *et al.* (2014a & 2015b), reported that higher soil organic carbon and total nitrogen contents resulted from 15 t/ha mulch rate. This produced taller plants than 0 and 10 t/ha mulch treatments. A third reason for the production of the tallest plants as a result of application of 15 t/ha mulch could be reduction in evaporative moisture loss or lower crop evapotranspiration with increase in mulch application rate (Eze *et al.*, 2014a; Eze, 2015).

Table 3: Interaction effects of land configuration x mulch on plant height (cm) of millet at 6 weeks after planting at Minna.

Mulch rate	Land configuration	
	Ridge	Flat
0 t/ha	69.70b	74.80b
10 t/ha	103.20b	101.00b
15 t/ha	158.90a	105.10b
SE ±	8.8	

Means with the same letter (s) in the rows and columns are not significantly different according to Duncan's Multiple Range Test (DMRT) at 5 % level of probability.

**Table 4: Effects of Land configuration and mulch on number of tillers and leaf area index of millet at Minna.**

Treatment	Number of tillers/plant				Leaf area index			
	3 WAP	6 WAP	9 WAP	12 WAP	3 WAP	6 WAP	9 WAP	12 WAP
<b>A: Land configuration</b>								
Ridge	2.2a	2.3a	2.7a	3.2a	0.14a	0.39a	0.52a	0.68a
Flat	2.0a	2.1a	2.2b	2.4b	0.11a	0.31a	0.48a	0.56b
SE ±	0.1	0.1	0.1	0.2	0.01	0.03	0.04	0.02
<b>B: Mulch rate</b>								
0 t/ha	2.0b	2.1b	2.2c	2.5b	0.07b	0.17c	0.31b	0.41c
10 t/ha	2.0b	2.1b	2.5b	2.6ab	0.13a	0.37b	0.52a	0.61b
15 t/ha	2.2a	2.4a	2.8a	3.3a	0.17a	0.52a	0.67a	0.84a
SE ±	0.1	0.1	0.1	0.2	0.02	0.03	0.05	0.03
<b>Interaction</b>								
A x B	NS	NS	NS	NS	NS	NS	NS	NS

Means with the same letter (s) in the columns are not significantly different according to Duncan's Multiple Range Test (DMRT) at 5 % level of probability.

NS: Not Significant.

WAP: Weeks After Planting

**Table 5: Effects of land configuration and mulch on panicle length and stover yield of millet at Minna.**

Treatment	Panicle length (cm)	Stover yield (kg/ha)
<b>A: Land configuration</b>		
Ridge	20.40a	3,432.10a
Flat	19.00a	3,333.30a
SE ±	0.80	275.10
<b>B: Mulch rate</b>		
0 t/ha	17.6b	2,222.20b
10 t/ha	19.80ab	3,277.80b
15 t/ha	21.60a	4,648.10a
SE ±	0.90	337.40
<b>Interaction</b>		
A x B	NS	NS

Means with the same letter (s) in the columns are not significantly different according to Duncan's Multiple Range Test (DMRT) at 5 % level of probability.

## Crop Yield Indices

### Panicle Length and Stover Yield

Table 5 displays the results of the effect of land configuration and crop residue-mulch on panicle length and stover yield. Although, land

configuration did not influence the two parameters significantly, they increased with increase in mulch rate. Fifteen t/ha mulch rate produced longer panicles and higher stover yield than both 0 and 10 t/ha rates. This observation could be attributed to lower crop evapotranspiration and

better microclimate resulting from higher soil surface cover and higher leaf canopy cover under 15 t/ha mulch treatment (Eze *et al.*, 2014b). Other reasons could be higher soil nutrient content. In a similar study, higher soil organic carbon and total nitrogen contents that resulted from the highest mulch application rate (15 t/ha) were found to be responsible for longest panicles and highest sorghum stover and grain yields (Eze *et al.*, 2015b).

## CONCLUSION

Findings in this study showed that planting on ridge resulted in taller ( $P \leq 0.05$ ) millet plants and higher number of leaves per plant, higher number of tillers per plant and higher leaf area index than planting on the flat. Plant height, number of leaves per plant, number of tillers per plant, leaf area index, panicle length and stover yield increased with increasing mulch application rate.

## REFERENCES

- Akanbi, O. S. and Ojeniyi, S. O. (2007). Effect of siam weed mulch on soil properties and performance of yam in Southwest, Nigeria. *Nigerian Journal of Soil Science*, 17, 120-125.
- Alhassan, A. B., Kundiri, A.M. and Folorunso, O. A. (1998). Water use efficiency of millet as influenced by different tillage and residue management practices in semi-arid Northeast Nigeria. *Journal of Arid Agriculture*, 8, 45-51.
- Bationo, A., Wani, S., Biielders, C. L., Vlek, P. L. G. and Mokwunye, A. U. (2000). Crop residue and fertilizer management to improve soil organic carbon content, soil quality and productivity in the desert margins of West Africa. In: Lar, R., Kimbe, J. M. and Stewart, B. A. (eds). *Global climate change and tropical systems*. CRC S. Press, Boca Raton, pp. 117-245.
- Biielders, C. L., Michels, K. and Bationo, A. (2002). On-farm evaluation of ridging and residue management options in a Sahelian millet-cowpea intercrop. 1. Soil quality changes. *Soil Use and Management*, 18, 216-222.
- Bray, R. H. and Kurtz, L. T. (1945). Determination of total, organic and available forms of P in soils. In: Page, L. A., Miller, R.H., and Keeney, D. R. (Eds) (1982). *Methods of Soil Analysis, Part 2. Chemical and Microbiological Properties* (2nd Edition), Agronomy monograph No. 9. Madison, Wisconsin, USA: American Society of Agronomy.
- Bremner, J. M. and Mulvaney, C. S. (1982). Total N in Soils. In: Page, L. A., Miller, R.H., and Keeney, D. R. (Eds) (1982). *Methods of Soil Analysis, Part 2. Chemical and Microbiological properties* (2nd Edition), Agronomy monograph No. 9. Madison, Wisconsin, USA: American Society of Agronomy.
- Chiroma, A. M. (1993). Effects of crop residue management and tillage on root growth, yield and water use efficiency of millet on a sandy loam soil. *M.Sc. Thesis*, University of Maiduguri, Nigeria.
- Chiroma, A. M., Folorunso, O. A. and Alhassan, A. B. (2006a). The effects of land configuration and wood-shavings mulch on the properties of a sandy loam soil in North-east Nigeria. 1. Changes in chemical properties. *Tropicultura*, 24(3), 129-134.
- Chiroma, A. M., Folorunso, O. A. and Alhassan, A. B. (2006b). The effects of land configuration and wood-shavings mulch on the properties of a sandy loam soil in North-east



- Nigeria. 2. Changes in physical properties. *Tropicultura*, 24(1), 33-38.
- Dugje, I. J. and Odo, P. E. (2006). Effects of planting pattern and variety on leaf area index and grain yield of Pearl millet intercropped with groundnut. *Nigerian Journal of Experimental and Applied Biology*, 7(1), 63-69.
- Eze, P. C. (2015). Effects of rice bran-mulch on soil properties and water use efficiency of sorghum and millet in Maiduguri, North-east Nigeria. *Unpublished Ph.D. Thesis*, University of Maiduguri.
- Eze, P. C., Alhassan, A. B., Tsado, P. A., Onyekwere, I. N., Kwari, J. D. and Dada, Y. M. (2014a). Effects of rice bran-mulch on the growth of sorghum and millet in Maiduguri, Semi-arid North-east Nigeria. *Nigerian Journal of Soil Science*, 24(1), 59-71.
- Eze, P. C., Alhassan, A. B., Kwari, J. D., Dada, Y. M., Tsado, P. A. and Lawan, A. N. (2014b). Effects of rice bran-mulch on the growth of sorghum and millet in Maiduguri, semi-arid north-east Nigeria. *Nigerian Journal of Soil Science*, 24(1), 72-79.
- Eze, P. C., Kwari, J. D., Kundiri, A. M., Tsado, P. A., Onyekwere, I. N. and Dada, Y. M. (2015b). Effects of rice husk-mulch on soil chemical properties under sorghum and millet in Maiduguri, Nigeria. *Production Agriculture and Technology Journal*, 11(1), 22-31.
- (FAO) (2004). *Soil fertility Maps of Nigeria*. National Special Programme for Food Security, Food and Agricultural Organization of the United Nations, Rome.
- FAO & ICRISAT (1996). The World sorghum and millet economics facts, trends and outlook. Food and Agricultural Organization of the United Nations, Rome and International Crops Research Institute for the Semi-arid Tropics, Patancheru, India.
- FDALR (1985). *The Reconnaissance Soil Survey of Niger State (1:250,000)*. Soils report. Federal Department of Agricultural Land Resources.
- Gee, G. W. and Bauder, J. W. (1986). Particle size analysis. In: Klute, A. (Ed) (1986), *Methods of Soil Analysis, Part 1. Physical and Mineralogical Properties*. Agronomy Monograph No. 9. Madison, Wisconsin: American Society of Agronomy.
- Grant, W. T. (1982). Exchangeable cations and exchangeable acidity. In: Page, L. A., Miller, R.H., and Keeney, D. R. (Eds) (1982). *Methods of Soil Analysis, Part 2. Chemical and Microbiological Properties* (2nd Edition), Agronomy monograph No. 9. Madison, Wisconsin, USA: American Society of Agronomy.
- Gupta, S. C., Ezeaku, J. E., Gwadabe, L. and Balarabe, S. (2000). Performance of pearl millet variety LC-ICMVI in a pearl millet-cowpea based system in Nigeria. *International Sorghum and Millet Newsletter*, 121pp.
- Hatfield, J. L., Allmara, R. R., Rehm, G. W. and Lowery, B. (1998). Ridge-tillage for corn and soybean production: Environmental quality impacts. *Soil and Tillage Research*, 48, 145-154.
- Haque, M. S., Sattar, M. A. and Pramanik, R. H. M. (2002). Land configuration and varietal effects on yield contributing traits and yield of garlic. *Pakistan Journal of Biological Sciences*, 5(10), 1024-1027.
- Khan, A. R. (2002). Mulching effect on soil physical properties and peanut production. *Italian Journal of Agronomy*, 6, 113-118.

- Kwari, J. D. and Bibinu, A. T. S. (2002). Response of two millet cultivars to sub-optimal rates of NPK fertilizer and sheep manure in different agroecological zones of North-east Nigeria. *Nigerian Journal of Soil Research*, 3, 33-38.
- Lal, R. (1995). Tillage and mulching effects on maize yield for seventeen consecutive seasons in a tropical Alfisol. *Journal of Sustainable Agriculture*, 5, 79-93.
- Materechera, S. A. and Mioza-Banda, H. R. (1997). Soil penetration resistance, root growth and yield of maize as influenced by tillage system on ridges in Malawi. *Soil and Tillage Research*, 41, 13-24.
- Mc Lean, E. O. (1982). Soil pH and lime requirements. In: Page, L. A., Miller, R.H., and Keeney, D. R. (Eds) (1982). *Methods of Soil Analysis, Part 2. Chemical and Microbiological Properties* (2nd Edition), Agronomy monograph No. 9. Madison, Wisconsin, USA: American Society of Agronomy.
- Ojeniyi, S. O., Odedina, S. A., Odedina, J. N. and Akinola, M. O. (2009). Effect of tillage and mulch combination on soil physical properties and sorghum performance on alfisol of Southwest Nigeria. *Nigerian Journal of Soil Science*, 19(2), 16-20.
- Radke, J. K. (1982). Managing early season soil temperatures in the northern corn belt using configured soil surfaces and mulches. *Soil Science Society of America Journal*, 46, 1067-1071.
- Sow, A. A., Hossner, L. R., Unger, P. W. and Stewart, B. A. (1997). Tillage and residue effects on root growth and yields of grain sorghum following wheat. *Soil and Tillage Research*, 44, 121-129.
- Statistix 8.0 (2005). *STATISTIX for windows* (version 8). Analytical software. Retrieved from support@statistix.com.
- Walkley, A. and Black, A. I. (1934). Carbon and organic matter. In: Page, L. A., Miller, R.H., and Keeney, D. R. (Eds) (1982). *Methods of Soil Analysis, Part 2. Chemical and Microbiological Properties* (2nd Edition), Agronomy monograph No. 9. Madison, Wisconsin, USA: American Society of Agronomy.
- Yandev, O. P. and Sachan, R. S. (1985). Some practical soil conservation measures in drought prone areas of Nigeria. *Noma*, 5, 4-7.