

Environmental Technology & Science Journal

Vol. 12 Number 1

June 2021

Aim and Scope

The Environmental Technology and Science Journal (ETSJ) is devoted to the publication of papers which advance knowledge of practical and theoretical issues that daily plague our society. The aim of the journal is to provide an avenue for the dissemination of academic research findings from various disciplines of the environment, engineering, pure and applied sciences, arts and social science which have materials that emphasize on environmental issues.

ETSJ Policy

ETSJ prohibits an author from submitting the same manuscript for concurrent consideration in two or more publications. The author must ensure that when a manuscript is submitted to ETSJ, the manuscript must be an original work. The Author should check the manuscript for any possible plagiarism using any program such as TURNITIN or any other software before submitting the manuscripts to ETSJ. Authors are not permitted to add or remove any names from the authorship provided at the time of initial submission without the consent of the Journal's Editor-in-Chief.

Guide for Authors

Please read the guidelines and follow the instructions carefully; doing so will ensure that the publication of your manuscript is as rapid and efficient as possible. The Editorial Board reserves the right to return manuscripts without review that are not prepared in accordance with these guidelines.

1. Manuscripts should not be more than 15 pages of A4 paper size (including abstracts, tables, figures, references and appendices) typed in double spacing, times new roman and font size 12. Additional fee of two thousand (₦2000) Naira will be charged for every additional page in excess of 15 pages.
2. Each paper must be compiled in one column; all text should be left and right justified. The page settings to be 25cm (1 inch) each for the top, bottom, left and right margins of an A4 paper. Decimal numbering of all sections of the article is recommended (with the exception of the reference and acknowledgement sections).
3. To ensure anonymity in the peer reviewing process, articles must be structured in such a way that the title of the paper, authors' names, address, and affiliation are written on the first page, followed by abstract and the body of the paper in that order.
4. Each article should contain an abstract of not more than 300 words.
5. Each article should contain between 5 and 7 key words (avoid use of phrases).
6. Use the SI (*Système Internationale*) units and symbols, while Unusual Characters and symbols should be explained in a list of nomenclatures.
7. The journal's referencing style is the APA format.
8. Figures, Tables and Graphs are to be placed at the appropriate places in the paper and not at the end of the paper.
9. Acknowledgement is optional, except for researches being funded through research grant awards.
10. Authors should avoid using headers, footers and special page breaks within the manuscripts.
11. The manuscript is to be sent via electronic means to: etsj@futminna.edu.ng for a blind peer review, which under normal circumstance will not exceed 4 weeks, after which the status of the article will be communicated to the Author(s).
12. Author(s) with positive review will be asked to effect corrections or revisions after which the camera-ready manuscript is to be emailed to etsj@futminna.edu.ng and to be accompanied by evidence of payment of publication fee.

13. The Editorial board is not responsible for the information or views expressed by the author(s).
14. The copyright of this journal belongs exclusively to the Editorial Board of Environmental Technology and Science Journal. No part of the publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise without prior permission of the School of Environmental Technology, Federal University of Technology, Minna-Nigeria or a licence permitting restricted copying.

Frequency of Publication

The journal is published twice a year in June and December

Subscription Details

The 2021 subscription rates for hardcopies of the journal including postage are:

Individual within Nigeria: N2,000 per copy and N1,000 for postage

Individual outside Nigeria including postage \$ 100

Institution within Nigeria: N5,000 per copy and N1,000 for postage

Institution outside Nigeria including postage: \$ 150

Correspondence

All correspondence should be addressed to

The Managing Editor

Environmental Technology & Science Journal

SET, FUT, Minna, Nigeria

Email: etsj@futminna.edu.ng

Phone: +234 805 170 3663, +234 803 653 4507

Published By

School of Environmental Technology, Federal University of Technology,
Minna-Nigeria

Managing Editor

Prof R. A. Jimoh

Department of Building,
Federal University of Technology,
Minna, Nigeria

Editorial Secretary I

Dr B. O. Ganiyu

Department of Quantity Surveying,
Federal University of Technology,
Minna, Nigeria
+234 803 626 9235

Editorial Secretary II

Dr E. B. Ogunbode

Department of Building,
Federal University of Technology,
Minna, Nigeria
+234 806 328 6122

Editorial Committee

Dr A. D. Adamu

Department of Quantity Surveying,
Federal University of Technology,
Minna, Nigeria

Prof I. C. Onuigbo

Department of Surveying &
Geoinformatics,
Federal University of Technology,
Minna, Nigeria

Prof O. A. Kemiki

Department of Estate Management
and Valuation,
Federal University of Technology,
Minna, Nigeria

Dr P. Ayuba

Department of Architecture,
Federal University of Technology,
Minna, Nigeria

Dr J. E. Idiako

Department of Quantity Surveying,
Federal University of Technology,
Minna, Nigeria

Dr A. Kawu

Department of Urban & Regional
Planning,
Federal University of Technology,
Minna, Nigeria

Dr I. B. Muhammad

Department of Architecture,
Federal University of Technology,
Minna, Nigeria

Editorial Advisers

Prof R. E. Olagunju

Department of Architecture,
Federal University of Technology,
Minna, Nigeria

Prof (Mrs) S. N. Zubairu

Department of Architecture,
Federal University of Technology,
Minna, Nigeria

Prof A. M. Junaid

Department of Urban & Regional
Planning,
Federal University of Technology,
Minna, Nigeria

Prof Y. A. Sanusi

Department of Urban & Regional
Planning,
Federal University of Technology,
Minna, Nigeria

Prof D. A. Muazu

Department of Building,
Baze University,
Abuja, Nigeria

Prof I. O. Aje

Department of Quantity Surveying,
Federal University of Technology,
Akure, Nigeria

Prof B. T. Aluko

Department of Estate Management,
Obafemi Awolowo University,
Ile-Ife, Nigeria

Prof P. C. Nwilo

Department of Surveying &
Geoinformatics,
University of Lagos,
Lagos, Nigeria

Prof U. O. Nkwogu

Department of Architecture,
Imo State University,
Owerri, Nigeria

Editorial

Two things have so far defined 2021 in giving hope to the populace; the 1st one is the Earth Day which was celebrated on 22nd April, 2021 while the 2nd one was the International Day of Co-operatives that was celebrated on 3rd July, 2021. According to the EarthDay.org President – Kathleen Rogers, “at the heart of Earth Day’s 2021 theme, Restore Our Earth, is optimism, a critically needed sentiment in a world ravaged by both climate change and the pandemic”. In expressing this sentiment, we have to “rebuild better together” which was the theme of this year’s International Day of Co-operatives. At the centre is the environment which continues to suffer one degradation or the other. It is on this basis that I welcome our readers to Volume 12, number 1, June edition of Environmental Technology and Science Journal (ETSJ). This edition is work-in-progress; articles will be added as they are reviewed and accepted. There are currently eight (8) articles that are published in the 1st instance.

The 1st article by Haruna *et al.* assessed the air quality of solid waste dumps in residential neighbourhoods of Makurdi Town. The result showed the mean concentration of carbon monoxide (CO) gas (16.0ppm), H₂S (M = 10.51ppm), and CH₄ (M = 12.17ppm) recorded in Wadata, and North Bank: CO (M=12.16ppm), H₂S (M = 4.17ppm), and CH₄ (M = 4.16ppm) recorded were above the regulatory standards recommended which if not checked will portend serious public health issues for the occupants of the areas studied.

Raheem *et al.* in the 2nd article also assessed protected green space of Eleyele Dam, Ibadan. The study concluded by recommending that further granting of approval on the land be stalled, non-compatible activities that could further lead to depletion of the green space should be discouraged and there should be thorough monitoring of the activities of people within the area.

According to Markus and Duniya in the 3rd article, researches have revealed that the fully-enclosed courtyard can enhance thermal performance provided that the appropriate Window-to-Wall-Ratio (WWR) is determined. To determine the WWR, the window-area and wall-area of the architectural space are required. Therefore, their study investigated the types of Window-to-Wall-Ratios (WWR) by identifying the types of window-area and room-area in fully-enclosed courtyard buildings in Kafanchan. The results indicated five types of window-areas, three types of room-areas and five types of WWRs. WWR3 has the highest percentage (39.1%), while WWR1 has the lowest percentage of (6%). The findings suggested that the larger WWRs are being used than the smaller ones.

The compressive strength and the durability of concrete containing an agricultural waste - *Acacia Nilotica* Seed Powder (ANSP) in 0.25%, 0.50%, 0.75% and 1% addition to the concrete mix design was evaluated by Dadu *et al.* in the 4th article. Results indicated that the summation of the major oxides of SiO₂ (27.7%), Al₂O₃ (4.04%) and Fe₂O₃ (6.21%) was about 38% indicating a Class C pozzolan. The study therefore recommended that 0.25% of ANSP can be added in concrete production as a retarder since it delays final setting time for about 8 hours; similarly increases the compressive strength of the concrete and improved the water absorption capacity from 2.4 to 7.4%.

The 5th paper by Yusuf *et al.* established the relationship between crime incidence and residential neighbourhood attributes towards improving safety and security in Kano Municipal Local Government Area (KMLGA). Findings from the multiple regression analysis for assault crime revealed that vacant lot/building has the strongest coefficient of determination of 0.140 and for theft crime, dead-ends-street has the strongest (0.360) as unique predictors that explained assault and theft crimes when the variance explained by other predictors in the model is controlled. The

study therefore recommended that there should be improvement programmes such as the provision for street lighting, pathway expansions, efficient waste removal, routine maintenance and repair of street lighting equipment in areas of high crime.

Communication enables exchange of ideas and instruction from one person to another. In the construction industry, effective communication among project participants helps to improve the level of understanding in order to achieve project objectives. Hence, Abdullahi *et al.* in the 6th paper assessed the communication practices adopted for managing construction projects in Abuja. Key findings of the study indicated that drawings were the most used mode of communication. Language was the most frequently stated as a barrier to effective communication by all the stakeholders. Furthermore, the site operatives were the most difficult people to communicate with during project construction phase. The study exposed the need for effective communication among stakeholders in the construction for achieving project goals.

Ganiyu *et al.* in the 7th paper assessed the organisational and behavioural factors influencing the retention of knowledge in Quantity surveying firms in Abuja with a view to establishing the relationship between the behavioural and organisational factors that influence knowledge retention. The results of analysis revealed that workers' low turnover is enhanced by proper motivation, effective trainings, good working environment, improved and effective organisational culture. Therefore, the study concludes that improvement of organisation values, beliefs and standards tend to improve workers behaviour towards achieving the goals of the organisation which encourages proper retention of the knowledge of effective workers which helps to ensure continuity in the flow of knowledge in organisations. The study recommended that employees should be properly trained by knowledgeable workers approaching retirement in order to ensure

the continuity of knowledge flow in the organisation.

Despite the importance of groundwater, quality assessment of groundwater has received little attention. Therefore, Mukhtar in the 8th paper assessed the physicochemical qualities of drinking water in Nguru town, Nguru LGA, Yobe State where eight (8) samples were collected from boreholes using purposive sampling technique. The result of the student t-test showed that all the parameters were within the acceptable drinking limit specified by WHO and NSDWQ with all p-values less than the t-critical of 2.14. The study concluded that the water is suitable for drinking.

R. A. Jimoh
Managing Editor

Contents

- 1-6 Air Quality Assessment of Solid Waste Dumps in Residential Neighbourhoods of Makurdi Town
Haruna D. M, Onoja O. O & Santali B. N
- 7-13 Assessment of Protected Green Space of Eleyele Dam, Ibadan
Raheem W. M, Adeboyejo A. T & Abolade O
- 14-20 Determination of Window-To-Wall Ratio of Fully-Enclosed Courtyard Residential Buildings in Kafanchan
Markus, B. & Duniya, S. D.
- 21-28 Effect of *Acacia Nilotica* Seed Powder on the Properties of Concrete
Dadu, D. W. Ibrahim, A. G. & Kida, A.T.
- 29-37 Urban Crime Incidence and Prevention Mechanisms in Kano Municipal Area Council
Yusuf, R, Adamu, H. I, Babanyara, Y. Y, Yahaya, A. U & Kawu, A. M
- 38-55 Assessment of Communication Practices adopted for Managing Construction Projects in Abuja
Abdullahi, A. H., Ibrahim, K., Momoh, A. S., Salawu, A., Ganiyu B. O., Kajimo-Shakantu K & Jimoh R. A
- 56-62 Organisational and Behavioural indices for knowledge Retention in Quantity Surveying Firms in Abuja Nigeria
Ganiyu B. O, Oyewobi L. O, Abdullahi L & Umar M. K
- 63-70 Assessment of Groundwater Quality and its Suitability for Drinking in Nguru
Mukhtar S, Usman N & Obadaki Y. Y

Air Quality Assessment of Solid Waste Dumps in Residential Neighbourhoods of Makurdi Town

*Musa H. D, Onoja O. O & Santali B. N
Department of Urban and Regional Planning,
Federal University of Technology Minna, Nigeria
musaharunad@futminna.edu.ng

Received: 9/4/2021

Revised: 14/5/2021

Accepted: 30/5/2021

The environs of urban centres in most developing countries are facing a serious depreciation in terms of quality of life caused by inadequate or poor solid waste management systems. Municipal solid waste management has been a challenging task for many cities. This study, therefore, assesses the effect of disposed neighbourhood solid waste on the air quality in Makurdi town. MSA Altair 5X Multi-gas Detector manufactured by Mine Safety Appliance Company USA was employed to measure the concentration of ambient air in the dumpsite's environment in the neighbourhoods. The ambient air quality measured were compared with regulatory standards of the Federal Environmental Protection Agency for Environmental Pollution Control in Nigeria which stipulated emission limits of pollutants from stationary sources such as from a site, process, stack and vent. Findings reveal poor ambient air quality in waste dumps due to the presence of concentrations of potentially harmful pollutants around solid waste dumpsites in Wadata and North Bank areas among the five selected neighbourhoods. The result shows the mean concentration of carbon monoxide (CO) gas (16.0ppm), H₂S (M = 10.51ppm), and CH₄ (M = 12.17ppm) recorded in Wadata, and North Bank: CO (M = 12.16ppm), H₂S (M = 4.17ppm), and CH₄ (M = 4.16ppm) recorded were above the regulatory standards recommended.

Keywords: Air quality; solid waste management; Quality of life; Pollution

INTRODUCTION

Globally, the effective management of urban solid waste has been a challenging task among developing countries, and in Africa where there are significant difficulties in urban solid waste collection and disposal (Maria, Gois & Leitao, 2019). The environs of urban centres in most developing countries are facing a serious depreciation in terms of quality of life caused by inadequate or poor solid waste management systems (Chen, 2018). The dearth of quality urban waste management framework and practices irregularities at the local level are the identifiable significant factors (Verma, Kaur & Tripathi, 2020).

The all-inclusive eco-friendly urban waste management system has become a global challenge due to rapid urbanisation, population increase, limited resources, growing industrialization and changes in the standard of living in urban centres (Ramachandra *et al.*, 2018). Rapid population explosion and urbanization has worsened the urban condition by producing a massive amount of solid waste at a rate that exceedingly surpassing that of urban growth (Ying-Chu, 2018). It is reported that two billion tons of generated municipal solid waste were recorded in 2011 worldwide (Amoo & Faglenle, 2013), and by 2050 it's projected to exceed 9.5 billion tons (Chen and Lo, (2015). The tremendous increase in the volumes of municipal solid waste led to mismanagement by local authorities (Verma, Kaur & Tripathi, 2020). Raised in the living standard of the urban inhabitants has

changed the pattern of goods consumed, which is directly proportional to the volume of solid waste generation (Chen, 2018; Verma *et al.*, 2020). Municipal solid waste generated encompasses all mixed wastes from urban, peri-urban regions, and the waste consists of waste discarded predominantly of household wastes with a minor portion of commercial, institutions, and industrial wastes (Chen, 2018).

Solid waste generation is an incessantly growing issue at the local scale and a challenging task for many cities. Studies revealed that the solid waste disposed is a source of greenhouse gases in an urban area. The municipal solid waste sector contributes 4% of the global total greenhouse gas emission (Metz, 2015). Greenhouse gases include carbon dioxide (CO₂), methane (CH₄), and volatile organic compounds (VOCs) etc. According to Stocker (2014), methane has a greater potent for global warming potential than carbon dioxide (CO₂) among all the greenhouse gases, and also, increase annual atmospheric concentration by 2% (Chen, 2018). In all the anthropogenic sources for the greenhouse gases, methane emission from landfill accounted for 3–9% in the world (Du *et al.*, 2017). Furthermore, the United Nation Framework Convention on Climate Change (UNFCCC), the Paris Agreement (PA2105) signed by all 197 members in 2018 target to reduce global warming by holding the increase in global average temperature to below 2° Celsius by effectively control the emission of greenhouse gases into the atmosphere (European Commission, 2019).

Nigeria has witness growth in population and economic development occasioned by mineral resources exploration and exploitation. As a result, cities are expanding in size and the populations accompanied by an increase in waste generation on a weekly or daily basis (Eche *et al.*, 2015). Rapid urbanization with the increasing urban population, changing lifestyle of the people and as well as an increase rate of development has led to a proliferation in the generation of solid waste in Makurdi town (Aguoru, 2015). Makurdi urban area is rapidly urbanizing and fast become characterized by indiscriminate refuse disposal, an uncleared heap of refuse that threatens the health of the residents (Eche *et al.*, 2015). This study, therefore, assesses the effect of disposed neighbourhood solid waste on the air quality in Makurdi town.

LITERATURE REVIEW

Municipal Solid Waste and Greenhouse Gas Emission

Greenhouse (GHG) gas are atmospheric gases such as water vapour, carbon dioxide, methane, and nitrous oxide that aid the absorptions and emissions of radiation within the thermal infrared range. Greenhouse gases, shield the earth surface from heat (LeTreat *et al.*, 2007). A naturally occurring shield of greenhouse gases comprising 1-2 percent of the earth's atmosphere, helps to absorb the radiated solar energy from the space and minimise the heat impact on the earth to a comfortable and liveable range (LeTreat *et al.*, 2007).

Human activities over the years have produced 40% increase in the atmospheric carbon emission concentration (Chen *et al.*, 2018), and has further increased despite the large uptake by various natural sinks. Anthropogenic carbon dioxide (CO₂) emissions source in the atmosphere includes fossil fuel combustion, natural gas, soil erosion, deforestation, wastes and livestock agriculture (Mora, 2013). A continuous increased in the emission of greenhouse gas on earth surface at the present rate may explicitly imply increased in earth temperature, with possibly harmful effects on environments, biodiversity and the livelihoods of people globally (Mora, 2013). The most abundant greenhouse gases in the earth's atmosphere are - Ammonia (NH₃), Carbon dioxide (CO₂), Methane (CH₄), Nitrogen dioxide (NO₂), Sulphur dioxide (SO₂), Carbon monoxide (CO), Hydrogen sulphide (H₂S) and Particulate matter.

Diverse wastes and management options have different implications for energy consumption, methane emissions, and carbon sequestration. The municipal landfills air pollutants emitted contributes to the greenhouse gases emission in the atmosphere

which causes severe harms to human health (Lazaridis & Chalvatzaki, 2010). Niloufer, Swamy, and Syamala (2014) assessed ambient air quality in solid waste dumpsites at Vijayawada, and Andhra Pradesh in India for possible health impacts using the handheld multi-gas analyser, Model-VEP-200 monitor found the presence of Methane (CH₄) and NO₂ gases which is within the allowable limits.

Weli and Adekunle (2014), examined the air quality in Rumuolumeni neighbourhood landfill in Port Harcourt to determining the possible influence of weather parameters on the concentration of air contaminants from the landfill. Air quality parameters examined from the study include nitrogen oxide, sulphur dioxide, methane, volatile organic compounds, ammonia (NH₃), and hydrogen sulphide (H₂S). The result reveals higher concentration of CH₄ (0.06 mg/m³), SO₂ (0.67 mg/m³), VOC (2.28mg/m³), H₂S (0.19 mg/m³), and NH₃ (0.12 mg/m³) except NO₂ at the dumpsite. Result reveal that relative dampness and temperature sways the concentrations of NO₂, NH₃ and H₂S. Similarly, Rim-Rukeh (2014) assessed the effect of municipal solid waste dump sites fire on atmospheric pollution in five (5) dump sites using hand-held air quality monitoring equipment. The results revealed a very high concentration of SPM (773 - 801 µg/m³), CO (133.7 -141.6ppm), CO₂ (401 - 404.5 ppm), NO₂ (21.0 - 27.3 ppm), SO₂ (27.7 - 37.1 ppm), NH₃ (14.7 - 19.5 ppm), Methane (2310 - 2771 ppm), and H₂S concentration (3.4 to 7.7 ppm) within the area of the dump site. Ubouh and Nwawuike (2016), evaluate the on-site and off-site ambient air quality at Nekede dumpsite in Owerri at a distance of 300 meters using the Crowcon Portable Gas Analyzer. The result confirms a high concentration of CH₄, NH₃, NO₂, SO₂ and Particulate Matter from the waste dumpsite except CO.

METHODOLOGY

Study Area

Makurdi is the administrative capital of Benue State, located on Latitude 7° 45' 50" N and Longitude 8° 32' 10" E, on the Benue River in the Niger-Benue trough (Figures 1). Makurdi urban area covers a landmass of approximately 16km² with a population of about 393,425 people (National Population Commission projected population 2020). Makurdi is growing at a rapid rate with the presence of the Federal institutions including Federal University of Agriculture, 72 Special Airborne Battalion Barrack, NASME Barrack, Nigerian Air Force Base, the Nigerian Navy Provost School and Benue State University, has been a pull force to its rapid urbanization in the town. The predominant occupation of the indigenous people is agriculture which ranges from crop cultivation at subsistence

and commercial scales, fishing which takes advantage of the location of the Benue River, and

small-scale animal husbandry, with others ranging from white collar-jobs to handcrafts and trading.




Figure 1: Location of Makurdi

Data Collection and Analysis

Data were collected on certain components of emission concentrates of pollutants from five (5) dumpsite within the neighbourhoods with the aid of handheld MSA Altair 5X Multi-gas Detector manufactured by Mine Safety Appliance Company, USA. The gas analyser device is equipped with four sensors that monitor gases in ambient air and display readings. The device has a dual sensor that provide readings for CO and H₂S, an oxygen sensor (it was not functional), pentane and methane. Three (3) air quality monitoring parameters - carbon monoxide (CO), Hydrogen sulphide (H₂S) and methane (CH₄) were determined. The level of concentrates

measured in the neighbourhoods were compared with regulatory Guidelines and standards for Environmental Pollution Control in Nigeria by the Federal Environmental Protection Agency (FEPA) in 1991 stipulates emission limits of pollutants (Table 1) from stationary sources (site, process, stack and vent), and the WHO standards for air quality parameters monitoring (Rim-Rukeh, A. 2014). The guidelines prescribe safe levels of common air pollutants like suspended particulate matter (SPM), oxides of carbon, nitrogen and sulphur, volatile organic compounds (VOC) and hydrogen sulphide. Measures above the stipulated regulation is considered highly polluted and harmful to the inhabitants.

Table 1 Nigerian Ambient Air Quality Standard

Air Pollutants	Emission Limits
Particulates	250 (µg/m ³)
SO ₂	0.1 (ppm)
Non-methane	160 (µg/m ³)
CO	11- 4 (µg/m ³) or 10 (ppm)
NOX	0.04 - 0.06 (ppm)
Photochemical Oxidant	0.06 (ppm)
H ₂ S	10 (ppm)
CH ₄	1000 (ppm)

Source: FMEnv, 1991

RESULTS AND DISCUSSION

Ambient Air Quality Municipal Solid Waste Management Site

The concentrations of suspended particulate in ambient air sampled at the vicinity of the solid waste

dump site in residential neighbourhoods in Makurdi metropolis is presented in Figure 2. The ambient air quality sampled shows the presence of a high concentration of toxic gases including carbon monoxide (CO), hydrogen sulphide (H₂S) and methane (CH₄) (Akpofure, 2014).




Figure 2: Mean Air Quality at Residential Neighbourhoods Solid Waste the Dump Sites

At the neighbourhoods' level, the mean concentration of carbon monoxide (CO) gas recorded at six different dumpsites in Wadata is 16.0ppm, Wurukum neighbourhood (8.17ppm), North Bank (12.16ppm), while both High-Level and Modern Market neighbourhoods recorded 2.64ppm of carbon monoxide in each. High CO concentration is observed in waste dump site at Wadata and North Bank neighbourhood areas (Figure 1) above the FMEnv limit of 10ppm and WHO limit of 10 – 20 ppm. At extremely high levels, CO can cause death (Kinoshita, 2020). Also, a high concentration of carbon dioxide which is a by-product of methane in the atmosphere causes the depletion of the ozone layer resulting in global warming.

Similarly, the mean concentration of hydrogen Sulphide (H₂S) gas recorded in the vicinity of solid waste dump site in the neighbourhoods shows the presence of a high concentration of H₂S (M = 10.51ppm) in Wadata. Whereas, low concentration of H₂S was recorded in Wurukum (M = 5.67ppm), North Bank (M = 4.17ppm), and Modern Market (M = 1.74ppm). There is no trace of H₂S at the refuse dump site in the High-Level neighbourhood. Hydrogen Sulphide (H₂S) is known to be immediately dangerous to life and health (IDLH). It

has a pungent smell which produces a very strong rotten-egg smell even at very low concentration, but at a high concentration, the odour will no longer be detected by the human nose. The health effect of H₂S in the environment includes eye, throat, and lung irritation, nausea, headache, nasal blockage, sleeping difficulties, weight loss, chest pain and aggravation of asthma (Mir, & Maurya, (2020). Humans are extremely sensitive to hydrogen sulphide odours and can smell such odours at concentrations as low as 0.5 to 1ppm (Akpofure, 2014). Hydrogen sulphide has both natural and man-made sources (such as biodegradable waste sites). Hydrogen sulphide does not have regulatory limits, because it is a “non-criteria” pollutant.

Also, the high concentration of Methane gas was recorded at the refuse dump site in Wadata (M = 12.17ppm), Wurukum (M = 3.00ppm), North Bank (M = 4.16ppm), Modern Market (M = 1.14ppm), and no trace of CH₄ at High-Level neighbourhood. Methane is an odourless gas and is lighter than air. Because of its lightness, it tends to rise and accumulate near the higher, stagnant parts of enclosed buildings and tightly closed manure storage pits, it is a major greenhouse gas (Akpofure, 2014).




Plate 1: Typical Solid Waste dump in the Neighbourhood in Makurdi

CONCLUSION

The study reveals that the CO and H₂S concentrations in the ambient air were found to exceed the FMenv and WHO regulatory recommended standards. Air quality within the neighbourhoods constitutes pollutants - Carbon monoxide (CO), and hydrogen Sulphide (H₂S), concentration from the solid waste dumpsites. Residents are at the risk pollutants emission from the solid waste dumpsites which threaten health due to exposure through inhalation of air that can cause respiratory disorders and lung cancer in humans. Wadata and North Bank are densely-populated neighbourhoods in Makurdi. The quality of ambient air is an important concern of habitants living in proximity to the dump sites. There is a need, therefore, to develop better practices in the management of solid waste, especially the open dump sites operation and emission control. Also, the need to constantly monitor the ambient air quality in Makurdi, especially around the dumping sites to avoid harmful air pollution for the resident's safety.

References

- Aguoru, C.U. & Alu, C.A. (2015). Studies on Solid Waste Disposal and Management Methods in Makurdi and its Environs North Central Nigeria. *Greener Journal of Environmental Management and Public Safety*, 4(2):019-027.
- Amoo, O.M. & Fagbenle, R.L., (2013). Renewable municipal solid waste pathways for energy generation and sustainable development in the Nigerian context. *Int. Journal of Energy Environ. Eng.* 4, 42e59.
- Babel, S. & Vilaysouk, X. (2015). Greenhouse gas emissions from municipal solid waste management in Vientiane, Lao PDR. *Waste Management & Research*. doi:10.1177/0734242x15615425
- Chalvatzaki, E. & Lazaridis, M. (2010). Estimation of greenhouse gas emissions from landfills: application to the Akrotiri landfill site (Chania, Greece). *Global NEST Journal*, 12(1), 108-116.
- Chen, C., Kotyk, J. F. K. & Sheehan, S. W. (2018). Progress toward commercial application of electrochemical carbon dioxide reduction. *Chem*, 4(11), 2571-2586.
- Chen, Y.C. & Lo, S.L., (2015). Evaluation of greenhouse gas emissions for several municipal solid waste management strategies, *Journal of Cleaner Production*, 1-7.
- Du, M., Peng, C., Wang, X., Chen, H., Wang, M., & Zhu, Q. (2017). Quantification of methane emissions from municipal solid waste landfills in China during the past decade. *Renewable and Sustainable Energy Reviews*, 78, 272-279.
- Getahun T, Mengistie E, Haddis A, Wasie F, Alemayehu E, Dadi D, Van Gerven T. & Van der Bruggen B. (2012). Municipal solid waste generation in growing urban areas in Africa: current practices and relation to socioeconomic factors in Jimma, Ethiopia. *Environmental Monitoring & Assessment*, 184(10), 6337-45.
- Hoornweg, D. & Bhada-Tata, P. (2012). What a waste: A global review of solid waste management. Urban Development and Local Government Unit. World Bank, Washington, DC. Available at: <https://goo.gl/WyZiK5> (accessed 2/04/2020)
- Kinoshita, H., Türkan, H., Vucinic, S., Naqvi, S., Bedair, R., Rezaee, R., & Tsatsakis, A. (2020). Carbon monoxide poisoning. *Toxicology reports*, 7, 169-173.
- Le Treut, H., Somerville, R., Cubasch, U., Ding, Y., Mauritzen, C., Mokssit, A., Peterson, T. & Prather, M. (2007). Historical Overview of Climate Change. In: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*
- Maria, C., Gois, J, & Leitao, A. (2019). Challenges and perspectives of greenhouse gases emissions from municipal solid waste management in Angola. *Energy Reports*, <https://doi.org/10.1016/j.egy.2019.08.074>.
- Metz, D. (2015). Peak car in the big city: reducing London's transport greenhouse gas emissions. *Case Studies on Transport Policy*, 3(4), 367-371.
- Mir, J. & Maurya, R. (2020). The primary gasotransmitters and their respective donors in the mission of vision (eye health): a comprehensive overview. *Authorea Preprints*.
- Niloufer, S., Swamy, A. V. V. S. & Syamala, D. K. (2014). Gaseous emissions from MSW dumpsites in Vijayawada. *AIJRSTEM*, 6, 67-73.
- Okiongbo, K. S. & Akpofure, E. (2014). Identification of hydrogeochemical processes in groundwater using major ion chemistry: a case study of Yenagoa and environs, southern Nigeria. *Global Journal of Geological Sciences*, 12, 39-52.
- Ramachandra, T. V., Bharath, H. A., Kulkarni, G. & Han, S. S. (2018). Municipal solid waste: Generation, composition and GHG emissions in Bangalore, India. *Renewable and Sustainable Energy Reviews*, 82, 1122-1136.
- Stocker, T.F. (2014). *Climate change 2013: the physical science basis: working Group I contribution to the Fifth assessment report of the Intergovernmental Panel on Climate Change*. Cambridge University Press.

- Ubouh, E. A., Nwawuiké, N. & Ikwa, L. (2016). Evaluation of the on-site and off-site ambient air quality (AAQ) at Nekede waste dumpsite, Imo state, Nigeria. *British Journal of Earth Sciences Research*, 4(1), 18-22.
- Verma, N., Kaur, M. & Tripathi A. K., (2020). Greenhouse Gas Emissions from Municipal Solid Waste Management Practice. in V. Shukla, N. Kumar (eds.), *Environmental Concerns and Sustainable Development*, Springer Nature Singapore. https://doi.org/10.1007/978-981-13-6358-0_17
- Weli, V. E. & Adekunle, O. (2014). Air quality in the vicinity of a landfill site in Rumuolumeni, Port Harcourt, Nigeria. *Journal of Environment and Earth Science*, 4(10), 1-9.
- Ying-Chu, C. (2018). Effects of urbanization on municipal solid waste composition. *Waste Management* 79 (2018) 828–836. <https://doi.org/10.1016/j.wasman.2018.04.017>